# MYCOLOGIA

OFFICIAL ORGAN OF THE MYCOGOGICAL SOCIETY OF AMERICA

Vol. XL

NOVEMBER-DECEMBER, 1948

No. 6

## CONTENTS

Notes on certain Gasteromycetes, including two new orders	
S. M. Zeller	639
Studies in the dark-spored agarics Alexander H. Smith	669
An undescribed species of Helminthosporium on sudan grass	
and sorghumC. L. LEFEBVRE AND HELEN S. SHERWIN	708
Two unusual fungi from Glacier National Park, Montana	
E. B. Mains	717
Six new indian Discomycetes	724
A study of the temperature and humidity requirements of	
Aspergillus niger	728
Two species of Coprinus with notes on their cultural characters	
Morten Lange	739
Mycological notes. IX	748

## PUBLISHED BIMONTHLY FOR THE NEW YORK BOTANICAL GARDEN

AT PRINCE AND LEMON STS., LANCASTER, PA.

Entered as second-class matter April 30, 1925, at the post office at Lancaster, Pa., under the Act of August 24, 1912,

## MYCOLOGIA

Published by

THE NEW YORK BOTANICAL GARDEN

MYCOLOGICAL SOCIETY OF AMERICA

OFFICERS OF THE
MYCOLOGICAL SOCIETY OF AMERICA

JULIAN H. MILLER,

President,
University of Georgia

F. K. SPARROW, JR.,

Vice-President,
University of Michigan

GEORGE B. CUMMINS, '48

Councilor,

Purdue University

JOHN A. STEVENSON, '49

Councilor,
U. S. Bureau of Plant Industry

J. C. GILMAN,

Secretary-Treasurer,

Iowa State College

A. J. MIX, '48

Councilor,

University of Kansas

RALPH EMERSON, '49

Councilor,
University of California

HARRY M. FITZPATRICK,

Historian,

Cornell University

## EDITORIAL BOARD

ALEXANDER H. SMITH,

Editor-in-Chief,
University of Michigan

NORMAN F. CONANT, '51
Duke University

WILLIAM W. RAY, '48 University of Nebraska DONALD P. ROGERS,

Managing-Editor,

The New York Botanical Garden

BERNARD O. DODGE, '49
The New York Botanical Garden

STUART M. PADY, '52 McGill University

EDITH K. CASH, '50 U. S. Bureau of Plant Industry easurer,

nia

ditor, al Garden , '49 al Garden

2



## **MYCOLOGIA**

OFFICIAL ORGAN OF THE MYCOLOGICAL SOCIETY OF AMERICA

Vol. XL November-December, 1948

No. 6

## NOTES ON CERTAIN GASTEROMYCETES, INCLUDING TWO NEW ORDERS 1

S. M. ZELLER \*

These notes have for the most part grown out of the study of type and other material at the New York Botanical Garden <sup>2</sup> and a certain few other types loaned from elsewhere.

## HYMENOGASTRALES

### 1. Gasterellaceae fam. nov.

Fructificationes parvissimae, depresso-globosae, epigeae; gleba uniloculata; sporis brunneis, verrucosis.

Fructifications very tiny, depressed globose, epigeous; campanulate development; gleba finally uniloculate, but at times with one circle of cavities formed by vertical, centripetal invaginations which reach the center forming a false columella; cavities lined with a basidial hymenium; spores verrucose, dark.

Type genus: Gasterella Zeller and Walker.

This family is erected to take the two uniloculate genera, Gasterella and Gasterellopsis, while Protogaster with coralloid development and smooth spores is retained in Protogasteraceae.

<sup>1</sup> Published as Technical Paper No. 535 with the approval of the Director of the Oregon Agricultural Experiment Station. Contribution from the Department of Botany and New York Botanical Garden, Cryptogamic Herbarium, co-operating.

<sup>2</sup> Reference to the New York Botanical Garden Herbarium hereafter in this paper will be by initials only, N.Y.B.G.

\* News has just been received that Dr. Zeller died from coronary thrombosis Nov. 4, 1948.

2. CORDITUBERA Hennings in Engler, Bot. Jahrbücher. 23: 557–558. f. V-8. 1897. [Syn. Hoehneliogaster Lohwag, Beih. z. Bot. Centralbl. 42 (Abt. 2): 299. 1926.]

We can see no particular reason for the genus *Hoehneliogaster* and thus it is being referred back to synonymy with *Corditubera* whence it came. The differences are more specific than generic. In *C. microspora* upon which Lohwag based his new genus, the basidia-containing lacunae are not quite filled with the netted reticulum of basidia-bearing hyphae, so that toward their centers are left slight cavities, whereas in *C. Staudtii* the lacunae are completely filled. Although the gleba is reddish in one species and yellowish in the other, the genus *Corditubera* is more closely related to *Leucogaster* than to any other genus.

3. The description of Hydnangium Wallroth given by Cunningham <sup>3</sup> as emended by Ed. Fischer (Nat. Pflanzen-familien **7a**: 30–31. 1933) is very misleading, especially so where he indicates the "columella dendroid, arising from a well-developed, sterile base," whereas the original description by Wallroth definitely states that a sterile base is not present and makes no mention of a columella. Fischer's description too was misconstrued and erroneously stated; thus Cunningham thought he might have reason to relegate Cavara's Arcangeliella to synonymy with Hydnangium. The facts are that Fischer presented informal taxonomic descriptions and actually pointed out a contrast between Hydnangium and Arcangeliella; that whereas in a species like Hydnangium carneum, the type of the genus, in which its primordial stages only have a columella, Arcangeliella retains the columella and pileate peridium to maturity.

After having collected and examined in the field and in the laboratory many specimens in both of these groups we are in agreement with Fischer on the relationships between Arcangeliella and Hydnangium and will abide by the published treatment of these two genera and Octaviania.<sup>4</sup>

<sup>&</sup>lt;sup>8</sup> Cunningham, G. H. The Gasteromycetes of Australia and New Zealand. See p. 63. 1942.

<sup>4</sup> Mo. Bot. Gard. Ann. 23: 574-575; 602-605. 1936.

## 4. Hydnangium nigrescens sp. nov.

Fructificationes  $2.5\text{--}4\times1\text{--}3$  cm. crassae, oblongae vel subglobosae, rhizomorpho affixae; superficie glabra, molli, alba dein nigrescente; fibrillis paucis, ad superficiem inferiorem applanatis; peridio simplici,  $130\text{--}150~\mu$  crasso, siccato  $45\text{--}60~\mu$  crasso, hyphis hyalinis magnis circa  $7.5\text{--}9~\mu$  crassis implicatis composito, superficie fuscata; gleba brunnea; locellis medii-magnitudinis; septis tenuibus, albis, hyphis compactis parallelibus compositis, circa  $60\text{--}85~\mu$  crassis (hymenio annumerato); basidiis 2-sporigeris; sporis subglobosis,  $12\text{--}16~\mu$ , pedicellatis, obscure brunneis, unigutulatis, episporio crasso, verrucoso, verrucis magnis, 10--12 circumferentiam quamquam ornantibus.

Fructifications oblong to subspherical, about 2.5–4 cm.  $\times$  1–3 cm., attached by a single rhizomorph; surface soft, glabrous, with some innate veins especially below, white becoming black on oxidizing, drying fuscous black; peridium 130–150  $\mu$  thick when fresh, drying 45–60  $\mu$  thick, spongy, composed of large loosely woven, hyaline hyphae, about 7.5–9  $\mu$  in diameter, the surface hyphae dark, forming a slightly more compact rind; gleba brown, drying Prout's brown; cavities relatively large; septa white, drying papery thin, of compact, parallel hyphae, about 60–85  $\mu$  thick (including hymenia); basidia 2-spored; spores subglobose to slightly ellipsoid, sometimes with remains of the stout sterigma, 12–16  $\mu$  in diameter, dark brown with one large vacuole, epispore thick, rough with very large blunt verrucae, 10–12 per circumference.

Partially exposed on the ground in mixed woods. **Type** collected in Cornell Plantations along Fall Creek, east of Floriculture Gardens, Tompkins county, New York, C. T. Rogerson, No. 1615, August 18, 1947. (In Cornell University, Plant Pathology Herb. No. 37257, and also in Zeller Herb.)

H. nigrescens has spore markings like those of H. purpureum but the spores are larger. It also differs in peridial characters, especially color, thickness, and size of hyphae.

5. **Hydnangium vesiculosum** (Coker & Couch) n. comb. (Syn. *Gymnomyces vesiculosus* Coker & Couch, Gasteromycetes of Eastern U. S. and Canada, p. 23. 1928.)

Fructifications subspherical, about 1 cm. diameter, drying to about half size, attached by a few basal fibrils which spring from a narrow depression; surface light buff yellow (about Naples yellow), spongy; peridium at first thin, of delicate, loosely woven, yellowish hyphae, quite evanescent, nearly absent at maturity, ex-

aster
ubera
neric.
, the

557-

h. z.

etted nters comand

re-

30s the

that ella. ted; gate The

and um,

the in

igenent

and.

posing the gleba in places; gleba pallid gray-brown (pale earthy buff), firm but pliable, not tough or elastic; cavities rather large and box-like, empty, about 0.4–0.8 mm. in diameter; septa about 0.17–0.2 mm. thick including the hymenia which are about 30  $\mu$  thick, composed of large bladdery cells (pseudoparenchyma) about 20–40  $\mu$  thick except for a very thin layer under the hymenium where the cells are small; basidia short, thick, 4-spored; sterigmata nearly half as long as the spore diameter; spores concolorous with gleba, spherical, 7.5–10  $\mu$  including the strong blunt spines, often stuck together in groups of 4, sometimes finely reticulate.

Type locality: Chapel Hill, North Carolina.

Habitat: Exposed by erosion on soil in frondose woods.

Distribution: Known from type locality only.

Illustrations: Coker & Couch, Gasteromycetes of Eastern U. S. and Canada. pls. 16, 17, 105, f. 17-19.

As stated previously there is no sound basis for retaining Gymnomyces since all the species have a peridium, even though it may be evanescent. The same argument holds for the retention of Chamonixia as separate from Gautieria since most, if not all, species of the latter have an evanescent peridium.

The species Gymnomyces vesiculosus is, therefore, transferred to Hydnangium. The spores are sculptured as described by Coker & Couch but they also have very fine reticulations which seem not to be at all raised (alveolate).

6. Elasmomyces Rodwayi (Massee) n. comb. . (Syn. Secotium Rodwayi Massee, Kew Bull. Misc. Info. 1901: 158. 1901.)

A part of the type collection from the Herbarium of George Massee is at the N.Y.B.G. The spores are subglobose, distinctly echinulate with blunt spines and also reticulate with large meshes, practically hyaline, short pedicellate, and  $9{\text -}11~\mu$  in diameter.

7. MacOwanites Kalchbrenner, Hedwigia 15: 115. 1876; Grevillea 10: 107. 1882. [Macowania Kalchbrenner, Gardeners' Chron., N.S. 5: 785. 1876, non Macowania Oliver in Hooker, Icon Pl. III 1: 49. 1870.—Hypochanum Kalchbrenner, Gard. Chron., N.S. 6: 140. 1876. nomen nudum.]

As was indicated previously 5 the spores of MacOwanites agaricinus Kalchbr. are borne "asymmetrically on the sterigmata as in

<sup>&</sup>lt;sup>5</sup> Zeller, S. M., and C. W. Dodge, Ann. Mo. Bot. Gard. 23: 636-637. 1936.

the Hymenomycetes." This matter has just been studied again and the part of the type in the herbarium of the N.Y.B.G. proves to be part of an aberrant Agaric belonging perhaps to *Russula*. The spores are slightly ellipsoid, echinulate, pedicellate with the sterigmatal scar asymmetrical to the main axis of the spore. There is also pseudoparenchyma in the tramal tissues as well as in the pileate tissue. According to nomenclatural procedure the genus is rejected and other species which have been assigned to this genus are disposed of as follows:

MacOwanites magnus Parks has lactiferous ducts in the sterile tissues and is like Arcangeliella in general morphological development. It, therefore, becomes Arcangeliella magna (Parks) n. comb. and M. alpinus Zeller is recombined as Elasmomyces alpinus n. comb.

8. GYMNOGLOSSUM Massee, Grevillea 19: 97. 1891. (Syn. Dendrogaster Bucholtz, Hedwigia 40: 316-318. 1901.)

Study of a part of the type collection of Gymnoglossum stipitatum Massee in the herbarium of N.Y.B.G. reveals that Cunningham's reduction of Dendrogaster to synonymy is justified. The specimen examined has smooth spores, however, as Massee reported. They are  $13-16\times 6-8.5\,\mu$ , definitely citriform with a very small apiculus and sometimes with a short pedicel. Since Dendrogaster is thus reduced to synonymy the following new combinations are necessary, Gymnoglossum majus (Z. & D.), G. megasporum (Z. & D.), G. cambodgense (Patouillard), G. candidum (Harkness), G. radiatum (Lloyd), G. connectens (Bucholtz), G. utriculatum (Harkness), G. globosum (Harkness), G. foetidum (Coker & Couch), G. olivaceum Zeller, and G. elasmomycetoides Zeller.

## HYSTERANGIALES

1. PROTUBERA Möller, in Schimper, Botanische Mitteilungen aus den Tropen 7: 10–22: 145. T VI, f. 1–10. 1895. (Syn. Protophallus Murrill, Mycologia 2: 25. 1910.)

There are two dried collections of Protubera Maracuja Möller in the Cryptogamic Herbarium of the N.Y.B.G., both ex Farlow

30 µ about nium erigorous oines, te.

arthy large

about

J. S.

may n of all,

erred d by which

tium

nctly shes,

876; ners' oker, Gard.

*gari*as in

1936.

<sup>&</sup>lt;sup>8</sup> l.c. See p. 71.

Herb., Harvard University and both collected by Rev. I. Rick as follows: Rick Fungi Austro-Americani, No. 36, taken at Sao Leopoldo, 1904, and Rick Expeditions in Brazil, No. 136, taken at Parecy Novo, Rio Grande do Sul, 1928. Both collections answer very closely Möller's description of P. Maracuja. After the fructifications were soaked in water for 10-12 hours the character and organization of the essential parts were readily observed. There were found but two minor differences between Protubera and Protophallus Murrill. The one is hypogeous, the latter epigeous, and the former has a columella branched from the base or point of attachment. In the two species of Protophallus the columella is simple, extending to the center of the fructification and from its summit the tramal plates radiate. Such differences are more specific than of generic rank and it would seem wise to unite the two genera and transfer the two species of Protophallus to Protubera as Protubera jamaicensis (Murrill) n. comb. and P. brunnea Zeller n. comb.

P. Maracuja and P. brunnea are quite closely related but they differ in color, and characters of columella and spores. Protubera and Calvarula will be retained as members of the family Proto-phallaceae.

### **PHALLALES**

1. CLATHRUS Persoon, Syn. Meth. Fung. p. 241. 1801. (Dycticia Rafinesque in Desraux, Jour. Bot. 2: 176. 1809.—Clathrus sect. Clethria Fries, Syst. Myc. 2: 288. 1823.—Ileodictyon Tulasne, Ann. Sci. Nat. Ser. III 2: 114. 1844.—Clathrella Fischer, in Engler & Prantl, Nat. Pflanzenfam. 1: [Ab. 1\*\*] 284. 1898.)

It is with some hesitancy that Clathrella Fischer is placed in synonymy with Clathrus, but there seems to be only one real distinction besides size to separate them, namely the longer meshes on the sides compared with those over the top of the receptacle of Clathrella while in Clathrus these meshes are more or less the same size throughout. Also the drawn-out stem-like base in some individual plants of Clathrella is not a constant character and may be as conspicuous in some individuals of Clathrella. It may or may not be present in all plants of a single collection according to Cunningham.<sup>8</sup>

c as

Sao

ken

an-

the

cter

ved.

epi-

or

the

tion

ices

vise

oto-

n.

hey

era

oto-

vc-

rus

Γu-

ner,

8.)

in

real

hes

of

the

me

nay

als

fa

Ileodictyon was proposed by Tulasne to receive those species of Clathrus with tubular, not chambered, arms of the receptaculum. Here again, Cunningham has found that "the type species, Clathrus cibarius, contains plants with both tubular and cellular arms, small plants as a rule possessing tubular, and large ones, cellular arms." Cunningham is, therefore, followed in grouping species of Clathrella and Ileodictyon under the genus Clathrus.

2. Lysurus Fries, Systema Mycologicum 2: 285–286. 1823. (Anthurus Kalchbrenner & MacOwan in Kalchbrenner & Cooke, Australian Fungi, Grevillea 9: 2. 1880.—Aserophallus Lep. & Mont. Ann. Sci. Nat. [Ser. 3] 4: 360. 1845.—Mycopharus Petch, Brit. Myc. Soc. Trans. 10: 281. 1926.—Pharus Petch, Bot. Gard. Peradeniya, Ann. 7: 59. 1919.—Lysurus sect. Desmaturus Schlechtendal in Linnaea 31: 180. 1861–62.)

Peridium duplex, outer layer thin and furfuraceous, inner thick, gelatinous, remaining at the base of the stem as a volva at maturity; receptacle a hollow cylindrical or flaring stem, carrying at its summit a number of simple arms apically free or united organically or by a delicate membrane, continuous with or somewhat distinct from the stem below; gleba borne on the inner surfaces and sides of the arms, olivaceous, mucilaginous, foetid; spores smooth, phalloid.

Type species: Lysurus Mokusin (Pers.) Fries.

The similarity between Lysurus and Anthurus has been discussed by Cunningham.<sup>3</sup> Early workers as well as workers of today have not been certain of the taxonomy of these two genera, and since the type specimen upon which Kalchbrenner & MacOwan erected the genus Anthurus no longer exists and the original description of Anthurus could just as well apply to Lysurus, Anthurus is reduced to synonymy. We cannot agree with Cunningham that Pseudocolus Lloyd is synonymous with Anthurus or Lysurus. It stands apart as a genus in which the arms of the receptacle are long and narrow and united at the apex.

The species that have been assigned either to Anthurus or Lysurus have at one time or another usually been referred to both genera so there are few if any new combinations to be made by the union of the two under the name Lysurus. There are three spe-

<sup>&</sup>lt;sup>8</sup> l.c. See pp. 101 and 104.

ce

gl

ti

5.

d

g

tl

cies in North America, namely, Lysurus Gardneri Berkeley, London Jour. Bot. 5: 535. 1846, Lysurus Mokusin (L. ex Pers.) Fries, Syst. Myc. 2: 286. 1823, and Lysurus pusillus Coker, Mycologia 37: 781–783. 1945.

3. Kupsura sphaerocephala Lloyd, in Lloyd, Myc. Writ. 7: 1303. f. 2903–2904. Oct. 1924.

The type specimen (No. 24893) (kindly loaned by Dr. J. A. Stevenson), upon which Lloyd based the genus Kupsura, proves to be a very mature, softened specimen of Simblum sphaerocephalum Schlechtendal. When the specimen was fresh the stem of the receptacle had been crowded up into the head and it was dried in that condition. This was misleading to Lloyd who took the brown interior (receptacle) of the head to be brown glebal tissue. Misleading also is a portion of the volva which remained over the top of the latticed head. Kupsura Lloyd thus becomes a synonym of Simblum.

4. STAHELIOMYCES CINCTUS Ed. Fischer, Mitteil. Nat. Ges. in Bern 1920: XXXV and 137. 1921.

This genus has been reported previously from British and Dutch Guiana. There is a very good pencil sketch made by A. F. Porter, Decorah, Iowa, of a specimen about 9 inches tall and 1.5 inches broad which he found in a dense jungle forest on the Bobanaza River, near Saryacu [Curicucha], Eastern Ecuador. The sketch is in the N.Y.B.G. Herb. This extends the known range fairly well across northern South America. There is no mistaking the identity of the fungus from which the sketch was made.

Somewhat more difficult, however, is the identification of the fungus from which was made a water-color painting to be found in the Ellis Collection at the same herbarium. Under the sketch in Ellis' handwriting is "Corynites Ravenelii B. & C. from Miss Lizzie Berk Hancock, Burlington, N. J., Autumn, 1881." The illustration is of a volvate fungus with reddish receptacle like a Mutinus (Syn. Corynites) but with the gleba in a zone around the receptacle some space below the apex, the tip of which had evidently been broken off. There was not the usual constriction of the receptacle at the glebal band as in S. cinctus. The plant evidently had all of the characters of Staheliomyces but if the re-

ceptacle was red and there was no constriction of the latter at the glebal zone it may have been a distinct species. There is a question, therefore, as to the species; and was the painting made from a specimen actually collected in New Jersey?

5. Itajahya galericulata Möller, in Schimper, Bot. Mitt. aus den Tropen 7: 79. pl. 5. 1895.

Our knowledge of the genus *Itajahya* in the United States is greatly enhanced by the careful field notes and compilations of the late Dr. W. H. Long <sup>6</sup> and his co-operators. He has reported the genus from Texas, New Mexico, and Arizona. In the herbarium of the N.Y.B.G. is a specimen of *Itajahya galericulata* collected by E. Bethel near Denver, Colorado. There are no further collecting data with the specimen.

#### LYCOPERDALES

1. The Lycoperdaceae of the past has been more or less a catchall for any gasteromycete having capillitium and so it seems best to divide the family as presented by Fischer (1933) into more unified entities. The genera with stromate fructifications have been referred to the new family Broomeiaceae and those with spiny capillitium are included in the new family Mycenastraceae.

#### Broomeiaceae fam. nov.

Lon-

ers.)

oker,

t. 7:

. A.

oves

pha-

f the

d in

the

sue.

nym

. in

utch

rter.

ches

irly

the

the

und

etch

**Iiss** 

The

e a

and

had

ion

ant

re-

Fructificationes stromatae ovoidae vel subglobosae; peridio duplici, exoperidio tenui subevanescenti, endoperidio membranaceo, stomatibus; capillitio simplici.

Fructifications single or many on a stroma, mostly ovoid, hemispheric or subspheric; exoperidium thin, wholly or partly disintegrated at maturity, endoperidium papery or thickish, laid bare at maturity, opening by an apical pore; capillitium present, threads more or less symmetrical, simple.

This family was created to care for the three genera with stromate fructifications, *Broomeia*, with a stalked stroma, *Diplocystis*, with a resupinate or patellate stroma, and *Lycogalopsis* in which the fructifications are borne singly on a stroma. Fischer (1933)

<sup>&</sup>lt;sup>6</sup> Long, W. H., and D. J. Stouffer. The genus *Itajahya* in North America. Mycologia 35: 620-628. *Illus*. 1943.

questioned whether these genera should be included in the Lycoperdales at all, but I am referring the *Broomeiaceae* to this order.

## Mycenastraceae fam. nov.

Fructificationes magnae, subglobosae; peridio duplici, exoperidio crasso, spongioso, glabro vel areolato, endoperidio crasso coriaceoque vel tenui tenacique, demum irregulariter dehiscenti; capillitio ramoso, spinoso; sporis globosis vel ellipsoideis, verrucosis.

Fructifications large, subglobose; peridium duplex, exoperidium thick, spongy, smooth or areolate, endoperidium thick and leathery or thin and membranaceous; capillitium branched with short, pointed, spine-like branches; spores globose to ellipsoid, verrucose.

This family is proposed to contain the genera Mycenastrum and Calbovista.

 LYCOPERDON ALBINUM Cooke, in Massee, Jour. Roy. Microsc. Soc. 1887: 723. Oct. 1887.

Lloyd  $^{7}$  must have applied from memory Cooke's name of this fungus to Porto Rican collections made by John A. Stevenson and used the misnomer L. albidum instead of L albinum. A slide (No. 58690) in the Lloyd Herbarium prepared by Lloyd from the type of L. albinum Cooke has been studied and it appears that Mr. Lloyd was justified in referring the Porto Rican collections to this species. Lloyd and Cooke, however, made the same mistake; the spores are finely asperate instead of smooth. An emended description follows:

Fructifications 4–12 mm. in diameter, subspherical to pyriform, sessile, yellowish with a whitish bloom, smooth to slightly checkered by tiny cracks; peridium thin, somewhat farinose, yellowish; gleba yellowish then turning grayish; capillitium scanty, hyaline, slender, flaccid; spores spherical to somewhat irregular, slightly asperate, almost hyaline, 2.5– $4.2\,\mu$ .

Type locality: Brazil, South America.

Habitat: On rotted wood or on humus in soil.

Distribution: Puerto Rico and Brazil.

Illustrations: Lloyd, Myc. Writ. 5: 582. f. 822. 1916.

7 Lloyd, C. G., Myc. Writ. 5: 582. f. 822. 1916.

## 3. Coilomyces Schweinitzii Berk. & Curtis.

A study of the type of this genus and species, which is to be found in the Herbarium at N.Y.B.G., verifies Fischer's <sup>8</sup> assumption that this genus was based on a collection of *Geastrum mirabile* Mont. from Surinam. The central cavity to which the authors referred and upon which the generic name was based is where the relatively large columella had collapsed. The Berkeley and Curtis specific name predates Montagne's name and necessitates the following new combination:

## Geastrum Schweinitzii (Berk. & Curtis) Zeller n. comb.

Coilomyces Schweinitzii Berk. & Curtis, Jour. Acad. Nat. Sci. Philadelphia (Series 2) 2: 279-280. 1853.

Geastrum mirabile Mont., Ann. Sci. Nat., Ser. IV 3: 139-140. pl. 6, f. 8.

Geaster papyraceus B. & C., Am. Acad. Arts & Sci. Proc. 4: 124. 1858. Geaster lignicola Berk., Linn. Soc. Bot. Jour. 18: 386. 1891. Geaster caespitosus Lloyd, Myc. Writ. 2: 315. pl. 100. 1907.

## 4. Bovistella atrobrunnea sp. nov.

Fructificationes depresso-globosae vel turbinatae, superne saepe collapsae, 3–4 cm. crassae, 2–3 cm. altae, rhizomorphis affixae; superficie impolitae leves vel leniter furfuraceae vel leniter rimosae, siccitate obscure brunneae; exoperidio tenui, fragili, superne squamis facile separabilibus discedisque vestito; endoperidio tenuissimo papyraceoque, impolito- vel nitido-glabro, apice osculo laciniato dehiscente; basi sterili cellulosa, convex inscula, nitido-brunnea septis instructa; gleba pulverulenta obscure vinaceo-brunnea vel obscuriora; capillitio discreto, longo, paulo ramoso, terminalibus longis attenuatis tenui-oribusque, obscure brunneo; sporis brunneis, sphaericis, verrucosis, 5–8  $\mu$ , longe pedicellatis.

Fructifications' oblate spheroid to turbinate, often collapsed above, 3–4 cm. broad, 2–3 cm. high, with a prominent attachment but not particularly radicate; surface dull, smooth to somewhat furfuraceous, somewhat rimose, dark brown (dry); peridium duplex, exoperidium thin, brittle, breaking up into small plates which easily separate and fall away; endoperidium very thin, papery, dull to shiny, a little lighter colored than the exoperidium, dehiscing by a torn irregular apical pore; sterile base prominent, somewhat convex above, occupying one-third to one-half of the lower portion of the fructification, of large cells, separated by thin, shiny, metallic-brown walls; gleba pulverulent, dark vinaceous brown or darker; capillitium free, long, slightly branched

erosc.

and

Lyco-

order.

crasso, i tena-

sporis

dium

eath-

short.

icose.

and

slide n the that ctions mis-

An

form, heckwish; aline,

ghtly

<sup>&</sup>lt;sup>8</sup> Fischer, Ed. Gasteromycetes, in Engler and Prantl, Die Nat. Pflanzenfam. 7a: 1-122. 1933. (See p. 76.)

or simple, with long tapering narrow (or even thread-like) terminals, dark brown, somewhat uneven; spores brown, spherical, verrucose,  $5-8~\mu$ , with a hyaline pedicel up to  $30~\mu$  long which is easily broken away.

On the ground, Ann Arbor, Michigan, October 6, 1936, A. H. Smith, 5048a, type (in U. Mich. Herb., portion in Zeller Herb.).

Bovistella atrobrunnea differs from other species of the genus in the very dark gleba, the spherical, verrucose spores and particularly in the capillitium. The latter is nearly simple but, now and then, dichotomously branched and the branches or terminals are tapering or drawn out to very long, narrow thread-like filaments very much narrower than the main stem of each unit of capillitium. It is named for the very dark brown gleba.

## 5. Morganella gen. nov.

Fructificationes parvae, subglobosae; peridio duplici, apice ore dehiscente; gleba pulverulenta, capillitio tenuissimis membranisque ad basim radiantes; sporis sphaericis, coloratis.

Fructifications small, subglobose; peridium duplex, dehiscing by an apical stoma; gleba pulverulent, with capillitium and filmy membranes radiating from base to peridium; spores spherical, colored.

Type species Morganella mexicana.

## Morganella mexicana sp. nov.

Fructificationes parvae, subglobosae, solitariae vel caespitosae, appendici radiciformi albo adfixae; peridio duplici; exoperidio furfuraceo, brunneo, in squamas furfuraceas fisso; endoperidio papyraceo-tenui, apice ore irregulari dehiscenti; basi sterili inconspicua, contextu flavido solidiusculo sine locello composito; gleba pulverulenta, pallido-grisea vel pallido-brunnea, tenuissima membrana et capillitio sporisque composita; membranae hyalinae, ut bullatis sacculis instructis, atque notis lunulatis appareant; capillitio hyalino, laevi, paulo ramoso; sporis rubro-brunneis, sphaericis, paulo verrucosis, 3.7-4.3 µ.

Fructifications from a white, mycelial rhizomorph, small, sub-globose, single or caespitose; peridium duplex; exoperidium fur-furaceous, becoming separated over the endoperidium in the form of tiny furfuraceous squamules, bay or natal brown becoming hair brown; endoperidium thin, papery, dehiscing by an apical irregular rupture; gleba with a very slight sterile base which is of yellowish solid tissue without apparent chambers, fertile portion pulverulent, pale gray, becoming light chocolate brown, composed at maturity

of very filmy membranes (in which capillitium is enmeshed) radiating from the base to the endoperidium and spores; membranes hyaline, with tiny pockets and swellings appearing as crescent-shaped markings; capillitium threads (embedded in the filmy membranes) hyaline, smooth, very slightly branched; spores reddish brown, spherical, slightly verrucose,  $3.7-4.3\,\mu$ .

Type locality: Near Guaymas, Sonora, Mexico. Habitat: In wet places among moss and leaves.

ter-

ical.

h is

H.

b.).

enus

par-

now

nals

fila-

t of

nte;

ites;

by

emred.

ndici

, in

ılari ello

sima

latis

nevi, .3 μ.

ub-

urorm

nair

ılar

rish

ent.

rity

Distribution: New Jersey, Mexico, Panama, and Colombia.

Specimens examined: New Jersey: Newfield, Aug. 1887, J. B. Ellis, No. 5013 (in N.Y.B.G. Herb.); Mexico: Sonora, near Guaymas, Thomas H. Macbride, type (in Morgan Herb. at University of Iowa); Panama: Canal Zone, Ft. Sherman Area, G. W. Martin, No. 6183; and South America: Colombia, Sierra Nevada de Santa Marta, Dept. Magdalena, Hacienda Cincinnati, G. W. Martin, No. 3440 (both in Mycological Collections of the University of Iowa).

In the N.Y.B.G. Herbarium there is correspondence between Thomas H. Macbride, J. B. Ellis, and A. P. Morgan relative to the collection of this fungus taken by Macbride in Mexico and to which the collector had given the tentative name *Lycogala mexicanum*. The following letter from A. P. Morgan to Macbride seems worth quoting:

"That is a queer thing you sent me, but I think it is a puffball—Lycoperdon. I cut it in two and scanned it internally and externally. The peculiar membrane with disk-like markings certainly resembles the membrane of Myxomycetes, it is very thin and irridescent, but the spores are those of a puffball. I managed to get off some bits of the rind, which I consider to be made up of true hyphae; I enclose them to you. The threads of the capillitium are unlike those of ordinary puffballs; but there are two puffballs that have similar threads and they have bits of membrane adhering to them in the same way; these are Lycoperdon Curtisii Berk. and L. acuminatum Bosc. Of course the membrane is not as marked a feature as in this specimen.

"But what is more, in cutting down through the peridium I arrived at a slender rooting mycelium, a thick strand of hyphae in it. Tear this up or mark it with the point of a knife blade and

you can see the hyphae comprising it plainly with about 500 diameter.

"I am quite sure that it is a Lycoperdon, but it is nothing I have ever seen before."

This truly is a most interesting Gasteromycete which belongs in the Lycoperdaceae or the Mesophelliaceae. Until more is known of the origin and development of the membranes in the fertile gleba it is placed in the Lycoperdaceae for convenience. It seems possible that the membranes mentioned above may originate from a very thin tramal tissue between the hymenia of the septa. The origin of the pockets and swellings in the membranes is beyond our present understanding.

Morgan's mention of membranes in the gleba of Lycoperdon Curtisii Berk. and L. acuminatum Bosc. is unexplainable unless he was dealing with very young specimens in which the tramal tissues are just breaking down. Such membranes, however, seem unrelated to such as found in Morganella at maturity.

At late maturity of *Morganella* the whole gleba disappears, leaving a more or less cupulate or discoid, empty peridium, exposing the silvery gray inner surface of the endoperidium.

## 6. Radiigera cinnamomea sp. nov.

Fructificationes 1–3 cm. crassae, depresso-globosae vel turbinatae, basi saepe radicanti; superficie cinnamomea, squamis minutis, concoloribus, saepe apice conniventibus obsita; basis sterilis superne convex inscula, inferne attenuata, cellulis parvis composita; columella conica vel subglobosa, mollospongiosa compacta, concolor; peridio duplici 0.5 mm. siccato; exoperidio squamoso; endoperidio suberoso, fragili siccato, 250–300 µ crasso; gleba cinnamomea, fasciculis hypharum capillitioque sporisque composita; capillitio ad columellam percursum radians hyalino molli, flexuoso, inaequali, 6–7 µ crasso; sporis pallidissine-brunneis, sphaericis, sparse-echinulatis, pedicellatis, 3.75–5 µ crassis, non guttulatis.

Fructifications 1–3 cm. diam., depressed globose to turbinate, with a slightly projecting basal attachment; surface dull cinnamon brown, with minute indistinct, concolorous, sometimes connivent scales; sterile base very slightly convex above, somewhat attenuate below, cells small but distinct, crowned by a broad subspherical to conical columella which occupies about one-half of the central portion of the fructification; columella concolorous with the gleba, composed of a compact, soft, pithy, homogeneous (?) tissue; peridium duplex, total drying about 0.5 mm. thick, exoperidium scaly

as described above, endoperidium corky, brittle, about 250–300  $\mu$  thick; gleba cinnamon brown throughout, composed of fascicles of hyphae and capillitium radiating from the surface of the columella to the inner wall of the endoperidium, readily separating from the former on drying; capillitium hyaline, various sizes of filaments up to 6–7  $\mu$  in diam., soft flaccid, sometimes with very filmy remains of hyaline membranes adhering; spores very dilute brownish, almost hyaline, spherical, minutely and sparsely echinulate, 3.75–5  $\mu$ , with a long flexuous (collapsed) pedicel, not guttulate.

Type locality: Near Philadelphia, Pa., collected by  $T.\ G.\ Gentry$ , type (in N.Y.B.G. Herb. and portion in Zeller Herb.).

Distribution: Known from type locality only.

## **SCLERODERMATALES**

 Scleroderma aurea Massee in Cooke, Grevillea 18: 26. 1889.

One-half of a specimen in the N.Y.B.G. Herb. is labelled in George Massee's handwriting as follows: "Scleroderma aurea, Mass., New Guinea, TYPE." It is a small-spored form of S. aurantium Pers. The spores are globose, alveolate-reticulate-echinulate,  $6.3-7.5 \mu$ . The echinulae are quite long and acute.

2. Scleroderma columnare Berkeley & Broome, Ceylon Fungi No. 726.—Areolaria columnaris (Berk. & Br.) DeToni in Sacc. Syll. Fung. 7: 144–145. 1888.

A supposedly authentic specimen from Ceylon purchased from George Massee by the New York Botanical Garden is the short stipitate form of *Scleroderma verrucosum* Pers. Cunningham referred *S. columnare* to *S. Bovista* and Lloyd referred it to *S. cepa*. The latter is more nearly correct since the spores are not reticulate. The spores are bluntly echinulate, the peridium warted, and otherwise the specimen appears like *S. verrucosum*.

3. Scleroderma Lycoperdoides Schw. Schrift. Naturf. Ges. Leipzig 1: 61. 1822.—Scleroderma tenerum B. & C. Cuban Fungi No. 512.

The type collection of *Scleroderma lycoperdoides* Schweinitz is in the Ezra Michener Collection, Vol. 17, Sheet No. 37 (U.S.D.A., Bureau of Plant Industry, Mycological Collections, Beltsville,

erdon inless ramal seem

t 500

ing I

igs in

nown

gleba

pos-

om a

The

eyond

pears, i, ex-

saepe ne atmolloperidio pa cinpillitio 6-7 µ ellatis,

inate, amon iivent nuate erical entral gleba, peridscaly Maryland). The collection is labelled "No. 2242, Syn. Fung. Scleroderma lycoperdoides Schw., on the earth, Carolina (ex Herb. Schw.)." This specimen has spores with closely set acute spines and they average about 12– $14~\mu$ . It seems in every respect like S. tenerum B. & C.

4. Scleroderma tuberoideum Speg. Anal. Mus. Nac. Buenos Aires 16: 28. 1906.

An authentic specimen of this species sent by Spegazzini to the N.Y.B.G. proves to be S. cepa Pers.

5. Astraeus hygrometricus (Pers.) Morgan is a cosmopolitan species throughout temperate climates. The species is quite variable in surface characters of the exoperidium. There are smooth, even glossy specimens, others that are rough and felty, while still others are fibrillose scaly. Otherwise there seems to be little variability except in size.

One collection taken by O. E. Jennings, in sand dunes, Presque Isle, Erie county, Pennsylvania (in Herb. Carnegie Museum, Pittsburgh), is a very smooth form to which a great deal of study has been given and merits further attention. The peridium has the same structure as that of typical A. hygrometricus, but when soaked in water the exoperidium does not dehisce along well-marked sutures radiating from the apex. On the contrary it cracks along irregular lines. If other collections may be found to show the same tendencies this form could hardly be included in Astraeus.

6. Astraeus pteridis (Shear) comb. nov. (Scleroderma pteridis Shear, Bull. Torr. Bot. Club 29: 451. 1902.—Geastrum hygrometricum Pers. var. giganteum Lloyd, Myc. Writ. 1: 68. f. 30. 1901.)

Most of the years since 1909 I have spent in western Oregon or Washington and have wondered why Scleroderma pteridis Shear could not be found. Recently a portion of the type was discovered at N.Y.B.G. It proved immediately to be an old acquaintance, the "Giant Astraeus" of the Pacific northwest. It is not unusual to find the buttons up to 2 inches in diameter. The largest expanded star observed was over 9 inches. Shear men-

tioned its similarity to *Scleroderma geaster*. It is in size only. The peridial characters are entirely distinct.

## CALOSTOMA MICROSPORUM Atkinson, Jour. Myc. 9: 16. 1903. Emended description:

Fructifications 4–7 cm. high; foot stalk 3–6 cm.  $\times$  1–1.5 cm., cylindrical or ventricose or enlarged below, sometimes compressed, rarely two or more footstalks joined throughout the length; endoperidium ovoid, 10–15 mm. broad, slaty or bluish gray to warm brown, mouth stellate with 5–7 prominently raised teeth, vermillion colored on their inner faces; exoperidium separating into numerous small, hard, adherent warts covering the middle and lower surface of the endoperidium and usually entirely wanting toward the apex where the endoperidium is quite smooth; spore sac pure white; spores pure white, smooth, oblong, some rarely ellipsoid, 6–10  $\times$  3.5–5  $\mu$ . Type locality: Rugby, Tennessee.

The type collection was kindly loaned by Dr. H. M. Fitzpatrick, and Atkinson's description has been emended to conform with other descriptions presented here. There are to us three outstanding characters of the species: The small ellipsoid spores which show little variability in size and shape, the very prominently raised peristome, and the prominence of the exoperidium left as warts on the sides of the endoperidium. The spores and the spore sac are also characteristically pure white.

A collection by L. W. Nuttall taken from wet mossy banks in Fayette county, West Virginia, extends the range. This collection predates the collection of the *type* by nine years. It was distributed as No. 881, Flora of Fayette county, West Virginia, collected by L. W. Nuttall, March 25, 1893. In this collection the endoperidium is warm brown while in the type collection it is slaty or bluish gray.

### 8. Sedeculaceae fam. nov.

Fructificationes coriaceae, sine basi sterili nec radicibus; peridio crasso, superne coriaceo, inferne paene obsoleto dehiscentique; gleba pulverulenta, cum venis crassis, a peridio ad centrum vergentibus; sporis brunneis, breve pedicellatis.

Fructifications leathery, without sterile base or radicle; peridium thick, leathery above, almost obsolete and dehiscing below; gleba becoming powdery at maturity, with broad veins extending in-

o the

ung.

(ex

acute

spect

ienos

e still

seum, al of peridricus, along arry it and to ed in

pteriygrof. 30.

eridis e was n old t. It The ward from the peridium; spores brown, pedicellate or with sterigmatal scar.

From the specimens so far examined it is evident that Sedecula develops centripetally and has complete reduction of stipe and sterile base. There is practically no sterile tissue remaining over the lower surface of the fructification where the gleba is almost if not usually naked. The genus was first placed for convenience in the Sclerodermataceae (Mycologia 33:212. 1941) merely because of the heavy, leathery peridium. The sterigmatal scar or pedicellate character of the spores indicates closer relationship to Pompholyx than to Scleroderma, but because of the evidence of centripetal development it is here placed in a family by itself. Sedeculaceae, so far containing Sedecula only, is referred to the Sclerodermatales.

#### **NIDULARIALES**

1. Sphaerobolaceae. The chief recent contributions to our knowledge of this family have been made by Walker <sup>9</sup> and Greis. <sup>10</sup>

Walker has found two distinct types, Sphaerobolus stellatus Tode and S. iowensis Walker. Both of these species have the lacunar type of glebal development in which large buffer cells or space formers expand the glebal chambers before basidia are formed. In S. stellatus the basidia grow into and fill these spaces forming nests of basidia, as in the Melanogastraceae and Sclerodermataceae. (See Zeller, S. M., Developmental morphology of Alpova. Oregon State College Monographs in Botany No. 2. illus. 1939.) In S. iowensis the basidia form hymenial linings to otherwise hollow chambers. There are other differences. There is a gelatinous layer in the peridium in S. stellatus, not in S. iowensis, while the gleba of the latter dries soft or gluey and gelatinous and that of S. stellatus dries firm and hard.

Greis found a form of *S. iowensis* in Europe which he called forma *europaea*. In addition he has described the genus *Nidulariopsis*. The fruiting body here rises from a mycelial cord and

<sup>&</sup>lt;sup>9</sup> Walker, Leva B. Development and mechanism of discharge in *Sphaero-bolus iowensis* n. sp. and *S. stellatus* Tode. Jour. Elisha Mitchell Sci. Soc. 42: 151-178. *pl. 16-25*. 1927.

<sup>&</sup>lt;sup>10</sup> Greis, H. Nidulariopsis melanocarpa Greis nov. gen. nov. spec. und eine neue Form von Sphaerobolus iowensis. Hedwigia 75: 255-266. illus. 1935.

the gleba is like that of *Sphaerobolus iowensis* with cavities lined with an even basidial hymenium. He says a special peculiarity is that the middle layer of the peridium consists of rounded brownish cells with thick walls, and this layer does not reach across the summit of the fruiting body. But this peculiarity was recognized by Lohwag <sup>11</sup> who believed that this layer indicates a cup-shaped primordial (tramal cup) development. Walker's illustrations of *S. iowensis* (*Pl. 17, f. 11 & 13*) show this interrupted middle layer very plainly.

It appears that any differences between *Sphaerobolus iowensis* Walker and *Nidulariopsis melanocarpa* Greis are merely specific rather than generic and that the two should be included in the same genus *Nidulariopsis*. In this arrangement then *S. iowensis* becomes **Nidulariopsis iowensis** (Walker) n. comb.

Since the two genera are so specialized in peridial structure and discharge of the gleba they are retained in the same family, *Sphaerobolaceae*.

Fischer (1933) referred this family to the Sclerodermatales as a special family. Its closest relationship in this order is to the Calostomataceae, because of the similarity in discharge of gleba. However, in Calostoma the spore sac is inverted and ejected through the stoma after the gleba has become a powdery mass, mostly spores. It would seem more reasonable to follow the usual procedure and include the Sphaerobolaceae with the Nidulariaceae in the Nidulariales.

### **PODAXALES**

## 1. Secotium albipes sp. nov.

Fructificationes turbinatae, stipitatae; pileo circa 4 cm. alto, 5 cm. crasso, subgloboso, inferne a stipite inseparabili; superficie laevi, glabra, viscida, rubra, siccitate obscuriore; stipite 1–1.5 cm. longo, sursum crasso inferne tenuiori, solido vel farcto, superficie siccitate alba, interne obscuriore, superne in columellam crassam percurrentem procurrente, peridio circa 650  $\mu$  crasso (sicc.), prosenchymate gelatinoso composito, siccitate duro obscuroque; gleba obscure brunnea, locellis parvis; basidiis 1-, 2-, et 4-sporigeris; sporis brunneis, ellipsoideis, laevibus vel subrugulosis, guttulatis, 6–8 × 12–18  $\mu$ ; cystidiis magnis conicis hyalinis.

<sup>11</sup> Lohwag H. Die Homologien im Fruchtkörperbau der höheren Pilze. I und II. Biologia generalis 2: 148–182, 575–608. 1926.

terig-

e and over ost if nce in

Pom-

Sede-

o our reis.<sup>10</sup> ellatus e the

a are spaces clero-

Io. 2. inings

in S. gelat-

called Nidud and

haeroi. Soc.

1935.

Fructifications turbinate, stipitate, up to 4 cm. tall and 5 cm. wide; pileus subglobose; smooth, glabrous, viscid, red, drying dark mineral red, not breaking from the stipe below; stipe short, 1–1.5 cm. long, broad above and tapered sharply downward, solid, drying dark within but whitish on the surface; columella percurrent, broad, but broader above and below; peridium about 650  $\mu$  thick (dry), of gelified, prosenchymatous tissue, drying dark and hard; gleba dark brown, locules small; basidia 1-, 2-, and 4-spored; spores brownish, narrowly ellipsoid, smooth but with occasional raised spots, guttulate, 6–8 × 12–18  $\mu$ ; cystidia large, conic, hyaline.

On the ground in rich forest duff.

California: Butte county, Merrimac, *Thelma Norman*, Nov. 9, 1932, type (in N.Y.B.G. Herb.).

The above collection had been labeled "Secotium erythrocephalum Tulasne." This drew our attention again to the original description of the latter in which Tulasne describes the stipe as 'white.' Cunningham has described the stem as 'bright yellow' and several collections from New Zealand in American herbaria show yellow slender stipes.

Secotium albipes differs from S. erythrocephalum in other respects, however, than in stipe color. The edge of the peridium (pileus) does not readily separate from the stipe below as in S. erythrocephalum, which not only separates but often the edge is turned back (repandus Tulasne), thin, and coriaceous. The latter is doubtless a part of the fundamental veil separating from the stipe. The dried plants of S. albipes are short and stout whereas those of S. erythrocephalum are slender and tall. The spores of the latter are smooth and those of S. albipes are smooth but have a tendency to slight roughness and are definitely guttulate. S. albipes is similar to S. tenuipes Setchell in color of the peridium but the latter has a brown stipe and the spores are dark brown, rough and citriform as in those of Hymenogaster.

## 2. Secotium nubigenum Harkness.

A recent note <sup>12</sup> indicated the loss of the type of this species, but in the N.Y.B.G. there is a collection with the label in Harkness' handwriting as follows: "Secotium nubigenum Hks., on logs of Pinus contorta, Summit of the Sierra Nevada." An additional

<sup>12</sup> Zeller, S. M., Mycologia 33: 210. 1941.

note that accompanied the original description is "7000 ft." Since this specimen was collected by Harkness at the type locality I have taken the liberty to label it "type." Unless Harkness distributed type material elsewhere, this at least is the only collection so authentic.

3. Secotium sessile Massee and Rodway, in Rodway, Roy. Soc. Tasmania, Proc. 1911: 31. 1912. (Elasmomyces sessile Rodw. Roy. Soc. Tasmania, Proc. 1924: 8. 1925.)

A part of the type of the above is in the N.Y.B.G. Herbarium. The packet containing the collection is inscribed with the following in George Massee's handwriting: "Secotium sessile Mass. & Rodw., Tasmania, Rodway 649. type." The specimens are in every particular the same as Elasmomyces Mattirolanus Cavara, with which it becomes synonymous.

### TWO NEW ORDERS

The genera of the Gasteromycetes for the most part fall into the older, established, and quite generally accepted orders, as follows:

1. The Hymenogastrales, which include mostly hypogeous species that retain the original glebal structures to maturity. Such genera have been assigned to the families *Protogasteraceae*, *Gasterellaceae*, *Melanogasteraceae*, *Rhizopogonaceae*, *Hymenogasteraceae*, and *Hydnangiaceae*. They contain about twenty-two genera.

2. The Hysterangiales include hypogeous or epigeous species with gelatinous or cartilaginous tissues and smooth, ellipsoid spores (phalloid) but without a so-called receptacle. They have been assigned to the families *Hysterangiaceae*, *Protophallaceae*, and *Gelopellaceae*. The three families contain seven genera.

3. THE PHALLALES are divided into the three families—Claustulaceae, Phallaceae, and Clathraceae. The species here have bacillar spores, the peridium a volva with at least one gelatinous layer (and that layer interrupted by sutures of fundamental tissue in the Clathraceae), and the gleba variously dispersed or elevated on a pseudoparenchymatous stem, or a clathrate or hollow structure known as the receptacle. There are about twenty genera referred to this order.

ov. 9,

5 cm.

g dark

1-1.5

, dry-

irrent,

thick hard:

pored;

sional valine.

cephanal deipe as rellow'

as in e edge
The from stout

The mooth guttu-of the dark

pecies, Harkon logs itional

- 4. The Lycoperdales, which include the "puff ball" types of fructifications having ordinary hyphal tissues that are not gelatinous or cartilaginous, the gleba disintegrating into a powder or into small hollow peridioles at maturity, and with a capillitium (except in Arachniaceae). The species here are assigned to the families Arachniaceae, Broomeiaceae, Mycenastraceae, Lycoperdaceae, Mesophelliaceae, and Geastraceae. These include about twenty-five genera.
- 5. The Sclerodermatales that include species with heavy peridial and other tissues, and gleba pulverulent at maturity; basidia symmetrically distributed or in nests or cavities arising through the dissolution of the tissue, at least without a well organized hymenium (except possibly in *Batarrea*). The order has been divided into the families *Sclerodermataceae*, *Pisolithaceae*, *Glischrodermataceae*, *Sedeculaceae*, *Astraeaceae*, *Tulostomataceae*, and *Calostomataceae*. These families include seventeen genera.
- 6. The NIDULARIALES, which are the distinctive "Bird's Nest" fungi and need no description here but are divided into the two families Nidulariaceae and Sphaerobolaceae, including six genera.
- 7. The Podaxales, the species of which have a percurrent columella or stem reaching to the summit of the fructification; peridium left at maturity in part as a pileus or volva, or annulus on the stem; gleba at first with hymenium of basidia covering the walls of chambers or pores or lamellae, persistent or pulverulent; spores colored. The order has two families, the Secotiaceae and the Podaxaceae, containing seven genera.

This treatment of the Gasteromycetes leaves four genera to be considered. These are Clathrogaster, Gasterosporium, Gautieria, and Tremellogaster. Tremellogaster and Clathrogaster may readily be placed in the same family because of their peridial and glebal characters. The peridia of these two genera are composed of at least two layers. The outer layer is of filamentous, fundamental tissue, forming a thin rind. Under this there is a thick gelatinous layer. In this layer the hyphae are far apart but are dispersed through a thick gel. This gelatinous layer is interrupted here and there by more or less radial plates (sutures) of more compactly interwoven hyphae, the same as the outer layer, and connecting the latter with the gleba, much as in the family Protophallaceae

and in the early button stages of the *Clathraceae*. In the peridium of *Tremellogaster* these sutures are more numerous and complicated than in *Clathrogaster*.

oes of

inous

except

, Me-

y-five

heavy

urity;

rising 11 or-

r has

iceae,

iceae.

Nest"

e two enera.

rrent

tion; nulus

g the

ilent:

and

to be

*ieria,* read-

debal of at

ental

ersed

and

actly cting

iceae

ra.

The peridium of Gastrosporium as previously reported <sup>13</sup> is very similar to that of the Gelopellaceae and to the volva in the button stages of the Phallaceae. The glebae of Gastrosporium and Tremellogaster are pulverulent at maturity. Whether the gleba of Clathrogaster retains its original structure to maturity is not known, but in other respects it is similar enough to Tremellogaster to be included in the same family with it. The type material of Clathrogaster was originally, at least, kept in preservative and the development of the gleba was arrested as at the time of collection. All of these genera, however, have spherical, echinulate or sculptured spores and, therefore, do not belong in the series of genera with phalloid spores, such as those of the Hysterangiales and Phallales.

To judge only from the gelatinous or cartilaginous character of the tissues in *Gautieria* this genus should take its place somewhere in the *Melanogastraceae*, *Hysterangiaceae*, or possibly the Sclerodermatales. Glebal characters, however, make it incompatible with the *Melanogastraceae* and the Sclerodermatales and the spore characters would keep it out of the *Hysterangiaceae*. One might stretch a point and place the genus *Gautieria* with the *Hymenogastraceae* having like spores, but it does not otherwise belong.

Two new orders are consequently proposed to receive these genera. The first, the **Tremellogastrales**, contains two families; the **Tremellogasteraceae** n. fam., including the genera *Clathrogaster* Petri and *Tremellogaster* Ed. Fischer, and the *Gastrosporiaceae* Pilat, including the one genus *Gastrosporium* Mattirolo. The second new order, the **Gautieriales**, contains the **Gautieriaceae** n. fam. with the one genus *Gautieria* Vittadini.

### TREMELLOGASTRALES n. ordo

Fructificationes hypogeae vel epigeae, sessiles; peridio duplici, strato externo hyphis intricatis composito, strato interno gelatinoso continuo vel laminis radiantibus interrupto investitoque; gleba primo carnosa, dein pul-

<sup>&</sup>lt;sup>13</sup> Zeller, S. M. New and noteworthy Gasteromycetes. Mycologia 31: 1-32. 1939 (see p. 17).

verulenta; columella simplici vel dendroidea, vel deficienti; sporis sphaericis, echinulatis vel verrucosis.

Fructifications hypogeous or epigeous, mostly sessile; peridium of two or more layers, the outer of fundamental tissue, the inner of a gelatinous nature, continuous or interrupted by sutures of fundamental tissue; gleba centripetally developed, pulverulent at maturity; columella simple or wanting; spores spherical, echinulate or verrucose.

#### KEY TO THE FAMILIES OF THE ORDER TREMELLOGASTRALES

- I. Peridium with a gelatinous inner layer interrupted by sutures of fundamental tissue; spores spherical, echinulate or rough.. Tremellogasteraceae

## A. Tremellogasteraceae n. fam.

Fructificationes hypogeae vel epigeae, sessiles; peridio duplici, strato externo hyphis intricatis composito, strato interno gelatinoso laminis radiantibus interrupto investitoque; gleba ochracea vel flavida, primo carnosa, dein pulverulenti; columella dendroidea vel deficienti; basidiosporis sphaericis, echinulatis vel cristatis.

Fructifications epigeous or hypogeous, subglobose, sessile; peridium of two or more layers; exoperidium of fundamental tissues; endoperidium of one or more layers of gelatinous tissue interrupted by plates of fundamental tissue; gleba developing centripetally from hyphae from the margins of the medulla of the primordium, pulverulent at maturity; capillitium rudimentary; columella sometimes present; spores spherical, echinulate or cristate.

#### KEY TO THE GENERA OF THE TREMELLOGASTERACEAE

- Tremellogaster Ed. Fischer, in Mitteil. Naturf. Ges. Bern 1923: 49-56. 1924.

Emended description: Fructifications epigeous, subglobose; surface drying coarsely flattened-tuberculate; peridium very thick, up

ridium inner res of ent at chinu-

aericis,

fundaeraceae utures ; riaceae

ato exantibus a, dein aericis,

peridssues; intertripee pricoluistate.

al, filan radiechinuogaster ore or eed, or

Bern

; surk, up to  $\frac{1}{6}$  the diameter of the sporocarp, with an outer layer of thick-walled, sclerotioid hyphae bordered internally by thin-walled, hyaline periclinal hyphae, a middle layer is brownish, gelatinous, reticulately divided by lighter colored nongelatinous, fundamental tissue (partitions or sutures), and a white inner layer is nongelatinous, of hyphae that intertwine and run parallel with the surface of the gleba; the latter ochre to brown, becoming pulverulent at maturity; pseudocapillitium various lengths, 2.5–4  $\mu$  in diameter, much branched, hyaline, sparsely warty-spinulose; basidia forming a palisade-like hymenium on the walls of lacunae through which many hyphae penetrate, 4-spored; sterigmata short; spores spherical, echinulate, dark brown, 5–6  $\mu$  in diameter.

Type species: Tremellogaster surinamensis Ed. Fischer.

Habitat: On moist sandy soil, near decaying wood.

Distribution: British and Dutch Guiana, South America.

There is but the one species in the genus, and the above description serves very well for the species.

2. CLATHROGASTER Petri, Malpighia 14: 125-126. 1900. Description emended.

Fructifications hypogeous, subglobose, radicate, rhizomorph attached in a depression in the base; peridium thick, of 2 layers, outer of fundamental, periclinal hyphae, thin, silky, reticulately furrowed; inner layer thick, gelatinous, interrupted by more or less radial sutures of tissue like the outer peridium and connecting the latter with the gleba; lactiferous ducts large, long, penetrating all sterile tissues; gleba chambered, spongy, yellowish, cavities spherical to irregular, larger toward the outside, with one to many very large radiating, gelatinous, sterile cavities; septa broad, with stipitate basidia and cystidia on both sides; columella or prominent tramal plates radiating from the base or from a pulvinate sterile base; basidia mostly 2-spored; spores spherical, yellowish, reticulate with interrupted ridges, pedicellate.

The type species: Clathrogaster volvarius Petri. Distribution: Sarawak, near Sibu, Borneo.

a. Clathrogaster volvarius Petri, Malpighia 14: 126. pl. 2, f. 1-2; pl. 3, f. 2, 3, 5-8, 10, 13; pl. 4, f. 1. 1900. (Arcangeliella volvaria (Petri) Z. & D. Mo. Bot. Gard. Ann. 22: 369. 1935; 23: 629. 1936.)

Fructifications irregular reniform about 4 × 6 cm., russet (in alcohol, 1934), surface irregularly reticulate-sulcate; sterile base scarcely more than a thickening of the peridium; columella conspicuous, branching near the base but branches percurrent or nearly so, gelatinous with many lactiferous ducts; peridium 1200-1440 \(mu\) thick (in alcohol), duplex, the outer layer (rind) of densely tangled hyphae, tough, somewhat gelatinous, the inner a thick gelatinous layer (tramal peridium), with many large lactiferous ducts, interrupted by radial plates (sutures) of tissue like the rind which connects the outer peridium and the gleba; the latter ochraceous-tawny, cavities ovoid, radiating from the columella and base; septa thick, of loose, gelatinous hyphae, with lactiferous ducts; basidia subcylindrical, 37-40 μ long, upper part collapsing after the separation of the spores; sterigmata short; spores spherical with short ridges and slender, blunt spines, yellowish, 9-12 µ in diameter.

Type locality: Sarawak, Borneo.

Distribution: Known from type locality only.

In his formal description of the species *C. volvarius*, Petri used the old spelling "vulvarius," but in all other cases throughout his paper he used "volvarius." Undoubtedly the use of the "u" in the one case was unintentional and thus our correction.

Petri did not designate a type species of Clathrogaster. C. volvarius Petri was chosen as the type since it was the better described and illustrated of the two species.

b. Clathrogaster Beccarii Petri, Malpighia 14: 126. pl. 2, f. 3-5,
7-9. 1900. (ut C. Beccari.) (Arcangeliella Beccarii [Petri]
Z. & D. Mo. Bot. Gard. Ann. 22: 366. 1935; 23: 635-636.
1936.)

Fructifications spherical to reniform, 1–3 cm. in diameter, raw sienna in alcohol, surface smooth but with low reticulated ridges; sterile base and columella not perceptible; peridium about 600–700  $\mu$  thick (in alcohol), duplex, the outer layer of compactly interwoven hyphae producing a rind, the inner layer thick, gelatinous of loosely woven hyphae mixed with lactiferous ducts, interrupted by radial plates (sutures) of fundamental tissue like the rind, and uniting the latter with the gleba; the latter amber-brown, cavities elongate, radiating from the base which is scarcely more than a thickened peridium; septa about 110  $\mu$  thick, similar to the inner peridium in structure; basidia clavate, 2-spored, about 80  $\times$ 

 $11~\mu$ , only the outer half collapsing after the separation of the spores; sterigmata short; spores  $11-15~\mu$  in diameter, spherical, with very high ridges, irregularly disposed over the surface, yellow.

Type locality: Sarawak, Borneo.

Distribution: Known from the type locality only.

B. Gastrosporiaceae Pilát, Bull. Soc. Myc. France 50: 45–46. 1934.

Description emended: Fructifications globose, hypogeous; peridium duplex; exoperidium of filamentous, fundamental tissues; endoperidium a continuous gelatinous layer, not interrupted by plates of fundamental tissue; gleba developing centripetally from anastomosing lamellae, produced subperidially, pulverulent at maturity; columella present; capillitium rudimentary; hymenium lining walls of cavities; spores spherical, minutely verrucose.

There is one genus, Gastrosporium.

Gastrosporium Mattirolo, Memoria Accad. Sci. Torino, Ser. II 53: 361. 1903. (*Leucorhizon* Velenovsky in Mykologie 2 (3-4): 49-51. f. 1-4. 1925.)

Description emended: Fructifications hypogeous, globose; surface soft, white, dry; peridium duplex; outer layer filamentous, easily separable; inner layer tough, gelatinous, cartilaginous, indehiscent, continuous, easily distinguished from the gleba; the latter filling the whole peridium (no sterile base), white, becoming ochraceous to subolivaceous, with main tramal plates extending from a columella which is simple but reaching beyond the center of the fructification; hymenium lining cavities at first, but the whole pulverulent at maturity; capillitium rudimentary; spores light ochraceous, spherical, slightly verrucose.

Type species: Gastrosporium simplex Mattirolo.

Habitat: In the soil around the roots of grasses and sedges.

Distribution: Northern Italy and Czechoslovakia.

There is but one species in the genus Gastrosporium. Velenovsky, however, described a species Leucorhison nidificum which becomes synonymous with G. simplex. It is peculiar that Velenovsky, Pilát 14 and Mattirolo overlooked the fine verrucosity of

<sup>14</sup> Pilát, A. Sur le genre Gastrosporium Mattirolo. Bull. Soc. Myc. France 50: 37-49. illus. 1934.

used ut his u" in

et (in

base

nt or

1200 -

ensely thick

erous

e the

a and

erous

psing

pheri-

 $-12 \mu$ 

volr de-

. 3–5, Petri] –636.

dges; 600ly ingelatinter-

more o the 80 ×

e the

the spores. A collection under the name *Leucorhizon nidificum* Vel. from Bohemia is to be found in the Lloyd collections (Smithsonian Inst. Cat. No. 148).

## GAUTIERIALES n. ordo

Fructificationes hypogeae, sessiles, basi rhizomorphico instructae; peridio deficienti vel hyphis laxe intricatis vel pseudoparenchymatibus composito; gleba primo albida translucento-cartilaginea, dein brunneola cartilaginea; columella simplici vel dendroidea, cartilaginea; septis hyphis gelificatis compositis, hymenophori; basidiosporis crasse fusiformibus, longitudinaliter costatis, brunneis.

Fructifications hypogeous, sessile; peridium usually wanting, when present stupose, loosely filamentous, or pseudoparenchymatous; gleba gristly-translucent, whitish, becoming brownish as spores mature, with a columella from a basal rhizomorph; basidia in a hymenium; septa usually gelatinous-cartilaginous, of gelified hyphae; basidiospores of various shapes, mostly broad fusiform, longitudinally costate, brown.

Gautieriaceae n. fam. Characteribus ordinis.

The order Gautieriales, represented by the one genus Gautieria. partakes of the characters of several other groups of gasteromycetes but differs fundamentally from each enough that it cannot logically be classified with any. The fructifications follow the centrifugal, coralloid pattern of development and the gleba has a conspicuously gelatinous-cartilaginous consistency such as found in the Hyster ngiales, but the gleba and spores do not partake of the phalloid nature of those of the latter. Nor does the peridium when present in species of Grazieria have the gelatinous or cartilaginous layers found in the me parts of the Hysterangiales. The spore type in the Gautieriales is similar to that found in Hymenogaster, Gymnoglossum, and Secotium, but the development of the fructifications is distinct as is the nature of the sterile The order Gautieriales differs from the Tremellogastrales in four fundamental respects. In the latter the gleba develops centripetally (lacunar), the peridium is characterized by gelatinous layers, the spores are spherical, and echinulate or verrucose, and the gleba becomes powdery at maturity.

To obtain the best concept of the genus Gautieria it is necessary to know several of its species as they occur in the field, so to mith-

ficum

peridio posito; ginea; comr cos-

nting, ymah as isidia lified form,

ieria, omyannot the has a found ke of dium

cariales. id in

terile rales elops nous

ssary so to

and

speak. Otherwise one does not become aware of the tough, gristly nature of the sterile tissues in most of the species. For instance, the type species G. morchelliformis Vitt. has such large glebal cavities and thin tramal septa that, although it has the same characters otherwise as the other species, the observer is not so impressed with its gelatinous-cartilaginous nature, as when he sees some other species of the genus with thicker septa, columella, etc. In fact the examination of dried specimens of G. morchelliformis may not reveal the real cartilaginous character of its tissues. In other words the type species of Gautieria does not represent the norm of the genus, but may be said to be almost marginal.

1. GAUTIERIA Vittadini, Monogr. Tuberac. 25–27. 1831.—not Gautiera Rafinesque Med. Fl. 1: 202. 1828. (Syn. Chamonixia Rolland, Soc. Myc. France Bul. 15: 76–77. 1899.)

Description emended: Fructifications subspherical to irregularly depressed, with a simple or branched rhizomorph sometimes persisting as a short stipe; columella variable, simple or branched, gelatinous to translucent-cartilaginous; peridium wanting, evanescent, or persistent, when early evanescent or wanting outer septa sterile over the surface; gleba white or gristly-translucent, becoming brownish as spores mature; cavities labyrinthiform; septa usually thick, gelatinous-cartilaginous, of interwoven, gelatinized hyphae; basidia in hymenium covering both sides of septa; spores ovoid, ellipsoid, fusiform, with longitudinal striae, brown.

Type species: Gautieria morchelliformis Vitt.

Habitat: Wholly or partially hypogeous under various kinds of shrubs and trees.

Distribution: Europe, Asia, Africa, North and South America, Australia, Tasmania, New Zealand.

The species of Gautieria have been published elsewhere.15

Eleven species have been reported from North America and eight others occur elsewhere in the world.

## A NEW FORM ORDER OF FUNGI IMPERFECTI

Lycoperdellon Torrend (in Broteria Sér. Bot. 11: 92. 1913) has had a very doubtful taxonomic position. Fischer (1933, p.

<sup>15</sup> Dodge, C. W., and S. M. Zeller, Mo. Bot. Gard. Ann. 21: 692–705. pl. 18, f. 51–66. 1934.

72) referred it to the doubtful list of genera under the Lycoperdaceae. To be sure, the fructifications of Lycoperdellon have the general appearance of a myxomycete, like Lycogala, or a lycoperdaceous gasteromycete but there are no basidia or basidiospores. Lohwag 16 accordingly recognized the sporophores as real conidiophores and thought them to represent the conidial stage of an ascomycete. The conidial stages of ascomycetes so far as we know, however, do not take on the form of a closed sporocarp similar to a gasteromycete. There is no doubt Lycoperdellon is one of the Fungi Imperfecti, but its closest relatives are doubtless among such forms as Leucophleps Harkness, all the species of which, except L. candida Harkness, have proved to be the conidial stage of one or another of the species of the gasteromycetous genus, Leucogaster Hesse. After considerable careful study of Lycoperdellon by Heim and Malençon 17 who continued to designate its spores as "conidia," Heim described the family Lycoperdellaceae to receive the genus Lycoperdellon with its two species L. Torrendii (Bresad.) Torrend and L. minutum Heim.18 Unfortunately he assigned the family to the Gasteromycetes whereas from our viewpoint it belongs in the Imperfecti. It is, therefore, proposed to transfer the family Lycoperdellaceae Heim to the Fungi Imperfecti and refer to it the form genera Lycoperdellon Torrend and Leucophleps Harkness. There will need also to be the Form Order Lycoperdellales 19 coordinate with the Phyllostictales, Melanconiales, and Moniliales. In a key like Martin's 20 the new form order would key out as follows: "Fructification determinate, gasteromycetoid; conidia borne in chambered cavities or nests."

Leucophleps candida Harkness has been found in western Oregon and central California. Neither species of Lycoperdellon has been reported from North America.

<sup>&</sup>lt;sup>16</sup> Lohwag, H. Zu Lycoperdellon. Ann. Myc. 32: 244-255. 1934.

<sup>&</sup>lt;sup>17</sup> Heim, Roger, and G. Malençon. Le genre Lycoperdellon: structure et position taxonomique. Rev. Gen. de Bot. 45: 53-67. 1933.

<sup>&</sup>lt;sup>18</sup> Heim, R. Fungi Ibirici. Treballs del Mus. Cien. Nat. Barcelona 15 (No. 3): 1-146. 1934. (See pp. 138-141.)

<sup>19</sup> Fructificationes gasteromycetoides, sed conidia gignentes.

<sup>&</sup>lt;sup>20</sup> Martin, G. W. Key to the families of fungi. Univ. Iowa Stud. in Nat. Hist. 17: 83-115. 1936. (See p. 90.)

## STUDIES IN THE DARK-SPORED AGARICS \*

ALEXANDER H. SMITH

(WITH 93 FIGURES)

The present study deals chiefly with notes on the microscopic characters of the types of some of the species of Coprinus, Naematoloma, Panaeolus and Psilocybe described from North America. As has long been recognized by investigators who are working in this group, the descriptive accounts, whether old or new, which do not place great emphasis on spore morphology, on the characters of the cystidia if such are present, and on the anatomical features of the fruiting bodies, are of little value in accurately delimiting taxonomic entities of any category. It naturally follows, then, that one of the first steps in a critical revision of these genera is to ascertain the diagnostic characters from the type specimens in so far as the latter are available. However, as I found out early in my work in this group, an extensive knowledge of these fungi as they occur in nature is an invaluable aid in interpreting the characters of dried specimens. Consequently, although my investigations in the group have been in progress for over ten years, it is only comparatively recently that I have established species concepts in many of the genera that are satisfactory to me, and which serve satisfactorily as the building units for a classification which is other than just haphazard.

Considerable progress has been made in the recognition of species in the field through the process of mass collections, *i.e.*, collecting material of *all* fruitings of dark-spored agarics found during a season. Through such methods one soon learns to recognize the frequently-encountered species and to ascertain the constancy of their characters even though he may be unable to apply binomials correctly to them. The studies of type specimens do much

coperve the coperpores. midioasco-

howr to a Fungi forms canone or

Heim nidia," genus Tored the it be-

gaster

refer the refer phleps Lyconiales, order astero-

on has

ona 15

in Nat.

<sup>\*</sup> Papers from the Herbarium and the Department of Botany of the University of Michigan. The cost of the illustrations was paid for by the University of Michigan Herbarium.

to clear up the application of names, and the final result should be a satisfactory classification in which the species are adequately described. This, at least, is the aim in the manuscript being prepared for the *North American Flora*.


Because of the interest being shown in dark spored agarics both by European and American workers at the present time, and because of the general impetus given to the study of fleshy fungi by those seeking new antibiotics for medicinal use, it is urgent to make available such data as that presented here. Since the characters of the spores are fundamental, and since they have been badly handled by past investigators, detailed descriptions as well as illustrations are included. Although the genera treated here are classed as "dark-spored" fungi, it must be remembered that in Stropharia, Naematoloma and Psilocybe this character breaks down, and species with earth-brown to cinnamon-brown spore deposits must be admitted if any emphasis at all is placed on obvious natural relationships. The more material I examine, the more I become convinced that actually the genus Agrocybe of modern authors should be placed here also.

For access to type specimens reported on here I am indebted to Dr. H. D. House, State Botanist, New York State Museum, Albany, New York, to Dr. Fred J. Seaver, of the New York Botanical Garden and to Dr. Rolf Singer, Farlow Herbarium, Harvard University, Cambridge, Mass.

#### **COPRINUS**

COPRINUS ANGULATUS Peck, Ann. Rep. N. Y. State Mus. 26: 60. 1874. (Figs. 1–2–type.) Spores 8.5–10 (11)  $\times$  5.7–6.3  $\times$  7–8.8  $\mu$ , dark bister to blackish revived in KOH, somewhat flattened, truncate-ellipsoid in side view, resembling a blunt arrow head in face view, apical hyaline pore broad and conspicuous; basidia 18–32  $\times$  7–9  $\mu$ , four-spored, trimorphic; paraphyses inflated, hyaline, readily collapsing; pleurocystidia voluminous, ellipsoid to cylindric (50) 60–100 (120)  $\times$  (15) 20–60  $\mu$ , hyaline, smooth, thin-walled and readily collapsing; cheilocystidia vesiculose and 15–40  $\mu$  in diam., or fusoid-ventricose [the latter 38–46 (55)  $\times$  (7) 9–14  $\mu$ ], both types thin-walled and hyaline; gill trama hyaline in KOH or colored cinnamon-brown toward pileus trama; pileus trama with a cuticle of vesiculose cells one cell deep, from

among them arise numerous pilocystidia (32)  $60-90 \times 9-16 \,\mu$ , subcylindric to ventricose at base, apices obtuse to subacute, the walls thin and hyaline or slightly thickened in basal part and either hyaline or tawny as revived in KOH; flesh beneath cuticle tawny to cinnamon-brown in KOH.



Figs. 1-9. Microscopic characters of dark-spored agarics.

uld be uately g pre-

debted iseum, York arium,

is. 26: -6.3 × t flatarrow cuous; es inus, elyaline, resicu-38-46 trama trama; from Discussion. Peck's original description and comments indicate that this species belongs in the *C. ephemerus* group, and the present study of the type makes this disposition unquestionable. The pilocystidia are abundant and unmistakable.

COPRINUS CALYPTRATUS Peck, Bull. Torr. Club 22: 205. 1895. (Fig. 4-type.) Spores  $17-21\times 10-13\,\mu$ , not appreciably flattened, hyaline pore small and slightly eccentric in side view of spore, walls only moderately thick and in KOH dark bister (no other details obtainable from type, and spores examined were slightly immature.

Discussion. I consider Coprinus asterophorus Long and Mentzer, Mycologia 37: 120. 1945 to be a synonym of this species. Its spores (Fig. 6) are not quite as broad in the one collection illustrated, but Long and Mentzer described them as  $14-20\times 10-12.7\,\mu$ , and in material sent me by the late Dr. Long considerable variation was evident. The patch of the volva which forms a yellowish cap on the pileus is apparently the important diagnostic field character. In its spore characters the fungus is very similar to Coprinus sterquilinus but differs sharply in the more compact organization of the veil tissue.

COPRINUS BRASSICAE Peck, Ann. Rep. N. Y. State Mus. 43: 18. 1890. (Fig. 3-type.) Spores 6.2– $7.8 \times 3.6$ – $4 \mu$ , dull cocoa brown when first revived in KOH, not flattened, ellipsoid in either view, hyaline apical pore distinct but small; basidia four-spored, hyaline in KOH; (no other microscopic characters determinable on material examined).

Discussion. Unfortunately the characters of the veil could not be determined. These are important in this group. The species has received considerable attention in the American literature.

Coprinus cinchonensis Murrill, Mycologia 10: 85. 1918. (Figs. 5 & 7-type.) Spores 9-11 × 5-6 × 6-7.5  $\mu$ , subelliptic in side view, ovoid in face view, terete or only slightly flattened, smooth, pale bister in KOH, apical pore hyaline and distinct; basidia hyaline in KOH, four-spored, 7-8  $\mu$  broad at apex; paraphyses hyaline, thin-walled, inflated, 10-14  $\mu$  diam.; pleurocystidia apparently present but remaining collapsed and details not clear, hyaline and thin-walled; cheilocystidia none seen; gill and pileus trama hyaline in KOH; universal veil remnants in the form of matted fibrils in patches over the disc (no globose cells seen),

the hyphae much-branched, with somewhat thickened walls and with scattered short thorn-like processes unevenly distributed over them, the cells nearly hyaline to slightly yellowish in KOH.

Discussion. This is a distinctive species if one considers the characters of the veil and spores in conjunction with the habitat on a log. It appears to be closely related to Coprinus Brassicae Peck but is readily distinguished on spore size.

COPRINUS EBULBOSUS Peck, Bull. Torrey Club 22: 491. 1895. (Fig. 8–type.) Spores 7–8.4 × 4.4–5  $\mu$ , dull chocolate-brown revived in KOH, smooth but often with a ventral hump as seen in side view (apparently a second pore), oblong to subellipsoid in face view, subellipsoid to slightly inequilateral in side view, apical pore present but obscure; basidia four-spored, hyaline in KOH, 18–24 × 7–8  $\mu$ , apparently trimorphic; paraphyses 10–16 × 10–16  $\mu$ , subglobose, hyaline in KOH; pleurocystidia scattered; subcylindric, 80–120 × 18–32  $\mu$ , thin-walled and readily collapsing; cheilocystidia none seen; gill trama of filamentous hyphae, hyaline in KOH; pileus trama hyaline in KOH, cuticle of vesiculose hyaline cells about 1 cell deep.

Discussion. Both the collection from Lyndonville, N. Y. and the one from Kansas show the curious ventral hump on at least some spores so there is no doubt but that both represent the same species. It is very closely related to C. quadrifidus Pk., but apparently differs in slightly narrower spores. However, this difference is not truly significant. From the information at hand it is not clear whether the ventral hump or second pore is an abnormality which occasionally occurs in the members of the quadrifidus group and is comparable to the bifid spores as reported by Lange (p. 743) for C. myceliocephalus, or whether it is a useful character.

COPRINUS HEXAGONOSPORUS Josserand, Bull. Soc. Myc. Fr. in press. (Fig. 9.) Pileus cylindric at first, 5–10 (15) mm. high, 4–6 (8) mm. across the base, very finely pubescent at least over the disc, disc "fuscous," marginal area pallid to "avellaneous," tinged "army brown" near the disc at maturity, surface closely folded-striate before expanding, becoming broadly convex to plane and plicate-striate before maturity; flesh very delicate and thin, odor none, taste mild to slightly bitterish; lamellae free but attached to apex of stipe, close, whitish, soon avellaneous and then black, narrow and equal (1–2 mm. broad), soon deliquescing; stipe 3–5 (10) cm. long, 0.5–1 (2.5) mm. thick, equal, hollow,

The 1895. flatew of

r (no

were

dicate

pres-

and f this e one em as Long which

ortant

gus is

s. 43: cocoa either pored, inable

ld not pecies re.

1918. otic in tened, stinct; araphestidia clear, pileus rm of seen), very fragile, sparingly pubescent at first from projecting caulocystidia, densely pubescent above, white at first but soon sordid brownish over lower two thirds, base inserted on the substratum.


Spores black in deposit,  $11-13\times5-6.2\times6.5-8\,\mu$ , black in KOH, flattened, subelliptic in side view, angular and obscurely six sided in face view, pore eccentric as seen in side view; basidia trimorphic,  $18-26\times8-10\,\mu$ ; paraphyses  $18-22\times9-16\,\mu$ ; pleurocystidia rare,  $60-90\times20-35\,\mu$ , becoming more elongated and narrower, readily collapsing; cheilocystidia similar to pleurocystidia or more vesiculose,  $40-80\times35-45\,\mu$ ; gill trama narrow and of very loosely interwoven cells; pileus trama with a cuticle formed by a row of vesiculose cells from which thin-walled pilocystidia project, the latter  $70-120\times10-15\,\mu$ , trama proper very thin but hyphae filamentous, those in the dark discal area dark sordid brown (almost vinaceous brown) when revived in KOH.

Discussion. This fungus, a close relative of C. ephemerus, is apparently quite common on horse dung in the United States. I have isolated it on numerous occasions. Brooks obtained it from horse dung collected in Kansas, and both Brooks and Lange isolated it from the same substratum collected near Ann Arbor. The spores in face view are the outstanding feature of the species, but once the fungus is known it can usually be identified at sight. It has not been previously reported from North America.

COPRINUS INSIGNIS Peck, Ann. Rept. N. Y. State Mus. 26: 60. 1874. (Fig. 10–type.) Spores 10–12.6  $\times$  7–8.4  $\mu$ , dark bister in KOH, the exospore wrinkled and cracked to form obscure warts or causing surface to appear decidedly uneven, in side view slightly inequilateral, in face view oblong or base somewhat pointed, apex with a protruding lens-shaped hyaline pore causing it to appear snout-like; basidia four-spored, hyaline in KOH, paraphyses inflated, readily collapsing; pleurocystidia apparently subcylindric, elongated and projecting across the gill cavity (but remaining collapsed); cheilocystidia none seen; gill and pileus trama hyaline in KOH, the latter homogeneous or nearly so (with no sharply defined cuticle in sections of revived material).

Discussion. This well-known and characteristic species of the C. atramentarius group is known from both North America and Europe. The occurrence in several sections of Coprinus of rough spored and smooth-spored species with almost identical macroscopic characters is interesting in the light of recent efforts to

place great emphasis on spore markings in determining natural relationships among the fleshy fungi. The markings on the spores of *Coprinus* form excellent characters for the recognition of species, but if a natural arrangement is desired, it would be a serious



Figs. 10-22. Microscopic characters of dark-spored agarics.

caulosordid tratum. KOH, x sided dia tripleurond narcystidia and of formed

cystidia hin but sordid

States. ined it Lange Arbor. species, t sight.

26: 60. ister in e warts slightly d, apex appear ses in-lindric, naining hyaline sharply

of the ica and rough macroorts to mistake to insist that all species with rough spores were more closely related to each other than to any with smooth spores. It is clearer in *Coprinus* than in almost any other genus that the character of ornamented spores has evolved independently on numerous occasions.

Coprinus Jalopensis Murrill, Mycologia 10:83. 1918. (Fig. 11–type.) Spores 6– $6.3 \times 3.1$ – $3.6 \,\mu$ , cocoa-color as revived in KOH but gradually becoming grayish, smooth, not flattened, slightly bean-shaped in side view, subelliptic in face view, smooth, apical pore hyaline and inconspicuous; basidia hyaline in KOH, 12– $18 \times 5$ – $5.5 \,\mu$ , four-spored, trimorphic; paraphyses thin-walled, hyaline, readily collapsing, 9– $12 \times 6$ –9 (12)  $\mu$ , enlarging to greatest size at time spores are discharged; pleurocystidia abundant and voluminous, hyaline, thin-walled and readily collapsing, extending across the gill cavity and 20– $40 \,\mu$  in diam.; cheilocystidia not reviving, the gill edges whitish apparently from filamentous hyphae; gill trama not reviving well but hyaline in KOH; pileus trama tawny yellowish to tawny-orange just under the cuticle, cuticle of a single layer of inflated cells, no pilocystidia or universal veil remnants found.

Discussion. This species must belong in the C. radians group, though its veil characters, if a veil is present, are not known. The oozonium and the small cocoa-colored spores along with the lignicolous habitat are certainly suggestive. The very small spores appear to characterize it in this group.

COPRINUS JONESII Peck, Bull. Torrey Club 22: 205. 1895. (Fig. 12–type.) Spores 7–9 × 6–7  $\mu$ , terete to very slightly compressed, subellipsoid in side view, subcircular in face view, black or nearly so in KOH, exospore separable from endospore only under considerable pressure (enough to break inner spore wall in many instances), apical hyaline pore broad and distinct; basidia four-spored, hyaline in KOH, 7–8  $\mu$  in diam., trimorphic; paraphyses hyaline, thin-walled, readily collapsing, 10–12  $\mu$  broad; pleurocystidia projecting across gill cavity but not reviving well, apparently long-cylindric and thin-walled; cheilocystidia none seen; gill and pileus trama hyaline in KOH; universal veil remnants filamentous, hyaline to sordid brownish in KOH.

Discussion. A member of the C. lagopus series, but distinct on spore characters. C. lagopus var. rotundisporus Kuhner & Josserand is either identical with or very close to Peck's species.

COPRINUS LANIGER Peck, Bull. Torrey Club 22: 491. 1895. (Fig. 14-type.) Spores dark sordid reddish brown when first revived in KOH but soon chocolate-gravish, 8-10.5 × 3.5-4.2 µ, smooth, not flattened, narrowly ellipsoid to subcylindric in face view, in side view almost straight to decidedly concave on inner side and the back distinctly to merely obscurely convex, lensshaped pore apical and conspicuous when not collapsed; basidia four-spored,  $14-24 \times 5-6 \mu$ , trimorphic, hyaline in KOH; paraphyses inflated,  $12-15 \times 8-12 \mu$ , hyaline, readily collapsing; pleurocystidia scattered, ellipsoid to subcylindric, 50-90 (or more) × 16-30 µ, thin-walled and readily collapsing; cheilocystidia not seen (cells had deliquesced); gill trama hyaline to faintly yellowish in KOH; pileus trama merely yellowish in KOH, the cuticle of inflated cells one cell deep; universal veil remnants distinctive: the layer next to cap surface of chains of globose to keg-shaped hyaline cells, the cells of the chains more elongated outward as well as more brownish and the filaments finally composed of more or less ellipsoid to cylindric, tawny-cinnamon (in KOH), somewhat thick-walled cells which are smooth except for obscure zones and lines of encrusting pigment, the end cell often somewhat fusoid and cystidium-like, the chains of cells showing a tendency to break up into short segments or individual cells.

Discussion. This species is very closely related to C. domesticus, but differs from it in the long narrow spores with the very broad germ pore. The clustered habit of growth is a less constant distinguishing character. It is one of the extreme variants of the domesticus-radians series which at least for the present is being recognized as a species.

COPRINUS MEXICANUS Murrill, Mycologia 10: 84. 1918. (Fig. 15–type.) Spores 4.7–5.3 (6)  $\times$  3.1–3.6  $\mu$ , pale cocoa-brown but soon changing to pale avellaneous when revived in KOH, many nearly hyaline, smooth, not flattened, ellipsoid to slightly inequilateral in side view, ellipsoid in face view, apical hyaline pore very small and inconspicuous; basidia and paraphyses not reviving; some fusoid-ventricose cystidia seen in crushed mounts of gills, these  $28–36\times9–12\,\mu$  and with obtuse to subacute apices; gill trama not reviving; pileus trama with a cuticle of narrow  $(3–5\,\mu)$  radially arranged hyphae heavily encrusted with a cinnamon-brown pigment (revived in KOH), beneath this occur scattered enlarged hyphae interwoven with typical filamentous strands but no true hypoderm differentiated; flesh proper hyaline in KOH; universal veil remnants hyaline to sordid yellowish in KOH, filamentous.

e more es. It nat the tly on

(Fig. ved in ttened, mooth, KOH, walled, great-nt and ending not re-yphae; trama icicle of al veil

group, mown. ith the spores

1895. y comblack e only vall in pasidia raphypleuroapparn; gill s fila-

istinct ner & pecies. Discussion. This is a most unusual species with a veil like that of C. myceliocephalus M. Lange (p. 742), but apparently not closely related to it. It very likely belongs in the C. phaeosporus group though the fibrillose cuticle is unusual even there. Its spores are among the smallest I have seen in the genus.

COPRINUS PSEUDORADIATUS Kühner and Josserand, Bull. Soc. Myc. Fr. 60: 26. 1944. (Fig. 13.) Pileus 3–5 mm. high, 3–4 mm. broad when expanded, expanding to plane and 5–7 mm. broad, at first completely covered by a dense hairy-fibrillose universal veil which breaks up into white squarrose squamules and soon disappears, surface grayish white before spores mature, soon becoming lead gray to blackish, wrinkled-striate to the smooth disc, soon splitting radially; flesh very delicate; lamellae free, white, becoming jet black before deliquescing, close, edges white-fimbriate; stipe 2–3 cm. long, 0.5 mm. in diam., equal, white at first, the lower part with recurved fibrillose squamules from the remains of the veil, glabrescent, base slightly tomentose.

Spores black in deposit,  $7-8.4 \times 4-4.5 \mu$ , blackish revived in KOH, ellipsoid, not flattened, apical hyaline pore small and inconspicuous; basidia four-spored; pleurocystidia and cheilocystidia similar,  $40-60 \times 10-18 \mu$ , ellipsoid to cylindric; fibrils of the veil of long-cylindric cells  $60-90 \times 12-18 \mu$ , next to the cap surface

the cells often short and ovoid.

Discussion. I first collected this fungus in Nova Scotia, July 27, 1931, in Colchester County. However, only a few caps were collected and a description was withheld pending the collection of more and better specimens. In the meantime Kühner and Josserand discovered it in France. During the summer of 1946 it was collected in the vicinity of the University of Michigan's Biological Station at Douglas Lake. The North American collections have all been from rabbit dung.

COPRINUS PULCHRIFOLIUS Peck, Ann. Rept. N. Y. State Mus. 29: 41. 1878. (Fig. 16-type.) Spores  $7-8.4\times4-4.2\,\mu$ , reddish brown, becoming gray in KOH, ellipsoid in face view, slightly curved in side view or merely straight on ventral line and convex on dorsal line, apical hyaline pore distinct; basidia  $6-7\,\mu$  in diam., four-spored, hyaline in KOH; paraphyses inflated and hyaline in KOH; pleurocystidia none found (careful search made); cheilocystidia none seen (gill edges had apparently deliquesced); gill trama hyaline in KOH or slightly yellowish where not revived well, cuticle of globose, hyaline, in-

flated cells; universal veil remnants also of hyaline, more or less globose-inflated cells readily separable from each other.

Discussion. This is one of the Coprini closely related to C. radians. A final disposition of it must await a critical study of the group based on numerous collections. I have been able to recognize both species in the vicinity of Ann Arbor.

COPRINUS QUADRIFIDUS Peck, Ann. Rept. N. Y. State Mus. **50**: 106. 1897. (Fig. 17–type.) Spores smooth, dark chocolate color revived in KOH, 7.5–9.5 (10.5) × 4–4.5 (5.5)  $\mu$ , very slightly compressed to terete, ellipsoid in side view, slightly ovoid in face view, apex truncate from a hyaline pore; basidia four-spored (12) 14–16 (18) × 6.5–7.5  $\mu$ , hyaline, trimorphic; paraphyses 9–11 × 8–10  $\mu$ , vesiculose, hyaline, readily collapsing; pleurocystidia abundant, subcylindric, 100–150 × 20–35  $\mu$ , hyaline, thinwalled, readily collapsing, at times extending across the gill cavity; cheilocystidia present at first and 50–80 × 15–25  $\mu$ , subellipsoid, soon collapsing; gill trama hyaline in KOH; pileus trama hyaline, cuticle of a layer of hyaline vesiculose cells several cells deep, veil remnants of filamentous hyphae 6–10  $\mu$  in diam., clamp connections present.

Discussion. This is one of our best known American species. In none of my collections have spores with a ventral hump or pore been present, and the species is common locally in June. In C. ebulbosus this character is quite pronounced. Aside from this one character both appear referable to a single species. The situation here parallels that found in C. atramentarius. See C. variegatus for further comments.

COPRINUS ROTUNDISPORUS Peck, Ann. Rept. N. Y. State Mus. 31: 35. 1879. (Figs. 18–19–type.) Spores 8–10 × 6–7 × 7–9  $\mu$ , black when first revived in KOH but the dark pigment of the exospore soluble in KOH and gradually dissolving in the mount to a bister solution leaving the spore with a dark bister endospore and a practically hyaline exospore, in side view subellipsoid, slightly flattened, in face view broadly ovate when first revived but when only endospore is colored appearing subcircular, apical hyaline pore broad and distinct, exospore separable from endospore if mounts in KOH are slightly crushed under a cover glass; basidia four-spored, 15–22 × 5–6  $\mu$ , hyaline in KOH, difficult to demonstrate and apparently either not projecting or only dimorphic; paraphyses greatly enlarged, 16–20 × 10–18  $\mu$ , globose

eil like tly not sporus e. Its

inconystidia ne veil surface a, July s were

ved in

s were ion of d Jos-946 it s Bioections

Mus. eddish ightly onvex diam., ine in cheilo-; gill ightly e, into ellipsoid, apparently more or less filled with a mucilaginous substance, not collapsing but instead reviving exceptionally well in KOH and very distinct by the manner in which the transmitted light is refracted; pleurocystidia scattered, not coprinoid but instead typically fusoid-ventricose as in most agarics and highly refractive like the paraphyses,  $28-36 \times 9-14 \mu$ , apices obtuse; cheilocystidia none seen; gill trama hyaline in KOH and reviving well, highly refractive, with a narrow central strand consisting of a few filamentous hyphae, these flanked on either side by subglobular to elliptic highly refractive cells resembling the paraphyses, cap obviously splitting down the backs of the gills along the layer formed by the filamentous hyphae; pileus trama nostly of highly refractive, large, well-revived hyaline (in KOH) cells, cuticle of narrow (4-7 \mu) radially arranged hyphae (which may possibly have been veil remnants) also highly refractive and hyaline in KOH; veil remnants filamentous and not sharply distinguishable from those forming the apparent cuticle; no clamp connections seen.

Discussion. This fungus has so many curious microscopic characters for a Coprinus that it should be easy to recognize it in spite of the lack of sufficient data on the macroscopic features. The behavior of the pigment in the exospore is unique, the type of pleurocystidium very unusual, and the refractive nature of the paraphyses anomalous.

COPRINUS SEMILANATUS Peck, Ann. Rept. N. Y. State Mus. 24: 71. 1872. (Fig. 21-type.) Spores  $12.5-15\times7-8.4\times9-12.5\,\mu$ , nearly coal black as revived in KOH, flattened, subellipsoid in side view, obscurely angular and broadly elliptic to subcircular in outline in face view, hyaline pore apical and inconspicuous or projecting slightly to form a snoutlike apex; details of gill trama and hymenium not obtainable; pileus trama hyaline in KOH; universal veil remnants of thin-walled readily collapsing globose to barrel-shaped cells.

Discussion. A member of the Coprinus niveus group, but a species rather easily distinguished in nature. Both the spore characters and the veil place it here.

COPRINUS SEYMOURII Peck, Ann. Rept. N. Y. State Mus. 28: 49. 1876. (Fig. 20-type.) Spores 6.2– $7.8 \times 3.1$ – $3.5 \times 4$ – $4.6 \mu$ , dull rusty bister in KOH, changing to chocolate-color, flattened, subellipsoid in side view, obscurely triangular in face view, hyaline

pore apical and distinct; basidia hyaline in KOH, four-spored, 5–6  $\mu$  in diam.; paraphyses hyaline, inflated, readily collapsing; pleurocystidia none seen (type had been pressed and gills did not revive well); cheilocystidia none seen; gill trama not reviving; pileus trama with a cuticle of vesiculose cells, the walls thickened in the angles at the base and brownish in KOH.

Discussion. A member of the C. micaceus series. I have a number of collections which have been referred here, but as yet am not satisfied that the species is more than an extreme variant of C. micaceus. If the habitat is truly terrestrial, that should be an aid in distinguishing it, but C. micaceus can be very deceiving in this regard.

COPRINUS SILVATICUS Peck, Ann. Rept. N. Y. State Mus. 24: 71. 1870. (Fig. 22-type.) Pileus 1-2 cm. high in buttons, 0.5-2.5 cm. across the base when expanding, ellipsoid to subovoid in buttons and then densely pubescent from cystidia, wrinkled-striate to disc but striae extending unequal distances, disc ochraceous tawny to tawny or this color over all when young, striate portion soon ochraceous buff to pale ochraceous buff and very atomate, disc glabrous at maturity, marginal area finally livid and splitting as the spores are shed; flesh very thin and fragile, no odor or taste; lamellae equal, attached at apex of stipe, moderately broad, close to crowded, pallid young, darkening over all before starting to deliquesce, edges white-fimbriate at first; stipe short, 1.5-2.5 cm, long, 1-2.5 mm, thick, equal, margin of cap attached at base but bulb if present soon evanescent (but expansion as in Bulbopodium of Cortinarius), densely pubescent at first, finally glabrous, pallid over all or at base slightly sordid.

Spores  $11.5-14 \times 7-8.4 \times 8-9 \,\mu$ , slightly broader in face view than in side view (slightly compressed), slightly inequilateral in side view, subovoid in face view, apical pore broad, hyaline, prominent and causing apex to appear more or less snoutlike, surface slightly wrinkled from a thin hyaline somewhat deciduous exospore, dark bister to blackish revived in KOH; basidia fourspored, hyaline in KOH, trimorphic; paraphyses inflated, hyaline in KOH; pleurocystidia and cheilocystidia none seen (sections revived poorly); gill and pileus trama dark brown when first mounted in KOH but rapidly clearing to sordid yellowish, cuticle of pileus of vesiculose cells about one cell deep, numerous hyaline, thin-walled, subcylindric to basally ventricose pilocystidia 80–120  $\times$  10–18  $\mu$  arising from this layer, their necks greatly elongated

and  $10-15 \mu$  in diam., their apices rounded to obtuse.

chare it in atures. e type

of the

us sub-

well in

smitted

but in-

highly

obtuse:

eviving ting of

y sub-

raphy-

ng the

stly of

cells.

h may d hva-

distin-

p con-

Mus. 4 × 9-bellipbellipincondetails yaline apsing

but a spore

s. 28:  $4.6 \mu$ , tened, yaline

Discussion. The description of the macroscopic characters is taken from numerous collections made by Conners, Groves, and Smith in the vicinity of the laboratories at the Petawawa Experiment Station near Deep River, Ontario, during the season of 1947. In this material the basidia varied from two- to four-spored and the spores  $14-20\times7.5-10\,\mu$ . The paragraph on the microscopic data given here was taken from the type. The species stands out as very distinct in the *C. ephemerus* series.

COPRINUS SPRAGUEI Berk. & Curt. Ann. and Mag. Nat. Hist. III 4: 292. 1859. (Fig. 23.) Spores bister in KOH, 9–11  $\times$  5.5–6.5  $\times$  6.5–7.8  $\mu$ , slightly flattened, suboblong in side view, in face view obscurely angular-ovate, base broadly pointed, the hyaline pore distinctly eccentric but small and inconspicuous; basidia hyaline in KOH, four-spored, 18–32  $\times$  6–7.5  $\mu$ ; paraphyses 14–18  $\times$  9–13  $\mu$ , hyaline and inflated; pleurocystidia and cheilocystidia none seen; gill trama hyaline in KOH; pileus trama homogeneous beneath a cuticle formed by a palisade of more or less ellipsoid cells, flesh tawny in KOH just beneath the cuticle in thick sections (material revived poorly and KOH test not truly distinctive), no pilocystidia seen.

Discussion. The microscopic data are taken from a collection Murrill compared with the type at Kew and pronounced identical with it. The species is clearly in the C. plicatilis series and distinct because of its smaller spores.

COPRINUS VARIEGATUS Peck, Bull. Buffalo Soc. Nat. Sci. 1: 54. 1873. (Fig. 24–type.) Spores 7–9 (10) × 5–6.2  $\mu$ , subellipsoid to ovoid, not flattened, often with a ventral hump or blister (germ pore?), dull cocoa-brown when first revived in KOH, gradually fading to near avellaneous or "wood brown" (possibly immature), apical hyaline pore distinct; basidia four-spored, hyaline in KOH, 14–22 × 5–7  $\mu$ , clavate to subcylindric on a short pedicel, trimorphic; paraphyses hyaline, thin-walled, readily collapsing; pleurocystidia cylindric, hyaline, thin-walled, readily collapsing, 60–120 × 15–28  $\mu$ ; cheilocystidia none seen; gill trama hyaline in KOH; pileus trama hyaline in KOH, cuticle of narrow (4–8  $\mu$ ) somewhat gelatinous hyphae as revived in KOH.

Discussion. This species is apparently easily distinguished from C. ebulbosus and C. quadrifidus by the filamentous somewhat gelatinous cuticle. However, further studies on either fresh or very carefully dried specimens of all three are needed to establish this

difference as a valid character. The general aspect of the fruiting bodies and the type of spore remind one of *C. ebulbosus* and *C. quadrifidus*. The white rhizomorphs and peronate-annulate stipe may be additional diagnostic characters.



Figs. 23-31. Microscopic characters of dark-spored agarics.

and peri-947.

and

out

Hist.

1 ×

7, in

hya-8 × none eous soid ions , no

tion tical dis-

soid erm nally re), OH, tribleu-60e in 8 μ)

rom gevery this Coprinus subpurpureus sp. nov. (Fig. 25-type.)

Pileus 1.5 cm. altus, demum 2–3.5 cm. latus, pubescens, purpureo-brunneus vel subpurpureus; lamellae anguste adnatae, confertae, pallidae demum nigrae; stipes 4–10 cm. longus, 1–2.5 mm. crassus, cavus fragilissimus, lilaceo-umbrinus demum pallidus, pubescens; sporae 12–14  $\times$  5.5–6.8  $\times$  7–8  $\mu$ .

Pileus 1.5 cm. high, up to 3.5 cm. broad when expanded, finely pruinose-pubescent when young, soon glabrous as cystidia collapse, "natal brown" to "Benzo brown" on disc, "light brownish drab" toward the paler ("light cinnamon drab") broad, striate, marginal area, in age "dark purple drab" over disc and dark gray to blackish over margin; flesh very thin and fragile, delicate, no odor or taste; lamellae narrowly adnate, close, becoming subdistant, narrow, near "tilleul buff," darkening to black over all before the edges deliquesce; stipe 4–10 cm. long, 1–2.5 mm. thick, equal, dull lilac umber young, pallid in age, densely pruinose-pubescent but soon

glabrescent, base white-strigose.

Spores black in deposits, 12– $14 \times 5.5$ – $6.8 \times 7$ – $8 \mu$ , smooth, germ pore eccentric, in face view spores with a ventral pore visible (but not humped as in *C. ebulbosus*), narrowly subovate to elliptic in face view, with a suprahilar depression in side view; basidia fourspored, tetramorphic, 18– $30 \times 6$ – $7 \mu$ ; paraphyses hyaline, inflated, not otherwise distinctive; pleurocystidia none seen; cheilocystidia vesiculose to broadly fusoid-ventricose, 10– $30 \mu$  in diam.; gill trama very thin and appearing cellular in sections, slightly colored vinaceous brown in KOH (young gills); pileus trama thin and appearing to be a mixture of narrow filaments and enlarged cells, cuticle a single layer of vesiculose to pedicellate cells from which thin-walled pilocystidia 60– $100 \times 10$ – $16 \mu$  project, layer beneath cuticle lilaceous-brown to sordid vinaceous brown in KOH, clamp connections present.

Gregarious on wet leaves in a springy area, under hardwoods, Colonial Point, Burt Lake, Cheboygan County, Michigan, July 31, 1947. Collected by Margaret Feigley (A.H.S. No. 26158—type).

Discussion. This is a very interesting species of the C. ephemerus series. The long narrow spores with the eccentric pore, the purplish color particularly in the stipes of young fruiting bodies, and habitat distinguish it from any other Coprinus known to me.

### **PANAEOLUS**

Panaeolus anomalus (Murrill) Saccardo & Trott. Syll. Fung. 23: 323. 1925. (Figs. 26, 27-type.) Spores 11–13 (14)  $\times$  6–

 $7 \times 7-9\,\mu$ , flattened, dark brown (darker than "warm sepia") when revived in KOH, narrowly subovate to elliptic in side view, broadly ovate in face view, some lopsided and some obscurely angled, apical hyaline pore present and distinct; basidia hyaline in KOH, four-spored,  $18-22 \times 8-9\,\mu$ ; paraphyses if present remaining collapsed; pleurocystidia abundant,  $60-80 \times 10-20\,\mu$ , apices acute, the walls thickened particularly in the apices which are usually solid, deep yellowish brown in KOH, broadly fusoid to fusoid-ventricose, smooth; cheilocystidia similar to pleurocystidia but smaller (all hyaline thin-walled cells remained collapsed); gill trama sordid yellowish in KOH; pileus trama sordid yellowish in KOH, the cuticle of vesiculose cells and scattered fusoid-ventricose thin-walled hyaline pilocystidia scattered between the vesiculose cells.

Discussion. This species was described as Campanularis anomalus by Murrill (Mycologia 10: 32. 1918). Some authors place the species of Panaeolus with brown thick-walled pleurocystidia in a separate genus, Copelandia, but in my estimation the group is better regarded as a section of Panaeolus.

**Panaeolus castaneifolius** (Murrill) comb. nov. (Fig. 28-type.) Spores  $12-16\times7-8.5~\mu$ , somewhat almond-shaped, obscurely to distinctly verrucose, dark tawny to pale russet when revived in KOH, hyaline apical germ pore present but small and inconspicuous under ordinary magnifications; basidia  $24-28\times10-12~\mu$ , hyaline in KOH, four-spored; pleurocystidia present as dark cinnamon-brown basidia-like or narrower bodies embedded in the hymenium,  $18-24\times6-10~\mu$ ; cheilocystidia abundant, fusoid-ventricose to subcylindric,  $24-38\times7-10~\mu$ , neck often flexuous and apices usually obtuse, thin-walled and hyaline; gill trama parallel to subparallel, dull cinnamon-brown when first revived in KOH; pileus trama with a cuticle of clavate, pear-shaped and vesiculose cells arranged into a somewhat irregular palisade, flesh proper floccose, interwoven and pallid to pale brownish when revived in KOH.

Discussion. Microscopically this species is almost identical with Panaeolus foenesicii, but the stipe, which is 4-6 mm. thick, the strong odor and unpleasant taste should distinguish it readily in the field. Murrill originally published it as Psilocybe castaneifolia (Mycologia 15: 17. 1923).

nneus grae ; aceo-

inely apse, rab" ginal kish aste; near delililac soon

erm

(but

ourated, tidia gill ored and tells, hich eath

ods, 31, pe). em-the

ing.

6-

lies,

me.

## Panaeolus fraxinophilus sp. nov. (Figs. 29-31-type.)

Pileus 8-15 mm. latus, conicus demum convexus, siccus et canescens demum udus et hygrophanus, fuscus vel niger, ad marginem Isabellinus; lamellae confertae, latae, cinereae demum nigro-maculatae; stipes 1-2 cm. longus, 1-2 mm. diam., cavus, fragilis, pruinosus, griseo-brunneus; sporae 9-11  $\times$  5-6  $\times$  6.5-7.5  $\mu$ . Ad. truncos Fraxini.

Pileus 8–15 mm. broad, conic to convex, surface appearing dry and hoary, typically becoming moist and hygrophanous, margin faintly translucent-striate and incurved at first, "fuscous black" to "fuscous" at first, nearly black on disc but soon "Benzo brown," "hair brown" to "drab" (dark gray), at or near margin "tawny olive" to "Isabella color" (sordid yellowish brown); flesh thin, dark, odor slightly farinaceous, taste mild, lamellae, close, moderately broad, adnate, dull drab becoming black-spotted, edges whitish; stipe 1–2 cm. long (up to 8 cm. around stumps), 1.5–2 mm. diam., hollow, fragile, equal, dark grayish brown with a tinge of red, densely pruinose over all and beaded with drops of moisture.

Spores 9–11 × 5–6 × 6.5–7.5  $\mu$ , smooth, slightly flattened, narrowly elliptic in side view, ovate in face view, dark reddish brown in KOH, black in mass, apical hyaline pore very small but distinct; basidia four-spored, a few two-spored, hyaline in KOH, 18–22 × 9–10  $\mu$ , clavate; paraphyses not differentiated; pleurocystidia none; cheilocystidia very abundant, filamentous and apparently with mucilaginous walls, hyaline, many crooked, 28–42 × 4–5  $\mu$ ; gill trama regular, pale dull brown in KOH; pileus trama with a dull brown region beneath the cuticle, remainder paler, cuticle of a layer of vesiculose hyaline cells one to two cells deep, numerous hyaline ventricose to subcylindric pilocystidia projecting and measuring 20–44 × 5–6  $\mu$ , filamentous in age.


On naked trunk of a fallen ash tree, Warrensburg, New York, September, 1934, A. H. Smith 778-type.

Discussion. The combination of relatively small spores and habitat distinguish it from any Panaeolus known to me.

Panaeolus reticulatus Overholts, Ann. Mo. Bot. Gard. 3: 195. 1916. (Fig. 32–type.) Spores 9–11 (12.5)  $\times$  5.6–6.2  $\times$  6.5–8  $\mu$ , slightly flattened, subelliptic to slightly inequilateral in side view, in face view ovate or a few very obscurely angular, dark bister revived in KOH, apical hyaline pore small but distinct; basidia four-spored, hyaline when projecting but the collapsed hymenium and subhymenium bister in KOH; pleurocystidia none; cheilocystidia fusoid-ventricose with obtuse apices, 18– $26 \times 8$ – $19 \mu$ , gill trama bister in KOH; pileus trama bister, cuticle not

reviving, apparently of collapsed vesiculose cells. The tissue of the gills and pileus revived poorly.

Discussion. The spores separate it easily from P. retirugis and the gray to umber colors from P. subbalteatus.



Figs. 32-38. Microscopic characters of dark-spored agarics.

mum nellae ngus, < 5–6

dry
rgin
" to
wn,"
wny
thin,
derwhitmm.
ge of
ture.
narown
inct;

one; nurama rown er of aline uring

ork,

and

1. 3: i.2 × al in dark inct; l hyione;

× 8– not

PANAEOLUS RUFUS Overholts, Ann. Mo. Bot. Gard. 3: 196. (Figs. 33, 34-type.) Spores  $12-14 \times 6.5-8 \times 7-9.5 \mu$ , slightly flattened, subovoid to subelliptic in side view, broadly ovate in face view, dark bister in KOH, hyaline pore apical to very slightly eccentric and inconspicuous; basidia hyaline in KOH (some collapsed basidia brownish), four-spored or a few twospored and then sterigmata abnormal, 20-24 × 10-12.5 μ; paraphvses apparently coprinoid but readily collapsing and basidia so numerous as to obscure them (hymenium in thick sections pale bister); pleurocystidia none; cheilocystidia abundant, hyaline in KOH, thin-walled, fusoid-ventricose to subvesiculose or subcylindric with a broadly rounded slightly enlarged apex, 26-42 × (8) 10-14 μ, in age becoming elongated and with flexuous walls; gill trama parallel, the cells broad and long  $(60-120 \times 15-20 \,\mu)$ , pale bister near and in subhymenium or finally sordid vellowish as revived in KOH; pileus trama sordid yellowish in KOH, cuticle of somewhat enlarged cells but not sharply distinct (as revived) from the remainder of the flesh.

Discussion. This species appears to me to be a synonym of Panaeolus subbalteatus as the latter is now understood. The only key character to separate them is the farinaceous odor and taste.

Panaeolus semiovatus (Fries) Lundell, Fung. Exsic. Upsal. No. 537. 1937. Spores 16–22 (23) × 8.5–11  $\mu$ , smooth, apices truncate from a hyaline pore, ellipsoid in side view, slightly lemonshaped to obscurely angled in face or back view, black in mass and dark blackish brown under microscope; basidia four-spored, 32–34 × 12–14  $\mu$ ; pleurocystidia rare to scattered, clavate to saccate or sometimes mucronate and with an irregular highly refractive content as revived in KOH (cystidia naematolomoid), 28–36 × 10–18  $\mu$ ; cheilocystidia similar to pleurocystidia or merely fusoid-ventricose and with homogeneous content, thin-walled, hyaline, 26–38 × 7–12  $\mu$ , apices obtuse, walls sometimes flexuous; gill trama apparently regular (not reviving well); pileus trama with a cuticle of inflated hyaline cells several cells deep but very soon gelatinous and their outlines difficult to ascertain.

Discussion. Strictly speaking this fungus is not within the scope of this paper, but is included in order that its microscopic details may be more readily discussed in relation to those of other species. It is evident, from a consideration of the aspect of the fruiting bodies and the similarity of the microscopic characters, that this fungus and P. solidipes Peck are very closely related.

The similarity in the pleurocystidia of each emphasizes this point. *P. semiovatus* was placed in a separate genus, *Anellaria* (*A. separata* of Karsten), primarily because of the annulate stipe. Lange placed it in *Stropharia*. In view of the similarity in all characters except the annulus between *P. semiovatus* and *P. solidipes* it seems more sensible to me to retain both in the same genus and group them together in a distinct section. A veil is present in many species of *Panaeolus* but in most the remains adhere to the margin of the cap when the latter expands.

Panaeolus solidipes Peck, Ann. Rept. N. Y. State Mus. 23: 101. 1872. (Figs. 35–36–type.) Spores (12.5) 14–17 × 6–8 × 8–11  $\mu$ , flattened, narrowly subelliptic in side view, broadly ovate to obscurely angular-elliptic in face view, very dark bister to blackish in KOH, apical pore hyaline but very small and inconspicuous; basidia four-spored; hyaline in KOH, 20–26 × 12–13  $\mu$ ; paraphyses not differentiated; pleurocystidia scattered to abundant, 36–54 × 10–20  $\mu$ , clavate from a narrow or a short, broad pedicel, hyaline in KOH and with an amorphous mass of a highly refractive substance in the enlarged portion, thin-walled and readily collapsing; cheilocystidia vesiculose, 18–24 × 9–14  $\mu$ , also with a highly refractive content revived in KOH; gill trama regular, sordid yellowish as revived in KOH, pileus trama hyaline to sordid yellowish; cuticle of a single row (in sections) of hyaline inflated cells.

Discussion. See P. semiovatus.

196.

.5 µ.

adly

l to

OH

wo-

1 50

pale

e in

(8)

gill

pale

as

ticle

red)

r of

only

iste.

sal.

ices

on-

and 32-

cate

tive

6 X

oid-

ine,

gill

vith

oon

the

opic

ther

the

ers,

ted.

PANAEOLUS VARIABILIS Overholts, Ann. Mo. Bot. Gard. 3: 197. 1916. (Figs. 37–38-type.) Spores  $11-12 \times 6-6.6 \times 7.5-9 \mu$ , flattened, subovate to slightly inequilateral in side view, ovate in face view, smooth, not obscurely angular, bister to dark bister as revived in KOH, hyaline apical pore distinct but not large; basidia hyaline in KOH, four-spored (a few with smoky brown sterigmata and some with brownish walls—the latter collapsed),  $20-24 \times 10$  $12 \mu$ ; paraphyses not distinctly differentiated until late maturity and then readily collapsing (young basidia also somewhat saccate); pleurocystidia none; cheilocystidia abundant, hyaline and thin-walled, fusoid-ventricose to subcylindric,  $26-35 \times 6-10 \,\mu$ , elongated and crooked in age, apices obtuse to subcapitate; gill trama bister in KOH; pileus trama with a cuticle of inflated cells about one or two cells deep, these soon collapsing and difficult to demonstrate, flesh proper hyaline to sordid yellowish as revived in KOH.

Discussion. P. variabilis most closely resembles the fungus currently known as P. papilionaceus, but differs in smaller spores. It needs further study.

Panaeolus venenosus Murrill, Mycologia 8: 186. 1916. (Figs. 39, 40–type.) Spores 10–12.6 (13) × 6–6.5 × 7–8.5  $\mu$ , dark bister revived in KOH, smooth, flattened, subovoid to elliptic in side view, not obscurely angular, apical hyaline pore small but distinct; basidia hyaline in KOH when isolated, brownish in thick sections, 18–24 × 7.5–8.5  $\mu$ , clavate; paraphyses not distinctive; pleurocystidia present only near gill edges and similar to cheilocystidia; cheilocystidia fusoid-ventricose, 26–38 × 6–10  $\mu$ , with a long, often crooked neck above a slightly ventricose basal portion, hyaline or bases brownish in KOH, thin-walled, apices capped with mucilage in some; gill trama bister in KOH, gradually becoming paler; pileus trama dark bister but gradually paler, cuticle formed by a palisade of clavate-pedicellate cells the bases of which are yellowish in KOH.

Discussion. In my estimation this species is not sufficiently distinct from P. subbalteatus to be recognized. The marginal belt in P. subbalteatus is not a constant character, in fact it is not visible on fresh entirely moist caps or on those completely faded. The spores in Murrill's type are slightly smaller than my measurements for P. subbalteatus, but the difference is too slight to be considered important until established from numerous spore deposits.


# Panaeolus fontinalis sp. nov. (Figs. 41, 42-type.)

Pileus 1-2 cm. latus, obtusus demum late conicus, impolitus vel pruinosus, griseo-olivaceus demum olivaceo-brunneus; lamellae pallide olivaceae demum cinereae et maculatae, confertae, latae, adnatae; stipes 5-10 cm. longus, 1-2 mm. crassus, fragilissimus, pallidus, deorsum argillaceus; sporae  $7-9\times4-5\times5-6.5~\mu$ 

Pileus 1–2.5 cm. broad, obtusely conic, expanding to broadly conic, surface unpolished to pruinose, "deep olive buff" over disc, paler toward margin, with a "buffy olive" marginal belt, finally "olive brown" or darker over all, fading eventually to "light grayish olive," with a silky appearance in age; flesh very thin and pallid, odor not distinctive; lamellae near "olive buff" young, soon drab, finally mottled with black, ascending adnate, close, broad, usually with 2 tiers of lamellulae, edges at first whitish and slightly floccose; stipe 5–10 cm. long, 1–2 mm. diam., equal, very fragile,

pallid and densely pruinose at apex, darker downward and "cinnamon buff" to "clay color" or tawny olive, pruinose over all at first and beaded with drops, naked in age.

Spores black in deposits, dark bister in KOH,  $7-9 \times 4-5 \times 5-6.5 \mu$ , smooth flattened, broadly ovate in face view, nearly elliptic



Figs. 39-48. Microscopic characters of dark-spored agarics.

ngus ores.

1916. 3.5  $\mu$ , liptic l but thick tive; reilotith a tion, pped benticle which

belt visiided. sureo be pore

osus, mum , 1–2 < 4–5

adly disc, nally light and soon oad, htly gile, in side view, pore apical and distinct; basidia four-spored, 18–24  $\times$  7–8.4  $\mu$ , hyaline in KOH; pleurocystidia absent to scattered (often abundant where insects have damaged hymenium), similar to cheilocystidia; cheilocystidia fusoid-ventricose to subcylindric, 26–34  $\times$  6–9  $\mu$ , apices obtuse, neck often flexuous; gill trama pale bister in KOH, the elements more or less interwoven; pileus trama darker than the gill trama, cuticle of a single layer of vesiculose cells, pilocystidia present and similar to cheilocystidia or filamentous and 15–30  $\times$  4–6  $\mu$ , hyaline.

Scattered on black muck among liverworts and a species of *Mnium* in a springy area in a cedar swamp, Burt Lake, Cheboygan Co., Mich., during July, 1947. July 4, 25437; July 5, 25493, 25494 and 25495. Collection 25437 is designated the **type**.

Discussion. The habitat of this Panaeolus is very unusual. A careful search was made for rabbit dung or that of other small animals, but none was found. The habitat, very small spores, and olive buff gills characterize the species.

## NAEMATOLOMA

Naematoloma campestre sp. nov. (Figs. 43-45-type.)

Pileus (6) 8-15 (20) mm. latus, obtuse conicus demum campanulatus vel plano-convexus, glaber, lubricus, subcastaneus demum pallide ferrugineus; lamellae latae, adnatae, subdistantes, subochraceae; stipes brevis, 2-3 cm. longus, 1-1.5 mm. crassus, subfibrillosus, deorsum fuscescens; sporae 7.8-9.3 (10)  $\times$  4.6-5  $\times$  5-6.7  $\mu$ .

Pileus (6) 8-15 (20) mm. broad, obtusely conic, expanding to broadly conic or subcampanulate, finally convex to nearly plane, surface glabrous or with a faint fringe of marginal fibrils when young, lubricous when moist, color "Vernona brown" to "warm sepia" or "auburn" to "pecan brown" on disc, fading first to "pale pinkish buff" or "cinnamon buff" (pale ferruginous), but in age darkening to pale tan, margin striate when moist; flesh pliant, thin, odor very faintly fragrant, taste mild to slightly bitterish; lamellae very broad, horizontally adnate with a decurrent tooth, readily seceding, subdistant (15-18 reach stipe), one tier of lamellulae, dull brown becoming near "snuff brown" to very sordid cinnamon-brown, edges even and not whitish; stipe short, 2-3 cm. long, 1-1.5 mm. thick, equal, cartilaginous-pliant, about "light pinkish cinnamon" over all at first from thin remains of light pinkish cinnamon veil, soon darkening from base upward to concolor or darker than the cap, only somewhat glabrescent.

Spores 7.8–9.3 (10)  $\times$  4.6–5  $\times$  5–6.7  $\mu$ , flattened slightly, angular-subspheric to angular-ovate or subelliptic with sides  $\pm$  parallel and with faintly angled shoulders in face view, subelliptic in side view, pore apical but small, pale bister as revived in KOH; basidia four-spored, hyaline in KOH, 18–23  $\times$  7–8  $\mu$ , subcylindric; paraphyses basidioid; pleurocystidia nearly embedded in hymenium and originating from gill trama or subhymenium, ventricosemucronate to fusoid-ventricose and with a highly refractive amorphous body as seen revived in KOH (*Naematoloma* type), 26–34  $\times$  9–14  $\mu$ ; cheilocystidia hyaline, thin-walled, homogeneous, 18–26  $\times$  4–7  $\mu$ , subcylindric to subfusoid-ventricose with obtuse apices; gill trama of subparallel hyphae, pale to rather dark bister as revived in KOH; pileus trama dark bister to sordid cocoa-color in KOH, homogeneous (no differentiated hypoderm or pellicle).

Gregarious on sod in pastures where Agaricus campestris is frequently found, definitely not on dung. May 24, 1946, Smith 21426-type.

Discussion. The spore characters and habitat are more like Psilocybe, but the pleurocystidia are typical of Naematoloma.

Naematoloma humidicola (Murr.) comb. nov. (Figs. 46-Spores 9-11  $\times$  5.5-6.3  $\mu$ , ovoid in front or back view, ellipsoid to slightly inequilateral in side view, pale tawny in KOH, smooth, apical pore present but very indistinct; basidia hyaline to pale yellow in KOH, four-spored, subcylindric,  $23-27 \times 5-6.5 \mu$ ; pleurocystidia scattered, yellow in KOH, fusoid ventricose to subclavate-mucronate, (44) 50-58 (64)  $\times$  8-10 (12)  $\mu$ , most of them with a highly refractive content, some with what appears to be a wrinkled inner wall, projecting 25-30 μ beyond the basidia; cheilocystidia 24–28 (32)  $\times$  5–8  $\mu$ , ventricose to subcylindric, hyaline, content homogeneous, thin-walled, abundant; gill trama subparallel, the cells equal in width throughout their length, pale yellow in KOH, subhymenium cellular but very thin and indistinct, pileus trama with a thin pellicle of narrow gelatinous hyphae 2.5-4  $\mu$  in diam.; beneath the pellicle is a region of enlarged cells (the hypoderm), the remainder of the flesh filamentous-floccose, all parts pale yellow in KOH, clamp connections present.

Discussion. Murrill described this species as Naucoria humidicola (North Am. Flora 10: 174. 1917). However, its natural affinities are unquestionably in the vicinity of Naematoloma elongatipes. It is distinguished among the slender species of Naematoloma by the conspicuous, elongated (for this genus) pleuro-

ama llose nen-

3-24

ered

nilar

dric,

pale

rgan 493, A

and

vel eus; cm. –9.3

to

ne, nen rm vale age ant, sh;

of ery ort, out

of

to

cystidia. In most species in this genus the cystidia do not project appreciably beyond the hymenium.

Naematoloma petasiforme (Murrill) comb. nov. (Figs. 49-Spores 8-10  $\times$  5-6  $\mu$ , ellipsoid to subovoid, smooth, smoky purple-brown under the microscope when fresh (Murrill), sordid to pale tawny when revived in KOH, with an apical hvaline germ pore: basidia four-spored, subcylindric to clavate. hyaline in KOH,  $20-24 \times 7-8 \mu$ ; pleurocystidia  $20-30 \times 9-12 \mu$ , mucronate, clavate or fusoid-ventricose and with a highly refractive body of amorphous material in enlarged part; cheilocystidia similar to pleurocystidia or fusoid-ventricose and 20-32  $\times$  8-11  $\mu$ with a hyaline homogeneous content, the former rare, the latter very abundant and also occurring near gill edge as pleurocystidia; gill trama regular, tawny brown in KOH, hyphae somewhat interwoven and with encrusted pigment; pileus trama with a cuticle of radially arranged hyphae mostly 4-6 µ in diam, and with clamp connections, the walls with encrusting pigment, some cells of the cuticle barrel-shaped and up to 15 µ in diam. (but no true cuticle of vesiculose cells visible in type), flesh proper floccose and golden tawny in KOH.

Discussion. Murrill described this as Psathyrella petasiformis (Mycologia 14: 276. 1922). It differs from other members of the Naematoloma dispersum series by its lack of a thin pellicle. There are other less striking differences such as the slightly larger spores and possibly a darker spore deposit. However, these cannot be given much emphasis until comparative studies of fresh material can be made and spores from deposits compared.

### **PSILOCYBE**

**Psilocybe bulbosa** (Pk.) comb. nov. (Figs. 52, 53–type.) Spores 6–7.5 (8)  $\times$  3.8–4.2  $\times$  4–4.8  $\mu$ , angular-ovate in face view, pointed at base, slightly inequilateral to subovoid in side view, flattened, dull yellowish brown when revived in KOH, with an obscure apical hyaline pore; basidia 20–24  $\times$  6–7  $\mu$ , four-spored, hyaline in KOH; pleurocystidia embedded and difficult to locate, 18–24  $\times$  6–7  $\mu$ , obtusely fusoid-ventricose, hyaline in KOH, content homogeneous; cheilocystidia abundant, 20–28  $\times$  4–8  $\mu$ , similar to pleurocystidia or more elongated, when revived in KOH often with a drop of a mucilaginous substance adhering to apex; gill trama parallel, hyaline to pale brownish in KOH; pileus trama with a thin, hyaline gelatinous pellicle of hyphae 1.5–3  $\mu$  in diam.

and bearing clamp connections, flesh proper of compactly interwoven hyphae, pallid tawny brownish in KOH.

Discussion. The distant gills, angular to ovate spores in face view, and thin gelatinous pellicle appear distinctive. I suspect the



Figs. 49-65. Microscopic characters of dark-spored agarics.

oject

. 49– ooth, rill), pical vate, 12 μ, fracstidia -11 μ atter

nterle of lamp f the aticle olden

idia;

f the here pores of be terial

ppe.)
riew,
riew,
riew,
rored,
cate,
consimiOH
pex;

ama am., pale color of the cap as described by Peck was caused by fading. The species was described as *Deconica bulbosa* by Peck (Ann. Rep. N. Y. State Mus. **46**: 107. 1893).

PSILOCYBE CAESPITOSA Murrill, Mycologia 15: 5. 1923. (Figs. 54, 55–type.) Spores 6.2– $7.5 \times 4$ – $5.2 \, \mu$ , smooth, subelliptic in side view, obscurely angular to ovate in face view, terete to very slightly compressed, pale ochraceous tawny when revived in KOH, truncate from a small apical germ pore; basidia four-spored, 16– $20 \times 6$ – $7 \, \mu$ , hyaline in KOH; pleurocystidia none; cheilocystidia abundant, 22– $28 \times 3$ – $6 \, \mu$ , narrowly fusoid, hyaline; gill trama parallel or nearly so, regular, the hyphae 4– $8 \, \mu$  in diam., sordid yellowish to hyaline in KOH; pileus trama homogeneous beneath a thin hyaline gelatinous pellicle of hyphae 2– $4 \, \mu$  in diam., flesh proper floccose and sordid yellowish in KOH, clamp connections present.

Discussion. This species is very closely related to P. subviscida Peck, but differs sharply in the spacing of the gills. Those of Peck's species are subdistant instead of crowded. P. caespitosa apparently has a more shaggy-fibrillose stipe and slightly smaller spores in addition, but the latter difference is very slight.

PSILOCYBE CASTANELLA Peck, Bull. N. Y. State Mus. 1: 7. 1888. (Figs. 56, 57–type.) Spores 6–7 × 3.5–4  $\mu$ , subellipsoid to ovoid, not flattened, with a small hyaline apical pore, dull yellowish brown revived in KOH; basidia four-spored, 18–22 × 6–7  $\mu$ , hyaline in KOH; pleurocystidia none; cheilocystidia 18–24 (28) × 3–5  $\mu$ , hyaline in KOH, subventricose to nearly cylindric, apices obtuse to subacute, apex often with an adhering drop or cap of a hyaline mucilaginous substance; gill trama parallel, yellowish brown in KOH or more tawny next to the subhymenium; pileus trama homogeneous beneath a thin, hyaline, gelatinous pellicle, remainder yellowish brown or toward the subhymenium paler.

Discussion. Psilocybe californica Earle (Bull. N. Y. Bot. Gard. 3: 301. 1904) appears, beyond the possibility of a doubt, to be this species. Its spores are also illustrated (FIG. 59). The habitat on sod appears to be distinctive as is also the habit of growing in rather dense masses. The latter feature is more characteristic of specimens from California. The cheilocystidia (FIG. 58) are also similar.

Panaeolus digressus Peck, Bull. Torrey Club 22 : 205. 1895. Spores sordid yellowish brown in KOH, 12–14  $\times$  6.8–7.8  $\times$  8–9  $\mu$ ,

slightly compressed, elliptic in side view, obscurely angled and broadly elliptic in outline in face view, hyaline apical pore small but distinct; basidia 30– $35 \times 10$ – $12.5 \,\mu$ , clavate, hyaline in KOH, four-spored; paraphyses not distinct from young basidia; pleurocystidia none, sections of hymenium pale sordid yellowish revived in KOH; cheilocystidia forming a sterile band on gill edge, hyaline, thin-walled, often crooked to contorted, typically fusoid-ventricose to subcylindric, 18– $32 \times (4)$  5– $8 \,\mu$ , apices obtuse and some capped with mucilage; gill trama pale sordid yellowish in KOH, subparallel next to subhymenium, central strand interwoven; pileus trama pale sordid yellowish in KOH except for a more tawny colored region just below the gelatinous cuticle, the hyphae of the cuticle hyaline to yellowish and 2– $3 \,\mu$  in diam., those of flesh proper 6– $8 \, (10) \,\mu$  in diam.

Discussion. A synonym of Psilocybe coprophila.

**Psilocybe fuliginosa** (Murrill) comb. nov. (Figs. 60, 61–type.) Spores  $6\text{-}7\times4.5\text{-}5.5~\mu$ , smooth, typically triangular in optical section but varying to ovoid or ellipsoid (as seen in face view), ellipsoid in side view, sordid yellowish brown when revived in KOH, with an obscure apical hyaline germ pore; basidia four-spored,  $18\text{-}20\times5\text{-}6~\mu$ , hyaline in KOH; pleurocystidia scattered to rare, similar to cheilocystidia; cheilocystidia  $16\text{-}28\times4\text{-}7~\mu$ , hyaline, narrowly ventricose, the neck flexuous and the apex acute to subacute or in the form of a small knob, many with a highly viscous substance adhering near or on the apex; gill trama parallel, sordid yellowish brown in KOH; pileus trama homogeneous; no pellicle seen, upper region darker yellowish brown in color than that near subhymenium.

Discussion. Dr. Burke has sent me material from Alabama which appears to belong here though the spores vary slightly, being somewhat longer. They range from 6.5 to  $8\mu$ . For P. fuliginosa the habitat on bare earth, the dry pileus, shape of spores in face view, and fuligineous cast of the carpophore are apparently distinctive. Murrill described it as Atylospora fuliginosa (Mycologia 10: 25. 1918).

PSILOCYBE GRAVEOLENS Peck, Bull. N. Y. State Mus. 167: 47. 1913. (Fig. 62–type.) Spores 8–10  $\times$  5–6  $\mu$ , ellipsoid in face view, in side view subelliptic to slightly inequilateral, smooth, pale sordid ochraceous tawny under the microscope in KOH, apical pore small but distinct; basidia hyaline in KOH, 22–26  $\times$  6–7  $\mu$ , four-spored; pleurocystidia rare to scattered, fusoid, 28–32  $\times$  7–

ding. Ann.

e in very OH, 16–tidia rama ordid leath flesh

scida e of itosa aller

: 7.

tions

yel-× 6– 8–24 dric, p or yelum; pelaler.

Bot. oubt, The

har-(FIG.

895. -9 μ,  $9\,\mu$ , the walls slightly brownish in some, hyaline in others when revived in KOH, buried in the hymenium; cheilocystidia hyaline,  $12\text{-}22\times7\text{-}9\,\mu$ , necks  $4\text{-}6\,\mu$  broad and contorted to merely flexuous; gill trama parallel, more or less ochraceous tawny when revived in KOH; pileus trama homogeneous, no pellicle differentiated, floccose, interwoven and pale to dark ochraceous tawny in KOH.


Discussion. The strong odor and clustered manner of growth on soil should be distinctive. The spores are almost identical with those of *P. silvatica*, and I suspect the species belongs in that group even though no bluish or greenish color is reported for it.

Psilocybe lateritia (Murrill) comb. nov. (Figs. 63, 64–type.) Spores ovoid to subellipsoid, terete or nearly so, 7.5–9.3  $\times$  5–6.2  $\mu$ , smooth, not angular, dull yellowish brown in KOH, apical pore present but very obscure; basidia 18–20  $\times$  6–7  $\mu$ , cylindric, hyaline or in thick sections yellowish, four-spored; pleurocystidia scattered, similar to cheilocystidia; cheilocystidia abundant, ventricose with acute apices but elongating to aciculate, 28–46  $\times$  6–9  $\mu$  (narrowest in the longest ones), neck or apex usually with an adhering mass of a highly viscous substance slowly soluble in KOH, content homogeneous and hyaline; gill trama of parallel hyphae, sordid pale yellowish in KOH; pileus trama with a thin cuticle of hyaline to pale yellowish gelatinous hyphae bearing clamp connections, beneath this a narrow region of cinnamon-brown hyphae (in KOH) but not otherwise distinct from the remainder of the flesh which is paler brownish to yellowish.

Discussion. This species was described as Psathyra lateritia, by Murrill (Mycologia 10: 33. 1918). It belongs in the group with P. atrorufa, but since this group is in need of a critical study, no attempt at further comparisons will be made here. It is not identical with any of my collections in that group.

PSILOCYBE LATISPORA Murrill, Mycologia 15: 10. 1923. (Figs. 65, 66–type.) Spores 6.2– $7.8 \times 4.6$ – $5.2 \times 6$ – $6.2 \mu$ , flattened, subovate to subcircular in outline in face view, often slightly angular, in side view slightly inequilateral to subelliptic, dark dull yellowish brown in KOH, apex truncate from an apical hyaline pore; basidia 18– $20 \times 7$ – $8 \mu$ , hyaline in KOH, four-spored; pleurocystidia none: cheilocystidia 22– $26 \times 4$ – $7 \mu$ , aciculate to narrowly ventricose, apex at times with a hyaline globule; gill trama regular, hyphae parallel to somewhat interwoven, with dull yellowish brown

walls in KOH; pileus trama homogeneous, surface hyphae slightly gelatinous in KOH but no true pellicle present, floccose, compactly interwoven and with dull yellowish brown walls when revived in KOH.



Figs. 66-83. Microscopic characters of dark-spored agarics.

when aline, dexun reifferawny

owth ntical gs in orted

ype.) 5.2 μ, pore hyastidia entri-5–9 μ h an le in rallel thin aring monthe

ritia, roup itical . It

923. flatghtly dull aline turoowly ular, own Discussion. This species is apparently closely related to P. phyllogena Peck but differs in its larger spores.

**Psilocybe mammillata** (Murr.) comb. nov. (Figs. 67, 68–type.) Spores 5–6 (7)  $\times$  4.5–5  $\mu$ , smooth, triangular in face view, some ovate-angular, in side view subelliptic to ovate, dull ochraceous tawny revived in KOH, with a small apical pore; basidia hyaline in KOH,  $13-15\times5-6~\mu$ , four-spored; pleurocystidia similar to cheilocystidia; cheilocystidia small and inconspicuous,  $12-18\times4-8~\mu$ , fusoid-ventricose or the apices subcapitate, hyaline; gill trama parallel, pale yellowish bister from a pigment which encrusts the hyphae, darker near subhymenium; pileus trama homogeneous, the hyphae near the surface a darker yellowish brown (bister) than those toward the subhymenium, clamp connections present.

Discussion. Murrill described the fungus as Psathyra mammillata (Mycologia 10: 33. 1918). The spores and cheilocystidia are distinctive. Psathyra cinchonensis Murrill (Mycologia 10: 33. 1918) is identical microscopically with P. mammillata and in the dried specimens the gill spacing did not appear distinctive. Psathyra cinchonensis is therefore considered a synonym of P. mammillata.

Psilocybe modesta (Peck) comb. nov. (Figs. 69, 72-type.) Spores 5.5-6.3  $\times$  3.6-4.2  $\times$  4-4.3  $\mu$ , subtriangular to broadly ovate in face view, elliptic in side view, pale bister to sordid tawny revived in KOH, apex truncate because of a small hyaline germ pore; basidia hyaline in KOH, four-spored, 14- $18 \times 4.5$ - $5.5 \mu$ ; pleurocystidia only near gill edge and similar to cheilocystidia; cheilocystidia abundant, hyaline, 22- $26 \times 3$ - $5 \mu$ , narrowly ventricose, often with drops of a mucilaginous substance adhering to apex; gill trama regular to somewhat interwoven and colored like the pileus trama or paler; pileus trama with a thin poorly formed pellicle of subgelatinous hyphae (in KOH), flesh proper floccose and dark, bright cinnamon in KOH (pale in very thin sections), pigment loosely encrusted on cell walls.

Discussion. In its small spores it resembles P. mammillata but the veil is pronounced and the cheilocystidia much longer.

PSILOCYBE NIGRELLA Peck, Bull. N. Y. State Mus. 139: 28. 1910. (Figs. 70, 71–type.) Spores 9–11 (12)  $\times$  6–7.5 (8)  $\mu$ , dark sordid yellowish brown revived in KOH, smooth, terete, broadly ovoid to ellipsoid, apical pore small and obscure; basidia

7, 68n face e, dull pore; oleuroinconbcapirom a enium;

darker

enium.

to P.

ystidia in 10: and in nctive. of P.

type.) ovate tawny germ -5.5 μ; stidia; wentring to ed like formed occose tions),

millata er. 9: 28. (8) μ, terete,

basidia

four-spored,  $22-26\times7-8.5~\mu$ , hyaline in KOH; pleurocystidia none seen (some collapsed basidia brown); cheilocystidia abundant and forming a sterile band on the gill edge, hyaline, narrowly ventricose to almost aciculate,  $20-26\times4-7~\mu$ , apices often somewhat flexuous; gill trama of long broad parallel cells with parallel walls, pale rusty brown in KOH, subhymenium cellular and dark rusty brown, the pigment encrusted on the cell walls; pileus trama with a gelatinous pellicle of narrow  $(2-4~\mu)$  hyaline appressed hyphae, beneath this a zone of dark yellowish brown hyphae with the pigment encrusted on the walls, paler toward the subhymenium, the hyphae interwoven and all with broad cells  $(8-15~\mu$  in diam.).

Discussion. The short stipe is the only character I can find to distinguish it from P. atrobrunnea, and length of stipe is a very variable character in bog-inhabiting species.

**Psilocybe pallidispora** (Murrill) comb. nov. (Figs. 73–75–type.) Spores 5.3–6.2 × 3.5–4.2  $\mu$ , terete, ellipsoid to ovoid, with a very small apical pore, dull yellowish brown in KOH; basidia 16–20 × 5–6  $\mu$ , four-spored, hyaline in KOH, pleurocystidia scattered, 34–44 × 7–10  $\mu$ , fusoid-ventricose with elongated necks and acute apices, the apex or neck frequently with a drop or coating of a highly viscous substance adhering to it, hyaline in KOH, thin-walled, content homogeneous; cheilocystidia 22–28 × 5–7  $\mu$ , fusoid-ventricose, in some subcapitate, hyaline in KOH, often with mucilaginous cap; gill trama regular, dark brown (near cinnamon-brown) when revived in KOH; pileus trama homogeneous, no pellicle seen, surface region a darker brown than that toward the subhymenium, pigment encrusted on the walls.

Discussion. Distinct by virtue of the small ellipsoid spores and moderately large pleurocystidia. The pileus is typically dry.

PSILOCYBE PANAEOLIFORMIS Murrill, Mycologia 15: 12. 1923. (Figs. 76–78.) Spores 9–11.5 × 6–7  $\mu$ , smooth, ellipsoid to subovoid, a few obscurely angular, not compressed, sordid yellowish brown to pale bister revived in KOH, apex truncate from a hyaline apical pore; basidia four-spored, short and fat, 14–16 × 8–9  $\mu$ , hyaline in KOH; pleurocystidia none seen; cheilocystidia very abundant and forming a sterile band on gill edge, 12–16 × 4–8  $\mu$ , ventricose and with a slender flexuous neck, apices subacute, hyaline in KOH; gill trama parallel to subparallel, pallid, brownish in KOH; pileus trama with a pellicle of narrow (1.5–3  $\mu$ ) filamentous hyaline hyphae subgelatinous in KOH, a few pilocystidia similar to cheilocystidia projecting from it, beneath this a hypo-

derm of enlarged cells cinnamon-brown in KOH, remainder of flesh also of enlarged hyphal cells but paler, clamp connections present on pellicle.

Discussion. This species appears to be a true Psilocybe and closely related to P. coprophila. It differs from the latter in terete somewhat narrower spores and short fat cheilocystidia. An imperfect was fruiting over the type and details of cap and gills could not be seen as clearly as desired. However, the data obtained were checked with the Alabama collection which is unquestionably the same.

PSILOCYBE PHYLLOGENA Peck, Bull. N. Y. State Mus. 157: 99. 1912. (Figs. 79, 80–type.) Spores 5–6.5 (7) × 4–4.6 × 5–6  $\mu$ , more or less triangular in face view, often top-shaped or merely ventricose near base, subelliptic in side view, apical hyaline germ pore present causing apex to appear slightly truncate, when revived in KOH sordid tawny brown in color; basidia four-spored, hyaline in KOH, 18–22 × 5–7  $\mu$ ; pleurocystidia none or present only near the gill edge and similar to cheilocystidia; cheilocystidia narrowly fusoid, apices often with a mucilaginous cap in KOH, content hyaline and homogeneous, 22–26 (34) × 4–6 (7)  $\mu$ ; gill trama regular, parallel or nearly so, pallid brownish in KOH to nearly hyaline; pileus trama homogeneous, no true pellicle present but surface hyphae only slightly gelatinous, flesh proper pallid sordid brownish in KOH, finally becoming nearly hyaline, interwoven and floccose.

Discussion. The lack of a veil in addition to the small compressed spores appears to be distinctive, though the absence of a veil needs to be confirmed from an examination of numerous collections of young carpophores.

PSILOCYBE PLUTONIA (Berk. & Curt.) Saccardo, Syll. Fung. 5: 1056. 1887. (Fig. 81–type.) Spores 5–6.2  $\times$  3.8–4.2  $\times$  5–6  $\mu$ , subtriangular in face view to angular-ovate or ventricose basally, often slightly indented around the apiculus, elliptic in side view, distinctly compressed, dull yellowish brown in KOH, with a small apical hyaline pore; basidia pale sordid yellowish to hyaline in KOH, 10–12  $\times$  5–6  $\mu$ , four-spored; pleurocystidia and cheilocystidia not seen (no good sections of gill edges obtained); gill trama parallel, dark sordid yellowish brown in KOH; pileus trama homogeneous, dark yellowish brown (near bister) in KOH, no pellicle observed.

Discussion. This species appears to be in the P. phyllogena series and is distinct on spore characters alone.

AGARICUS POLYTRICHOPHILUS Peck, Ann. Rep. N. Y. State Mus. 30: 42. 1878. (Fig. 83–type.) Spores 6–7.5 × 4–5  $\mu$ , terete, ellipsoid to ovoid, smooth, not angular but an occasional spore ventricose at base, dull yellowish brown in KOH, apical pore present but small; basidia four-spored, 16–20 × 6–7.5  $\mu$ , hyaline in KOH; pleurocystidia very rare and similar to cheilocystidia; cheilocystidia 16–23 × 5–7  $\mu$ , narrowly fusoid-ventricose with obtuse apices, hyaline and with homogeneous contents when revived in KOH; gill trama parallel, yellowish brown; pileus trama with a thin gelatinous pellicle of hyaline hyphae, flesh proper compactly interwoven and yellowish brown in KOH.

Discussion. I am not sure at present as to the status of this species. Our common Psilocybe on Polytrichum has no odor and so I am not inclined to consider Peck's type identical with it.

Psilocybe pyrispora (Murrill) comb. nov. (Figs. 82, 84type.) Spores  $6.5-7.5 \times 4-4.5 \mu$ , smooth but some obscurely angular, terete, somewhat pear-shaped to ellipsoid or ovoid (angularity best seen in face view, dull yellowish brown when revived in KOH (pale smoky purplish brown in mounts of fresh material -Murrill), apical pore present but small; basidia four-spored, hyaline in KOH,  $12-14 \times 6-7 \mu$ ; pleurocystidia scattered to fairly abundant, 18-26 (32)  $\times$  10-15  $\mu$ , sessile or with a short pedicel, broadly ovate to elliptic in optical section, hyaline and highly refractive in KOH but content homogeneous, the walls only slightly thickened, smooth; cheilocystidia  $18-26 \times 4-7 \mu$ , narrowly ventricose to subcylindric, the necks straight or flexuous and apices often capped with mucilage, hyaline in KOH; gill trama not reviving well but apparently subgelatinous in KOH; pileus trama with a broad band of indistinct hyaline hyphae forming the upper region, lower region yellowish brown (upper portion apparently subgelatinous in KOH but details were not distinct).

Discussion. A very curious species by virtue of the pleurocystidia. It is the wall which refracts the light in the characteristic manner, and since the content of the cystidium is homogeneous and hyaline, the species cannot be properly regarded as belonging in Naematoloma. Murrill described it as Deconica pyrispora (Mycologia 14: 261. 1922).

der of ections

be and


tter in ystidia. ap and ne data is un-

merely e germ en repored, present ystidia KOH, µ; gill OH to present pallid inter-

come of a

ng. **5**: 5–6 μ, asally, view, small ne in locystrama nomo-ellicle

PSILOCYBE SUBVISCIDA (Peck) Kauffman, Agar. Mich. p. 275. 1918. (Deconica subviscida Peck, Ann. Rep. N. Y. State Mus. 41:70. 1888). (Figs. 85–87–type.) Spores 7–8 × 4–4.5 (5)  $\mu$ , smooth, ellipsoid to somewhat ovoid, not appreciably angular, terete, yellowish brown in KOH, with a small apical germ pore; basidia 18–20 × 6–7  $\mu$ , four-spored, hyaline in KOH; pleurocystidia clustered between the gills, rare elsewhere, 34–50 × 7–9



Figs. 84-93. Microscopic characters of dark-spored agarics.

(10). $\mu$ , ventricose, elongating to subcylindric with flexuous necks and obtuse apices, hyaline, thin-walled; cheilocystidia variable, ventricose to ellipsoid at first but soon developing a flexuous projection up to 30–40  $\mu$  long and 6–7  $\mu$  thick or fusoid-ventricose from the first, the former 20–26  $\times$  10–12  $\mu$  at first, 36–50  $\times$  7–10  $\mu$  in age, the others usually smaller and 20–32  $\times$  5–8  $\mu$ ; gill trama regular or nearly so, color not distinctive; pileus trama with a

thin gelatinous pellicle of narrow hyaline hyphae, remainder compactly interwoven and floccose, not distinctively colored in KOH.

Discussion. The occurrence of pleurocystidia between the gills should not be given much emphasis. In many species such structures develop along the line where a new gill is forming and hence they could also be interpreted as cheilocystidia. I suspect that such is the situation here.

Psilocybe tomentosa (Murrill) comb. nov. (Figs. 88-91type.) Spores  $8-10 \times 5-6 \mu$ , smooth, slightly compressed to terete, angular to elliptic in face view, slightly bean-shaped to subelliptic in side view, some ventricose basally and with a median constriction (apparently abnormal), dull tawny revived in KOH, apical pore hyaline and very inconspicuous; basidia mostly four-spored, hyaline in KOH; 24-28 × 5.5-7 µ; pleurocystidia rare and embedded in the hymenium, similar to cheilocystidia or shorter; cheilocystidia  $34-48 \times 5-8 \mu$ , subcylindric to subfusoid, the apices obtuse, broadest at or above the middle and neck very short and indistinct, a few with the apex embedded in a mucilaginous substance; gill trama parallel to somewhat interwoven, pale yellowish in KOH; cuticle of pileus consisting of a turf of upright pilocystidia and chains of ± globose cells with a pale tawny encrusting pigment on the walls, the pilocystidia more or less clavate, 20-52  $\times$  8-12  $\mu$  and irregularly arranged so as to project to varying distances above each other, hyaline or with encrusted pigment, the chains of vesiculose cells not highly colored in KOH (as they are in genus Cystoderma), masses of vesiculose cells and pilocystidia aggregated into fascicles to form scales on the cap; flesh of cap of interwoven hyphae which are pale yellowish in KOH, clamp connections present.

Discussion. Although the microscopic characters of this species depart radically from those of most psilocybes in respect to the cuticle of the pileus, the differences do not seem to me sufficient to justify erecting a new genus.

PSILOCYBE VIALIS Murrill, Mycologia 15: 11. 1923. (Figs. 92, 93-type.) Spores pale yellowish in KOH, 7-8.4  $\times$  4-4.6  $\mu$ , terete, ellipsoid to somewhat ovoid, some obscurely angular in face view, apex truncate from a small hyaline germ pore; basidia 17-20  $\times$  5-6.5  $\mu$ , four-spored, hyaline in KOH; pleurocystidia embedded in hymenium or subhymenium, 20-26  $\times$  8-10  $\mu$ , clavate to obovoid above a thick pedicel, brownish in KOH and with a granular content or some with a highly refractive amorphous sub-

p. 275. te Mus.  $5(5) \mu$ , ungular, n pore; urocys- $0 \times 7-9$ 

/ C/C

necks , venrojecfrom O μ in trama stance; cheilocystidia abundant,  $20\text{--}26 \times 4\text{--}6\,\mu$ , aciculate to narrowly fusoid, hyaline, apex often with an adhering globule of a hyaline viscous substance; gill trama regular, the hyphae somewhat interwoven, pale ochraceous tawny in KOH; pileus trama lacking a distinct pellicle, surface region nearly hyaline, more or less ochraceous tawny toward the gills.

Discussion. The pleurocystidia are not those of Naematoloma, and I doubt if the species is any more closely related to members of that genus than are the other psilocybes.

### **EXPLANATION OF FIGURES**

The illustrations of spores, cystidia, and veil elements were drawn with the aid of a camera lucida. As reproduced the spores are approximately  $1650 \times$  natural size; the cystidia and veil elements are approximately  $700 \times$  natural size unless otherwise stated.

FIGS. 1-9. Coprinus angulatus: 1, pilocystidia; 2, spores in side and face view. Coprinus Brassicae: 3, spores in side and face view. Coprinus calyptratus: 4, spores in both views. Coprinus asterophorus: 6, spores in both views. Coprinus cinchonensis: 5, cells from veil showing spines; 7, spores in both views. Coprinus ebulbosus: 8, spores—some showing ventral hump (or pore). Coprinus hexagonosporus: 9, spores in both views.

Figs. 10-22. The spores are shown in both side and face views. Coprinus insignis: 10, spores. Coprinus jalopensis: 11, spores. Coprinus Jonesii: 12, spores. Coprinus pseudoradiatus: 13, spores. Coprinus laniger: 14, spores. Coprinus mexicanus: 15, spores. Coprinus pulchrifolius: 16, spores. Coprinus quadrifidus: 17, spores. Coprinus rotundisporus: 18, pleurocystidia; 19, spores. Coprinus Seymourii: 20, spores. Coprinus semilanatus:

21, spores. Coprinus silvaticus: 22, spores.

Figs. 23-31. The spores are shown in both side and face views. Coprinus Spraguei: 23, six spores. Coprinus variegatus: 24, ten spores, two of which show ventral hump or pore. Coprinus subpurpureus: 25, spores. Panaeolus anomalus: 26, spores; 27, pleurocystidia. Panaeolus castaneifolius: 28, spores. Panaeolus fraxinophilus: 29, spores; 30, pilocystidia; 31, cheilocystidia.

Figs. 32-38. The spores are shown in both views. Panaeolus reticulatus: 32, spores. Panaeolus rufus: 33, cheilocystidia; 34, spores. Panaeolus solidipes: 35, spores; 36, pleurocystidia. Panaeolus variabilis: 37, cheilo-

cystidia; 38, spores.

Figs. 39-48. Panaeolus venenosus: 39, spores; 40, cheilocystidia. Panaeolus fontinalis: 41, spores; 42, cheilocystidia. Naematoloma campestre: 43, spores; 44, pleurocystidia; 45, cheilocystidia. Naematoloma humidicola: 46,

spores; 47, pleurocystidia; 48, cheilocystidia.

Figs. 49-65. Naematoloma petasiforme: 49, pleurocystidia; 50, spores; 51, cheilocystidia. Psilocybe bulbosa: 52, spores; 53, cheilocystidia. Psilocybe caespitosa: 54, spores; 55, cheilocystidia. Psilocybe castanella: 56, cheilocystidia; 57, spores. Psilocybe californica: 58, cheilocystidia: 59, spores. Psilocybe fuliginosa: 60, spores; 61, cheilocystidia. Psilocybe

to narile of a

somestrama

toloma, embers

with the 1650 × natural

and face s calypin both spores al hump

oprinus lonesii: er: 14, spores. urocysanatus:

which maeolus spores.

naeolus cheilo-Panae-

re: 43, la: 46,

pores;
Psiloa: 56,
a: 59,
locybe

graveolens: 62, spores. Psilocybe lateritia: 63, pleuro- and cheilocystidia; 64, spores. Psilocybe latispora: 65, spores.

FIGS. 66-83. Psilocybe latispora: 66, cheilocystidia. Psilocybe mammillata: 67, spores; 68, cheilocystidia. Psilocybe modesta: 69, spores; 72, cheilocystidia. Psilocybe nigrella: 70, spores; 71, cheilocystidia. Psilocybe pallidispora: 73, pleurocystidia; 74, spores; 75, cheilocystidia. Psilocybe panaeoliformis: 76, spores; 77, cheilocystidia; 78, pilocystidia. Psilocybe phyllogena: 79, cheilocystidia; 80, seven spores. Psilocybe plutonia: 81, spores. Psilocybe pyrispora: 82, pleurocystidia. Agaricus polytrichophilus: 83, spores.

Figs. 84–93. Psilocybe pyrispora: 84, spores. Psilocybe subviscida: 85, cheilocystidia; 86, cheilocystidia; 87, spores. Psilocybe tomentosa: 88, pilocystidia; 89, veil elements; 90, cheilocystidia; 91, spores. Psilocybe vialis: 92, pleurocystidia; 93, spores.

# AN UNDESCRIBED SPECIES OF HELMIN-THOSPORIUM ON SUDAN GRASS AND SORGHUM

C. L. LEFEBURE AND HELEN S. SHERWIN

(WITH 2 FIGURES)

In 1939, G. W. Burton collected and sent us (7) leaves of a common Sudan grass × Leoti sorghum hybrid (now Tift Sudan grass), bearing a leaf spot that was defoliating some of his breeding material. Lesions on these leaves varied considerably, the smaller ones being 0.5 to 1.0 mm, in each direction, whereas the longer ones measured up to 15 mm, in length and 1 to 6 mm, in The lesions were round to elliptical, the long axis parallel to the leaf veins, and oftentimes the spots seemed to be limited in their spread by the veins. Many of the lesions had coalesced so that comparatively large portions of the leaves were affected (FIG. 1. A). The most characteristic features of the lesions, however. were the alternate light tan and darker, more narrow bands of tissue, producing a zonate appearance which suggests the common name "target spot" (FIG. 1, A). Lesions were usually surrounded by a narrow tannish-brown border setting off the affected area from the healthy tissue. The symptoms on Tift Sudan grass are strikingly similar to the lesions on corn and teosinte caused by the fungus originally described as Ophiobolus heterostrophus Drech. (5), but later transferred to the genus Cochliobolus (6).

Although the fungus causing the target spot of Tift Sudan grass has never been collected on common Sudan grass, the latter was found to be susceptible when inoculated. On common Sudan, however, the lesions appear less zonate (FIG. 1, B). The spots are often uniformly dark, or when older they may have a straw-color center surrounded by a reddish-purple border; others have an additional band of straw-color tissue that in turn is surrounded by a dark border. The striking difference in the leaf spots of the two varieties of Sudan grass is in the color because Tift Sudan

IN-

of a udan reed-, the s the m. in rallel ed in ed so (FIG. ever, ds of nmon inded area s are d by phus (6). grass was udan, spots trawhave

nded ts of udan



Fig. 1. Target-spot on Sudan grass.

grass has the tan pigment, hence, the tan leaf spot, whereas common Sudan grass contains the dark pigment, hence, the purplishblack lesions.

Under field conditions the fungus sporulates rather sparsely on both the upper and lower surfaces of infected leaves. Furthermore, sporulation is restricted to the dead, brown tissue of the lesions, and this paucity of fruiting may account for the slow spread of the disease. When infected leaves are placed in a moist chamber, however, sporulation is profuse, at first restricted to the lesions but later, as the leaf tissue becomes flaccid, the leaf surfaces may be completely covered with spores.

The conidia of the fungus are usually moderately curved, widest at or near the middle and taper slightly toward the rounded ends, regardless of the host from which they were collected (Fig. 2, A, B, C, D). Although the conidia of this Sorghum fungus are golden yellow to light olivaceous, resembling the conidia of Cochliobolus heterostrophus (Helminthosporium maydis Nisikado and Miyake) in color, they are less strongly curved, smaller, and have fewer septa than the corn fungus. The average number of septa in the conidia of the Sorghum fungus is 5.1, whereas in the corn fungus it is 7.7. The greater average number of septa in the conidia of the corn fungus is one of the more striking morphological characteristics that can be used to distinguish this fungus from the one here described on Sorghum.

The conidia are borne on conidiophores that usually emerge singly from a stoma, or directly through the epidermis (Fig. 2, D, a), but under favorable conditions two to four may emerge from a single stoma (Fig. 2, D, b). Comparatively few conidia seem to be borne on each conidiophore, there being often only one conidium under field conditions, whereas in a moist chamber two to four are often produced, as indicated by the number of geniculations (Fig. 2, D, a, b). In a moist chamber, or on agar where there is abundant moisture, a conidium borne on the primary conidiophore, and while still attached to it, may germinate and produce a secondary conidiophore bearing one or two conidia (Fig. 2, D, c). This may be repeated, the secondary spore germinating to produce a tertiary conidiophore, which in turn bears one or two conidia, giving the appearance of the conidia being

superimposed and produced in a chain-like manner. These secondary conidiophores are usually much shorter than the primary ones, but the secondary conidia may be as large as the primary ones. This type of conidial proliferation is very characteristic of this fungus and aids in distinguishing it from several other species of *Helminthosporium* that occur on grasses (4). In water or on agar, the conidium germinates usually by producing two polar germ tubes (Fig. 2, D, d), and oftentimes a germ tube-like hypha emerges from an attached conidiophore (Fig. 2, D, e).

The fungus grows well on corn-meal and potato-dextrose agars forming a compact mass of aerial hyphae and conidia. As the colony becomes older, sporulation becomes usually more abundant near the margin. On both corn-meal and potato-dextrose agar, the growth is whitish at first, turning light gray or light gravishbrown when the fungus begins to sporulate and later grayish-olive to slate olive (8). Small whitish upright tufts of hyphae are sometimes formed within or on the border of the mat. The growth on potato-dextrose agar is more profuse than that on corn-meal agar. When the fungus grows from diseased leaf fragments plated on potato-dextrose agar, the surrounding agar becomes a vinaceous-cinnamon or fawn color. On corn-meal agar the color is much fainter, being light vinaceous-cinnamon when distinguishable. In a solution of carrot decoction plus one per cent dextrose, the fungus grows well, forming a fluffy mat of mycelium on the surface of the liquid. The color of the mycelial mat is white at first and later grayish-olive to slate olive.

### TAXONOMY OF THE FUNGUS

Helminthosporium sorghi was described by Schweinitz (10, p. 279) in 1832 from decaying leaves of Sorghum collected near Lititz, Pennsylvania. Later, Cooke (3, p. 141) used the same name for a fungus that he found on Sorghum that had been collected by Ravenel and sent to him for identification, which Ravenel issued later as No. 167 of his Fungi Americani Exsiccati. Saccardo (9, p. 420), seeing that the specific name had been preoccupied, set up the new name Helminthosporium cookei Sacc. Hence, H. sorghi Cooke became a synonym of H. cookei Sacc.

complish-

of the slow in a ricted e leaf

videst ends, IG. 2, is are Cocho and have septa corn in the ologifrom

merge ag. 2, merge onidia y one r two enicuwhere mary

midia gerbears being

and

The writers examined the type specimen of Helminthosporium sorghi Schw. from the Schweinitz Collections in the herbarium of the Academy of Natural Sciences of Philadelphia, and were unable to find conidia of a Helminthosporium. We also examined what undoubtedly is co-type material, in the Mycological Collections of the Bureau of Plant Industry and in the Farlow Herbarium, Harvard University, and again it was impossible to find conidia. There were, however, black setae of a Colletotrichum sp. in typical anthracnose lesions present in abundance on these leaf speci-Whether Schweinitz observed these setae and mistook them for conidia of Helminthosporium cannot be determined now, although he described the Helminthosporium "pustules" as being black with conidia that were concolorous. In addition, the lesions on these leaf specimens do not resemble those caused by our fungus. so it seems obvious that the organism causing the leaf spot on Sorghum here described is clearly distinct.

We have also examined Ravenel's Fungi Americani Exsiccati specimen No. 167 and found a dark, crust-like, effuse fungus growth on the surface of a *Sorghum* culm. The fungus was superficial, produced no lesions, indicating the organism was probably a saprophyte. A few dark, thick-walled, straight conidia were found that looked wholly different from those produced by our fungus. These conidia were not associated with conidiophores that resemble those of the genus *Helminthosporium*.

Ciferri and Gonzalez Fragoso (2) described Helminthosporium sudanensis Frag. and Cif. on the flower parts of Sudan grass [Sorghum vulgare var. sudanense (Piper) Hitchc.]. Examination of specimens of the above material from the Farlow Herbarium and the Mycological Collections of the Bureau of Plant Industry reveals conidia that are mostly sharply curved, with the central cell larger and darker than the others, and the end cells practically hyaline. According to Boedijn (1), such fungi should be placed in the genus Curvularia. Since the fungus on Sudan grass described by Ciferri and Gonzalez Fragoso has, in addition to the above characteristics, conidia that are 4-septate and the conidial measurements agree with those of C. geniculata (Tracy and Earle) Boedijn, it is likely that species.

mm of nable what as of ium, idia. typi-pecitook now, being sions agus, t on

was was couced idio-

grass nina-Her-Plant n the cells

nould udan lition I the Cracy A

Fig. 2. Conidia and conidiophores of Helminthosporium sorghicola.

It appears, therefore, that the fungus that produces the target spot on Sorghum spp. described in this paper is widely different from the species of Helminthosporium previously reported on these hosts, and we believe it is an undescribed Helminthosporium. Since it occurs on several species of Sorghum, and because sorghi and sudanensis have already been used as specific names in the genus Helminthosporium, we propose to use the specific epithet sorghicola for the new species described hereafter.

### Helminthosporium sorghicola sp. nov.

Conidiophoris singulis vel 2-4-caespitosis, typice simplicibus interdum ramosis, olivaceo-brunneis, 5.5–10.5  $\mu$  in diam., 115–700  $\mu$  longis vel aliquando in madore longioribus; conidiis secundariis vulgaribus, brevioribus, 5–7  $\mu$  in diam., 25–200  $\mu$  longis; conidiis aureo-flavis usque pallide olivaceis, 20–105  $\mu$   $\times$  8.5–20.6  $\mu$ , plerumque curvatis, circa medium latissimis et apices rotundatos versus attenuatis, 2–8-septatis, tenui-tunicatis, hilo inconspicuo praeditis.

Hab. in foliis glumisque Sorghi spp., U. S.

tissue is placed in moist chamber.

Producing well-defined spots on *Sorghum* spp., mostly in the Southern United States. Spots small, tan or reddish-purple, depending on host pigment—tan on Tift Sudan grass, reddish-purple on common Sudan grass; older lesions usually show target or zonate pattern, with light centers, surrounded by narrow band of darker tissue, and this bordered by a wider band of light tissue, etc. Lesions range in size from barely visible to  $1 \times 15$  mm.; these may coalesce to involve the whole leaf. Lesions at first round to elliptical, becoming more elongated as they are somewhat limited by the leaf veins. Fruiting of fungus on leaves and glumes sparse under field conditions, abundant when infected

Conidiophores arising singly or in groups of 2 to 4, usually from stomata or occasionally singly between ordinary epidermal cells; typically simple, occasionally branching; dark, olivaceous brown; measuring 5.5 to  $10.5\,\mu$  in diameter and 115 to  $700\,\mu$  in length, sometimes longer under very moist conditions; secondary conidiophores common, especially under moist conditions, shorter, 5 to  $7\,\mu$  in diameter and 25 to  $200\,\mu$  in length. Conidia on Tift Sudan grass leaves in moist chamber golden yellow to light olivaceous, measuring 20 to  $105\,\mu \times 8.5$  to  $20.6\,\mu$  (means =  $59.2\,\times 14.1\,\mu$ ), usually curved, widest near the middle, and tapering slightly toward rounded ends; 2 to 8 septa (mean = 5.1); peripheral wall thin; hilum moderately broad, not conspicuous; germinat-

ing by the production of two polar germ tubes.

arget

erent

these

ium.

orghi

the

ithet

n ra-

μ in

105 µ

datos

the

de-

rple

t or

d of

sue, m.;

first

me-

and

cted

ally

mal

eous

in lary

ter,

Tift oli-

 $2 \times$ 

ing

ph-

nat-

Found on diseased leaves of Sorghum vulgare var. sudanense (Common) × Sorghum vulgare (Leoti) hybrid, Tifton, Ga., February 15, 1939; on leaves of Sorghum vulgare var. sudanense (Piper) Hitchc. (Tift variety), Gainesville, Fla., August 12, 1942; Tifton, Ga., August 19, 1943 (type), Herb. No. 896; on leaves of Sorghum vulgare Pers., Baton Rouge, La., July 1944, and on leaves of Sorghum halepense (L.) Pers., Cordele, Ga., August 22, 1945.

In addition to the above, Helminthosporium sorghicola has been found on the following specimens in the Mycological Collections of the Bureau of Plant Industry under the name Helminthosporium sorghi Schw.; on Sorghum halepense, College Station, Texas, fall, 1889, H. S. Jennings No. 80; on Sorghum halepense, College Station, Texas, January 10, 1890, H. S. Jennings; on Sorghum halepense, Austin, Texas, August, September, October, November, 1900, W. H. Long No. 447; on Sorghum, Etheridge, N. C., August 14, 1905, F. L. Stevens No. 404; on Sorghum halepense, San Antonio, Texas, July 29, 1910, W. P. C.

The writers wish to express their thanks to Dr. Francis W. Pennell, The Academy of Natural Sciences of Philadelphia, Mr. A. P. D. Piquet, Farlow Herbarium, Harvard University, and Mr. John A. Stevenson, Bureau of Plant Industry Station, for placing pertinent material at our disposal for study; to Miss Edith Cash for preparation of the Latin diagnosis; and to Dr. D. C. Bain, Louisiana State University, for sending a specimen of Sorghum M. N. 608.

Type specimens of *Helminthosporium sorghicola* have been deposited in the Mycological Collections of the Bureau of Plant Industry, Beltsville, Maryland, and in the Farlow Herbarium, Harvard University, Cambridge, Massachusetts.

DIVISION OF FORAGE CROPS AND DISEASES, PLANT INDUSTRY STATION, BELTSVILLE, MD.

### LITERATURE CITED

- Boedijn, K. B. Ueber einige phragmosporen Dematiazeen. Bull. Jard. Bot. Buitenzorg (Ser. 3) 13 (1): 120-134. 1933.
- Ciferri, R., and Gonzalez R. Fragoso. Hongos parasitos y saprofitos de la República Dominicana. Bol. R. Soc. Espan. Hist. Nat. 26: 491-500. 1926.

Cooke, M. C. Ravenel's American Fungi. Grev. 6: 129-146. 1878.
 Drechsler, Charles. Some graminicolous species of Helminthospo-

rium: I. Journ. of Agric. Res. 24: 641-740. 1923.

 Leafspot of maize caused by Ophiobolus heterostrophus, n. sp., the ascigerous stage of a Helminthosporium exhibiting bipolar germination. Journ. of Agric. Res. 31: 701-726. 1925.

 Phytopathological and taxonomic aspects of Ophiobolus, Pyrenophora, Helminthosporium and a new genus, Cochliobolus.

Phytopathology 24: 953-983. 1934.

 Lefebvre, C. L. An undescribed leaf spot of Tift Sudan Grass. Phytopathology 34: 992. 1944.

Ridgway, R. Color standards and color nomenclature. Washington. 1912.

9. Saccardo, P. A. Sylloge Fungorum. 4: 420. 1886.

 Schweinitz, L. D. de. Synopsis Fungorum in America Boreali media degentium. Trans. Am. Phil. Soc. of Phil. N. S. 4: 141– 318. 1832.

### **EXPLANATION OF FIGURES**

Fig. 1. Helminthosporium sorghicola on leaves of A. common Sudan  $\times$ 

grass Leoti (Sorghum) hybrid; B. common Sudan grass, × 2.

Fig. 2. Helminthosporium sorghicola, drawn with the aid of a camera lucida, A, B, C and D  $\times$  500, except D, a, b  $\times$  250. A, Conidia from Johnson grass, Sorghum halepense; B, conidia from Sorghum vulgare, variety Rex; C, conidia from Sorghum vulgare, variety M. N. 608; and D, conidia and conidiophores from Sorghum vulgare var. sudanense (Tift); a, single conidiophores emerging directly through the leaf epidermis; b, group of conidiophores emerging through a stoma; c, primary conidia that have germinated and produced secondary conidiophores and conidia; d, germinating conidia, producing polar germ tubes; and e, primary conidium having produced a secondary conidiophore, a germ tube-like hypha now emerging from the distal end of this conidiophore.

# TWO UNUSUAL FUNGI FROM GLACIER NATIONAL PARK, MONTANA

E. B. MAINS

(WITH 1 FIGURE)

In 1941, the writer spent several weeks in Glacier National Park, Montana. Among the fungi collected, two, *Mesopsora Hypericorum* and *Mitrula gracilis*, were of more than ordinary interest. Both were obtained on the continental divide along the trail from Logan Pass to Hidden Lake at altitudes around 7,000 feet.

### MESOPSORA HYPERICORUM (Winter) Dietel

As far as the writer is aware this is the first report of M. Hypericorum for North America. The following description has been prepared from the collection.

Uredinia hypophyllous and caulicolous, golden to orange yellow when fresh, whitish when dried, pulverulent, flat to somewhat pulvinate, circular to elliptic in outline, 0.2–0.8 mm. wide, ruptured epidermis fairly conspicuous; urediniospores catenulate in short chains, variable in shape, broadly ellipsoid, obovoid to globoid, 18–24  $\mu$  long, 14–18  $\mu$  wide, wall colorless, 1.5–2.0  $\mu$  thick, moderately verrucose with irregular warts; telia not seen.

On Hypericum Scouleri Hook., Logan Pass, Trail near Hidden Lake, Glacier National Park, Montana, September 20, 1941, E. B. Mains (6121).

The rust was first described by DeCandolle as *Uredo Hypericorum*. Winter (16) apparently first described the telia and placed the species in *Melampsora* because the telia are typical of that genus. For a number of years the rust was treated as a *Melampsora* having only uredinia and telia. According to the Sydows (14), Gobi and Tranzschel were the first to note that the uredinia differed in several important aspects from the uredinia

1878. ospo-

us, n. g bi-5.

oolus, oolus.

irass.

hing-

oreali 141-

lan ×

mera nnson Rex; and onidionidi-

nated nidia, ed a the

<sup>&</sup>lt;sup>1</sup> Paper from the Herbarium and the Department of Botany, University of Michigan.

of species of Melampsora. Instead of having pedicellate spores and paraphyses as in other species of Melampsora the sori lack paraphyses and the spores are catenulate in short chains. Fischer (5) also found the same. They concluded therefore that the sori are caeomoid aecia and that the species has aecia and telia and lacks uredinia, and therefore is a melampsoropsis. In support of this interpretation Fischer states that intercalary cells occur in the spore-chain and the sorus is bordered by a slight layer of thin-walled cells suggesting a peridium. In the Montana collection it has not been possible to determine whether intercalary cells occur. Sections through the sori show a slight layer of large thin-walled cells at the margin sometimes extending a short distance upward under the ruptured epidermis. It is questionable that a peridium is formed.

Klebahn (9) found uredinia having paraphyses on Hypericum humifusum and concluded that they belonged to M. Hypericorum and that therefore the species is an autoeumelampsora. However, the uredinia on H. humifusum were not associated with any other spore-form and although on the other collections the sori designated as "caeoma" were found with telia the "uredo" sori did not occur. He reports an experiment in which somewhat scanty telial material giving weak germination was sown on several species of Hypericum, Abies pectinata, Picea excelsa and Larix decidua without results. He expressed considerable doubt concerning his conclusions and in 1914 he (10) described the rust on H. humifusum as a new species, Uredo (Melampsora?) hyperici-humifusi, and treated Melampsora Hypericorum as a melampsoropsis. The Sydows (14, 15) in their monograph of the rusts have followed this latter treatment.

Dietel (2) has decided that the sori with catenulate spores are uredinia. He points out that they have never been reported associated with pycnia and that they are produced in successive generations during the summer. Also they are similar to the uredinia of species of *Coleosporium* and *Chrysomyxa* which produce aecia on conifers. He concludes that the rust is probably heteroecious with aecia on conifers and with uredinia and telia on species of *Hypericum* and that it belongs in a genus intermediate between

Melampsora and Coleosporium for which he proposes the name Mesopsora.

The correct interpretation of the sori of this rust and its relationship to other species of the Melampsoraceae cannot be determined with certainty until its life history is worked out through cultures. Until such is done it seems best to follow Dietel's treatment. The absence of pycnia strongly indicates that the sori are uredinial in function. If the rust is an autoecious melampsoropsis as suggested by the Sydows and Fischer it would be necessary to assume that only secondary or repeating aecia (aecidioid uredinia according to Arthur's terminology) have been collected since pycnia have not been reported. Such a species would be unique in Melampsora: Since the rust appears to be fairly common in Europe, primary aecia accompanied by pycnia would be expected in some collections. If the rust is heteroecious as suggested by Dietel pycnia would not be expected on Hypericum. Also the negative results obtained by Klebahn in cultures of germinating teliospores on species of Hypericum would be expected. Since intercalary cells are known to occur in the uredinia of Coleosporium (Moss, 11) their presence as noted by Fischer in the uredinia of Mesopsora Hypericorum might be expected.

The Sydows have reported M. Hypericorum from Europe, Asia Minor and Africa. It also has been reported from Australia and Japan but such reports may be based on misdeterminations of Melampsora Kusanoi Diet. As far as I am aware Mesopsora Hypericorum has not been previously reported for North America. Therefore the isolated occurrence of the species on an indigenous species of Hypericum is an interesting instance of disjunct distribution.

# MITRULA GRACILIS Karsten (Fig. 1, A-E)

In his monograph of the Geoglossaceae of North America Durand (3) includes two species of Mitrula on mosses, M. gracilis Karsten and M. muscicola E. Henn. Only a few localities are given for each. M. gracilis is listed for Labrador and Newfoundland "attached to and evidently parasitic on Paludella squarrosa." It is stated that Rostrup also has reported the species from Greenland. Apparently the only report of the species for the United

lack scher

and rt of r in thinon it

ccur. alled vard lium

rum ever, ther esignot

s of dua his mi-

are asennia

ecia ous of een

States is that of Seaver (13) who found it "on some species of bog moss in Geneva Creek Canvon, Colorado, at an elevation above 8000 ft." Only one collection of M. muscicola was reported by Durand in his monograph "on mossy stems (Webera nutans) . . . at about 7000 feet elevation, Laggan, Alberta." In 1915 Durand (4) also collected it "on wet moss at the water's edge, Lake Agnes, Alberta." In the United States it is known only from collections made by Kauffman (8) at Tolland, Colorado, at 9,500 feet and Leal, Colorado, at 8,600 feet. The two species are therefore rare for North America. Durand (4) states that he searched for them carefully but in vain at various points along the Alaskan coasts as far north as Skagway. In the United States only the three collections from Colorado have been made. The writer has sought for them for a number of years without success. However, in 1940 a Mitrula was found in moss around the margins of pools near the Hidden Lake Trail a short distance from Logan Pass. The fungus probably was parasitic since it was fruiting in patches of dead moss. The following description has been prepared.

Ascomata capitate, very variable, 10–30 mm. long, the ascogenous portion irregularly globoid, obovoid, ovoid, ellipsoid, reniform, flattened globoid, or occasionally almost cylindric or spathulate, 2–6 × 1.5–7 mm., smooth, convoluted, somewhat cerebriform or ridged when fresh, the ridges and convolutions more pronounced when dry, light orange-yellow to orange-buff (Ridgway) when fresh, darker when dried, cinnamon-rufous, the stipe slender, 1 mm. wide, equal or somewhat wider above, concolorous with the head or somewhat lighter, smooth when fresh, longitudinally furrowed when dried; asci clavate, 60–74 × 6–7  $\mu$ , pore staining blue with iodine, the ascospores fusoid-cylindric, 9–12 × 2–2.5  $\mu$ , oblique or biseriate, the paraphyses filiform, slightly wider above, 1.5–2  $\mu$ , as long as the asci.

Growing in and apparently killing moss, Aulacomnium palustre (Web. & Mohr.) Schwaegr. around margins of pools, Trail to Hidden Lake, Logan Pass, Glacier National Park, Montana, 8–20–41, E. B. Mains (6117).

The Montana collection is similar to those reported by Kauffman (8) as M. muscicola from Colorado. Since he did not give a description the following has been prepared from his collections in the Herbarium of the University of Michigan and from his notes.

f bog

above

ed by

. . .

irand

gnes,

tions and rare them as far tions them trula dden prob-

ogerenithuform nced when mm. head wed with e or

stre 1 to -20-

man e a

s in

tes.

Ascomata capitate, 10–20 mm. long, the ascogenous portion obovoid to subgloboid, 3–5 × 3–4 mm., pale ochraceous, darker when dried, even, rugose or lacunose-wrinkled, sharply differentiated from the stipe, the stipe slender, even, creamy-whitish, subpellucid, elastic, glabrous when fresh, yellowish brown and longitudinally furrowed when dried; asci clavate, 65–90 × 6–9  $\mu$ , the pore staining blue with iodine, the ascospores fusoid-cylindric, 10–14 × 2  $\mu$ , biseriate, the paraphyses filiform, slightly thickened above.




Fig. 1. Mitrula gracilis from Glacier National Park, Montana. A. Ascomata arising from moss, approximately  $1 \times$ ; B-E. Ascomata showing some of the variations in shape and surface,  $5 \times$ .

Growing on living moss in a swampy ground, Leal, Colorado (8,600 ft.), Aug. 11, 1917, C. H. Kauffman; on mossy stump, Tolland, Colorado (9,500 ft.), Sept. 14, 1920, C. H. Kauffman.

According to Kauffman's notes the stipes of his collections were much lighter in color than those of the Montana collection. When dried they show little if any difference.

Mitrula gracilis and M. muscicola do not differ greatly according to the descriptions of Karsten (7) and Henning (6). In treatments where the species are separately maintained the principal differences noted are that M. gracilis has an orange-brown more or less even head and a lighter colored stipe whereas M. muscicola has a more convoluted, cinnamon-brown head and concolorous stipe. Such variations may occur within a collection, as in the Montana collection. Although Durand treated them as separate species he did so doubtfully. Seaver also has expressed considerable doubt concerning their separation. Nannfeldt (12) in his recent article concerning the Geoglossaceae of Sweden has concluded that they are synonymous. With this the writer agrees. M. gracilis appears to have been more frequently collected in Europe than in North America, although it is far from common. Its arctic and high alpine distribution probably accounts to some extent for the few collections.

### LITERATURE CITED

- 1. De Candolle, M. Flore française 6: 660 pp. 1815.
- Dietel, P. Kleine Beiträge zur Systematik der Uredineen. Ann. Mycol. 20: 29-33. 1922.
- Durand, Elias J. The Geoglossaceae of North America. Ann. Mycol. 6: 387-477. 1908.
- 4. —. New or noteworthy Geoglossaceae. Mycologia 13: 184-187.
- Fischer, Ed. Die Uredineen der Schweiz. Beiträge Kryptogamenfl. Schweiz. 2, Heft 2; xciv + 590 pp. 1904.
- Henning, Ernst. Bidrag till svampfloran i Norges sydligare fjelltrakter. Overs. Kongl. Vet. Akad. Fördh. 1885. No. 5: 49-75.
- Karsten, P. A. Fragmenta mycologica scripsit I. Hedwigia 22: 17-18, 1883.
- Kauffman, C. H. The mycological flora of the higher Rockies of Colorado. Papers Mich. Acad. Sci. Arts, Letters 1: 101-150. 1921.
- Klebahn, H. Kulturversuche mit Rostpilzen XII. Zeits. Pflanzenkrank. 15: 65-108. 1905.

- 10. . Uredineen. Kryptogamfl. Mark Brandenburg 5A: 69-904.
- Moss, E. H. The uredinia of Melampsora and Coleosporium. Mycologia 21: 79-83. 1929.
- Nannfeldt, J. A. The Geoglossaceae of Sweden. Arkiv Bot. 30A, No. 4: 1-67. 1942.
- Seaver, Fred J. Studies in Colorado fungi—I. Discomycetes. Mycologia 3: 57-66. 1911.
- Sydow, P. & H. Melampsora. Monographia Ured. 3: 334-393.
   1915.
- 15. —. Uredo. Monographia Ured. 4: 384-612. 1924.
- Winter, Georg. Die Pilze Deutschlands, Oesterreich und der Schweiz in Rabenhorst's Kryptogamen-Flora I Abt., Band I. 924 pp. 1881–1884.

Ann.

lorado

stump,

ffman.

s were

When

ccord-

). In

prin-

brown

as M. I conon, as em as ressed
(12)
n has grees.
n Eumon.
some

Ann.

–187. nenfl.

fjell-

9-75. 22:

es of

-150. izen-

# SIX NEW INDIAN DISCOMYCETES

EDITH K. CASH

Among specimens of Discomycetes collected by Sultan Ahmad in various localities in India and Pakistan and sent to the writer for examination during the past few months, a specimen of Dasyscyphella, two of Humaria, and three of Humarina appear to differ from any species of these genera that could be found in the literature and are therefore named here as new. Type specimens of these fungi are deposited in the Mycological Collections of the Bureau of Plant Industry, Beltsville, Maryland, and duplicate material of some of them in the herbarium of the collector at Lahore, Pakistan.

### 1. Dasyscyphella indica sp. n.

Apothecia substipitata, ceracea, turbinata, flava, 1-2 mm. diam., margine et extus e pilis echinulatis, hyalinis,  $40-50 \times 3-4 \mu$  albopruinosa; asci cylindrici, apicibus attenuatis, octospori,  $100-120 \times 7-8 \mu$ ; ascosporae hyalinae, anguste clavatae, multiguttulatae,  $35-45 \times 2 \mu$ ; paraphyses tenues, usque  $2 \mu$  inflatae; excipulum hyalinum, plectenchymaticum.

Apothecia substipitate, waxy, turbinate, 1–2 mm. in diameter, hymenium plane, Capucine yellow, ferruginous when dry, exterior concolorous, white-pruinose, margin fimbriate; asci cylindrical, narrowed and with wall thickened at the tips, gradually attenuated to a short pedicel, eight-spored,  $100-120\times7-8~\mu$ ; ascospores hyaline, narrow-clavate, multiguttulate, acute at the lower end, often arcuate or sigmoid when free from the ascus,  $35-45\times2~\mu$ ; paraphyses numerous, delicate, only slightly enlarged at the tips up to  $2~\mu$ ; exciple hyaline, plectenchymatous, hairs echinulate, hyaline or subhyaline,  $40-50\times3-4~\mu$ .

On mossy bark of trees, Mussoorie, July 20, 1940, S. Ahmad 434. D. indica differs from Erinella corticola Mass., described from India (Kew Bull. 1898, p. 115, 1898), in the flexuous, instead of fusoid paraphyses, and from several other somewhat similar species of Erinella in the dimensions of the asci and spores.

<sup>&</sup>lt;sup>1</sup> Except where noted, color readings are from Ridgway, R. Color standards and nomenclature, Washington, D. C., 1912.

# 2. Humaria ahmadii sp. n.2

Apothecia sessilia, carnosa, 1–3 mm. diam., rubra, patelliformia; asci cylindrici, octospori, ad apices rotundati, 180–190 × 18–20  $\mu$ ; ascosporae hyalinae, late ellipsoideae, uniseriatae, verrucosae, 2-guttulatae, saepe solum quatuor maturascentia, 15–20 × 9–12  $\mu$ ; paraphyses flexuosae, numerosae, ad apices usque 5–6  $\mu$  incrassatae; setae paucae, pallide brunneae, subacutae, simplices vel 1–2-septatae, 75–100 × 8–11  $\mu$ .

Apothecia sessile, fleshy, 1–3 mm. in diameter, bright red,<sup>3</sup> fading to pale olive gray or nearly white in dried specimens, patelliform, margin nearly even, excipular hairs short, inconspicuous; asci cylindrical, rounded at the apex, gradually attenuated toward the base, eight-spored, 180–190 × 18–20  $\mu$ ; ascospores hyaline, broadly elliptical, uniseriate, coarsely verrucose, two-guttulate, usually only four becoming mature,  $15–20 \times 9–12 \mu$ ; paraphyses numerous, flexuous, unbranched, enlarged at the tips to  $5–6 \mu$ ; setae inconspicuous, pale brown, subacute, simple or 1–2-septate,  $75–110 \times 8–11 \mu$ .

On the ground, Lahore, Pakistan, Nov. 24, 1947, S. Ahmad 2226. The very short, inconspicuous setae constitute the most distinctive character of this species. It appears to be close to *Cheilymenia calvescens* Boud., from which it differs in the shorter setae and coarsely verrucose, rather than minutely echinulate spores.

# 3. Humaria pallidisetosa sp. n.

Apothecia patelliformia, carnosa, 6–8 mm. diam., avellanea, margine denticulato; asci cylindrici, apicibus subobtusis, octospori,  $200-250\times13-15~\mu$ ; ascosporae uniseriatae, hyalinae, late ellipsoideae, glabrae,  $15-18\times9-11~\mu$ ; paraphyses numerosae, granulosae, ad apices 3–4  $\mu$ ; setae pallidissime brunneae vel subhyalinae, flexuosae, angustae, usque  $250-300\times5-8~\mu$ , glabrae vel minute echinulatae, septatae.

Apothecia patelliform, soft fleshy, 6–8 mm. in diameter, wood brown to drab, drying cinnamon drab, margin minutely denticulate; asci cylindrical, slightly flattened at the tips and narrowed below, eight-spored, 200–250 × 13–15  $\mu$ ; ascospores uniseriate, hyaline, broadly ellipsoid, smooth, 15–18 × 9–11  $\mu$ ; paraphyses numerous, granulose, 3–4  $\mu$  at the apices; setae very pale brownish to subhyaline, flexuous, narrow, smooth to finely echinulate, septate, narrowed and rounded at the tips, 250–300 × 5–8  $\mu$ .

<sup>2</sup> As pointed out by Kanouse (Mycologia 39: 655. 1947), Humaria is the valid generic name under the International Rules of Nomenclature for the species generally known as Lachnea or Patella.

<sup>3</sup> Ridgway color reading not made on fresh material.

hmad er for yscydiffer itera-

f the mahore,

ns of

rgine cylinlinae, e 2 µ

hyerior rical, ated hyoften arap to e or

rom d of spe-

and-

On the ground, Rohtak, Punjab, Jan. 17, 1947, S. Ahmad 1776. The fungus is close to *Patella gilva* (Boud.) Seaver in dimensions, but differs from the latter in the nearly hyaline setae and in the color of the hymenium.

### 4. Humarina plumbeo-atra sp. n.

Apothecia cupuliformia, sessilia vel partim immersa, carnosa, glabra, margine leniter crenata, 1.5–2 mm. diam., violaceo-ardosiaca usque plumbeo-atra; asci cylindrici apicibus obtusis, octospori, 130–175 × 12–15  $\mu$ ; ascosporae uniseriatae, late ellipsoideae, glabrae, utrinque guttulatae,  $13–14\times7–9~\mu$ ; paraphyses tenues, ascos superantes, apicibus 2.5–3.5  $\mu$  crassis; excipulum ex hyphis tenuibus plectenchymatice intertextis, extus cellulis angulosis  $10–15~\mu$  diam. compositum.

Apothecia sessile or partly immersed in soil, fleshy, smooth, margin slightly crenate, 1.5–2 mm. in diameter, hymenium and exterior dark violet slate to plumbeous black, flesh purple when crushed; asci cylindrical, obtuse at the tips, gradually attenuated below, eight-spored, 130– $175 \times 12$ – $15 \,\mu$ ; ascospores uniseriate, broadly ellipsoid, smooth, with one small guttule at each end, 13– $14 \times 7$ – $9 \,\mu$ ; paraphyses slender, longer than the asci, gradually enlarged to 2.5– $3.5 \,\mu$  at the tips; exciple composed of fine plectenchyma within, the outer layer of thin-walled, subglobose to angular cells 10– $15 \,\mu$  in diameter.

On the ground beside a water course, Ladhar, Sheikhupura, Punjab, July 8, 1946, S. Ahmad 1640.

Among the species nearest in color of the hymenium, *Humarina* purpurea Seaver has rough spores, and both *Humaria plumbea* Fr. and *Peziza violacea* Pers. ex Fr. have much larger apothecia.

# 5. Humarina umbrina sp. n.

Apothecia patelliformia usque plana, aliquantus plicata, carnosa, umbrina, 7–8 mm. diam., extus glabra vel minute pustulata, margine subundulato; asci cylindrici, apicibus obtusis, octospori,  $200-250 \times 11~\mu$ ; ascosporae uniseriatae, hyalinae, ellipsoideae, glabrae,  $12-15 \times 7-9~\mu$ ; paraphyses flexuosae, apicibus usque  $4-5~\mu$  incrassatis et dense granulosis; excipulum e cellulis tenuibus, subglobosis usque  $25~\mu$  diam. compositum.

Apothecia patelliform to applanate, sessile, more or less plicate, fleshy, 7–8 mm. in diameter, hymenium Saccardo's umber, drying sepia, exterior concolorous, smooth to minutely pustulate, margin slightly undulate; asci cylindrical, obtuse at the tips, narrowed below, eight-spored,  $200-250 \times 11~\mu$ ; ascospores uniseriate, hyaline,

ellipsoid, smooth,  $12-15 \times 7-9 \mu$ ; paraphyses flexuous, unbranched, tips swollen to  $4-5 \mu$  and filled with fine yellowish granules; exciple of thin-walled hyaline subglobose cells,  $25 \mu$  in diameter.

On the ground, Lahore, Pakistan, Nov. 24, 1947, S. Ahmad 2225. H. umbrina may be distinguished from Peziza saccardiana Cke. by the smooth spores. In P. sepiatra Cke. and P. sepiatrella Sacc., two species also similar in color, the spores are decidedly longer.

# 6. Humarina zizyphi sp. n.

Apothecia obconica usque turbinata, substipitata, 1–2 mm. diam., margine subundulato, sicco involuto, armeniaco-lutea usque ochraceo-aurantia, sicca daucino-ochracea; asci cylindrici apicibus deplanatis, longe pedicellati, octospori,  $100-150\times13-15\,\mu$ ; ascosporae uniseriatae, hyalinae, glabrae, oblongo-ellipsoideae, guttulis parvis numerosis impletae,  $20-25\times10-11\,\mu$ ; paraphyses abundantes, hyalinae, granulosae, ad apices  $2-3\,\mu$  inflatae, circa  $50\,\mu$  infra apices ramosae; excipulum subhyalinum e cellulis irregulariter angulosis  $15-25\,\mu$  in diam. compositum.

Apothecia obconic to turbinate, substipitate, 1–2 mm. in diam., margin slightly undulate and inrolled when dry, hymenium apricotbuff to ochraceous orange, carrot red to ochraceous tawny when dry, exterior concolorous, smooth; asci cylindrical, flattened at the tips, long pedicellate,  $100-150\times13-15\,\mu$ ; ascospores hyaline, smooth, oblong-ellipsoid, filled with many small oil globules,  $20-25\times11-12\,\mu$ ; paraphyses abundant, hyaline, granulose, branched about  $50\,\mu$  below the tips, slightly swollen to  $2-3\,\mu$  at the tips; exciple subhyaline, composed of irregularly angular cells  $15-25\,\mu$  in diameter.

On stones and on dead branches of Zizyphus jujuba, Ladhar, Sheikhupura, Punjab, Sept. 23, 1941, S. Ahmad 419.

In the form and size of the spores this fungus resembles *Humarina waterstonii* Seaver described from Bermuda on seeds of *Livistona chinensis* (Mycologia 31: 533. 1939), but differs from that species in the host plant and in the paler hymenium.

U. S. PLANT INDUSTRY STATION, BELTSVILLE, MARYLAND

maratra; e uni-; pam ex 0-15  $\mu$ 

1776.

men-

nd in

marl exwhen uated riate, 3–14 arged hyma cells

arina a Fr.

Pun-

brina, ; asci riatae, picibus uibus,

icate, rying argin d bealine,

# A STUDY OF THE TEMPERATURE AND HUMIDITY REQUIREMENTS OF ASPERGILLUS NIGER 1

JOHN TYLER BONNER

(WITH 3 FIGURES)

There are a number of reasons why the temperature and humidity relations of fungi are important. First they are of physiological interest for they affect and limit the fundamental processes of growth and development. Also, of a more practical nature, they both provide information helpful in the problem of fungus or mildew control, and they are useful in laboratory tests of vulnerability or resistance of items to fungus attack, where the optimum conditions of temperature and moisture must be known.

Aspergillus niger was chosen as the organism for this study both because of its widespread occurrence and because of its frequent use as a test organism. (The latter is especially true of the strain chosen: United States Department of Agriculture No. Tc 215–4247.)

#### MATERIALS AND METHODS

The technique used was basically that of Galloway (1935). Briefly, small squares of cellophane streaked with fungus spores were suspended on threads over solutions of different concentrations of calcium chloride, giving atmospheres of different relative humidities. Since details of the methods play an important role in the reproducibility of results, they will be described here.

Aspergillus niger was grown in Petri dishes on the following media: Dextrose, 30 gm.; NaNO<sub>3</sub>, 30 gm.; K<sub>2</sub>HPO<sub>4</sub>, 1.0 gm.; MgSO<sub>4</sub>, 0.25 gm.; KCl, 0.25 gm.; agar, 15 gm.; distilled water,

<sup>1</sup> This study was carried out during the war as part of a tropical deterioration research program at the Aero-Medical Laboratory, Wright Field, Dayton, Ohio.

The author wishes to express his gratitude for the invaluable technical assistance of Mrs. Roberta Dingwall.

1000 ml. (pH adjusted to 6.8). The spore load was obtained from cultures that had grown at room temperature from seven to fourteen days.

D

huysioesses ture, is or

lnernum

both

uent train 215–

35). ores atraative role


wing gm.;

ater, iora-Day-

nical

16 OZ. FRENCH \_ SQUARE BOTTLE

The squares, onto which the spores were placed, were made



METHOD OF PLACING CELLOPHANE SQUARES IN TEST CHAMBERS

CA CL, SOLUTION

Fig. 1. Diagram illustrating the technique used.

from strips of cellophane  $^2$  which had been immersed in five per cent malt extract (Difco) solution for ten minutes, dried in room air, and cut into squares  $1 \times 1$  cm. Each square was strung onto a piece of number fifty white cotton thread and held by a simple knot. They were not sterilized because the germinating spores could always be identified as *Aspergillus niger* under the microscope. In the few instances where contaminants appeared, the squares were discarded.

To place a streak of spores on a square, the cellophane square was folded back so that two corners met and the fold was brought in contact with the surface of the fungus culture. This fold,

### TABLE 1

CONCENTRATION OF CALCIUM CHLORIDE SOLUTIONS (AS DETERMINED BY THE HARVEY (1910) METHOD) AND THE ESTIMATED RELATIVE HUMIDITY OVER THESE SOLUTIONS (OBTAINED FROM DAFA OF THE DOW CHEMICAL COMPANY)

Concentration of Solution	Estimated Relativ			
26.5%	70%			
21.9%	78%			
17.1%	85%			
10.0%	93%			

studded with spores, was then rubbed gently against a piece of filter paper to even their distribution. The inoculated squares were than suspended over various concentrations of calcium chloride solution in 16 oz. French square bottles so that there were 4 squares in each bottle (Fig. 1).

Four different concentrations of calcium chloride solutions were prepared and each solution was divided so that 30 ml. were dispensed in a number of the 16 oz. French square bottles. The concentrations of the solutions were carefully checked by the Harvey (1910) technique. From information generously supplied by the Dow Chemical Company (1945) it was known what relative humidities could be expected over such solutions. This information is recorded in table 1.

The bottles containing the squares were placed in an incubator in which a fan had been installed to ensure constant temperatures

<sup>&</sup>lt;sup>2</sup> The cellophane "used was Du Pont PUT-O (30 gms. per square meter) which has received no special surface coating treatment of any sort, but is merely pure, regenerated cellulose.

and to prevent any temperature gradients. The length of time for the temperature inside a French square bottle (kept at room temperature,  $24^{\circ}$  C.) to reach equilibrium with the oven temperature was determined. By use of thermocouples it was found that at the near extreme incubator temperature of  $45^{\circ}$  C., in one-half hour the temperature inside the bottle was  $1^{\circ}$  C. low, and in one hour it had reached the incubator temperature.

The technique for making the germination observations was to remove a square from the bottle and fix the spores on the cellophane for staining by holding the surface containing the spores over steam rising from boiling water for about one-half to one minute. The square was then immersed in Linder's Lacto-phenol-cotton-blue, which was used as a mounting and staining fluid, and heated gently on a slide, under a cover-slip, to flatten out the cellophane.

The germination times were established by making stained preparations of replicates of any one experiment at different intervals of time approaching the expected germination time as determined by preliminary tests. The four sets of slides of each determination were carefully examined microscopically and an estimated germination time was established. The times are expressed  $a \pm b$ . The a indicates the most probable time;  $\pm b$  the range of time within which germination certainly occurs (TABLE 2).

In this study "germination" is defined as that stage of development where the sides of the germ tube are first parallel, and "germination time" is defined here as the length of time required for at least ten spores to germinate on a cellophane square. This is contrary to the procedure used by many (see, for instance, Wellman and McCallan, 1942) where the per cent germination is plotted against time, and "germination time" is arbitrarily chosen as the time when fifty per cent, or some other per cent, germination has occurred. The reason for not using such a method is twofold: (1) it is difficult to get an even distribution of spores on the cellophane squares when using Galloway's (1935) technique, which is necessary to establish per cent germination, but more important, (2) from the practical point of view it is of greater value to know when the first spore germination will occur than to know the average time for all spores. This point was clearly shown, when in some

room onto mple

ores

the uare

fold,

e of

vere dis-The the supwhat

ator ures

at is

instances the per cent germination remained extremely low, but the few that did germinate grew and sporulated. (For example see figure 3, F.)

IX

v

fı

### RESULTS

The germination time of Aspergillus niger USDA No. Tc 215–4247 was determined at 10°, 20°, 30°, 40°, 45°, 50° C. and at the following relative humidities for each of those temperatures: 100%, 93%, 85%, 78%, 70%. Most determinations were checked in at

TABLE 2
GERMINATION TIME OF Aspergillus niger at DIFFERENT
TEMPERATURES AND RELATIVE HUMDITIES

Temperature in °C.	Relative Humidity	Estimated Germination Time in Hours	Number of Experiments Performed		
10	100	>100	1		
20	100	12 ±1	4		
20	93	12 ±1	4 4		
20	85	17 ±3	4		
20	78	>100	1		
30	100	4 ±1	4		
30	93	5.5±1	4		
30	85	9 ±1	4		
30	78	48 ±3	4		
30	70	>100	1		
40	100	5.5± .5	4		
40	93	3.5± .5	4		
40	85 -	8 ±2	4		
40	78	33 ±3	4		
40	70	>100	1		
45 .	100	17 ±5	2		
45	93	7 ±2	4		
45	85	9 ±2	4		
45	78	50 ±5	2		
45	70	>100	1		
Mi.					
50	100	>100	1		

least four separate experiments. This checking was found necessary after preliminary experiments which showed that the variability of results was large. However, in a few instances the results were sufficiently definite to warrant fewer checks. The data from these experiments are recorded in table 2.

In order to appreciate fully the results given in table 2 they must be displayed in the form of a graph, but since there are three variables, ideally a three-dimensional graph is required. Taking heart from the ecologists (see Shelford, 1929) who have been confronted with the same problem, a two-dimensional graph has been




Fig. 2. Germination curves.

but the

at the 100%, l in at

of nts ed

cesarire-

relata drawn (FIG. 2) in which the abscissa is relative humidity, the ordinate temperature, and "germination time" is represented by a family of curves.

The salient characteristics in relation to temperature and humidity that can be observed for Aspergillus niger in figure 2 are primarily the following: the optimum conditions for growth are at a relative humidity near 93 per cent, and at a temperature near 40° C; at 100 per cent relative humidity the optimum temperature is near 30° C. Thus we find that the optimum condition for growth for this species is not at saturation and that the optimum temperature varies with different humidities.

It should be clearly understood, when interpreting the graph in figure 2, that for two reasons the accuracy of the curve is limited: (1) as stated above the points on the graph are the "estimated" germination time within a range of time where it is certain germination occurred. Although these "estimated" points are believed to be fairly accurate, they are to a limited extent arrived at by human judgment and not entirely by objective measurement. (2) There are relatively few points on the graph and therefore there has been a considerable amount of interpolation and extrapolation and the possible error is multiplied by the fact that it is in three dimensions.

During the course of these experiments it was noted that the different conditions imposed on the germinating spores of Aspergillus niger definitely affected its morphology. It was especially true that under extreme conditions abnormalities in pattern occurred. These abnormalities were of two types. One occurred at high temperature, 45° C., and at high humidities, 100 per cent and 93 per cent, where the spores were greatly swollen (FIG. 3, A). Under optimal conditions slight swelling occurred (FIG. 3, B), but in these extreme cases the spore swelled to over ten times its normal volume. This swelling in no way prevented germination, as evidenced in figure 3, A. At a higher temperature, 50° C., where no germination occurs, there was no evidence of any swelling, in fact the spores showed signs of degeneration. The second type of abnormality appeared at low relative humidities, 70-78 per cent at all viable temperatures. In these instances the spore appeared to be of normal diameter, but the germ tube was enlarged (FIG.

Fig. 3. Germinating spores.

, the

l hu-2 are n are near ature n for mum

oh in ited: ated" ermi-eved at by (2) e has and e di-

difrgiltrue red. emper nder nese volevinere , in

red FIG. 3, C), sometimes even spherical (FIG. 3, D), rather than the normal thread-like construction. If the experiment was allowed to run for extended periods of time, often different types of peculiar segmented, flat, fanlike hyphal clusters resulted (FIG. 3, E, F). This apparently does not inhibit sporulation, as seen in figure 3, F where a sporangiophore rises from the center of the abnormal mycelium.

#### DISCUSSION

A comparison has been made between the results obtained here on Aspergillus niger with those to be found in the literature. Very little complete work has been done on the temperature and humidity requirements of any one fungus. Although Groom and Paniset (1933) give some data, the only work I have found that is sufficiently parallel to this study so that comparisons can be fruitfully made is that of Tomkins (1929).

Tomkins made detailed studies of two fungi: Alternaria citri and Trichoderma lignorum. His results, which are essentially similar for both of these fungi, differ in three major respects from Aspergillus niger: (1) The optimum relative humidity for Tomkins' fungi is 100 per cent, whereas for Aspergillus niger it is nearer 93 per cent. (2) In his fungi the optimum temperature is the same for all relative humidities, which is not true for Aspergillus niger. (3) In his fungi the temperature minimum becomes lower the higher the relative humidity, whereas in Aspergillus niger the temperature minimum is approximately constant between relative humidities of 85 to 100 per cent.

The reason for these major differences could be found in a number of directions. Tomkins' technique is very different—for instance he measures growth rate instead of germination time. Furthermore his actual testing conditions differ radically from those used here.

But also it is quite conceivable that different fungi vary in their response to temperature and humidity. To examine this point a survey was made of the minimum, optimum and maximum temperatures and humidities of numerous fungi. Briefly the following information on temperatures was obtained:

e nor-

red to

culiar

, F).

3, F

ormal

here

Very

l hu-

and

nat is

ruit-

and

nilar

rgil-

ungi

per

(3)

gher

ture

ities

um-

in-

ur-

iose

neir

t a

ering In a fairly representative group of mildews the range of minimum temperature for forty-four fungi is from  $-6^{\circ}$  C. to  $15^{\circ}$  C., the range of optimum temperatures for fifty-one fungi is from  $10^{\circ}$  C. to  $45^{\circ}$  C., and the range of maximum temperatures for thirty-two fungi is from  $25^{\circ}$  C. to  $51^{\circ}$  C.

Similar data on humidities show that the minimum relative humidity for fifty-eight fungi is from approximately 70 per cent to approximately 98 per cent. Furthermore, the frequency distribution of minimum relative humidities for the fifty-eight fungi is fairly even between the limits of the range.

We have then ample evidence that fungi differ greatly in their response to temperature and humidity.

Unfortunately the basic problem of how temperature and humidity affect the processes of growth and development in any fungus, and what it is in different fungi that is affected so specifically by these environmental conditions, remains completely obscure. If anything is indicated it is that the relationship between the three variables, (1) species of fungus, (2) temperature, and (3) humidity, must be extremely complex. The results on *Aspergillus niger* will serve only to expose more fully the fundamental problem though their practical value will be more immediate.

#### SUMMARY

The combined temperature and relative humidity requirements of Aspergillus niger were studied with the following results: The optimum conditions for growth are at a relative humidity near 93 per cent and at a temperature near 40° C.; at 100 per cent relative humidity the optimum temperature is near 30° C. Thus for this species the optimum condition for growth is not at saturation and the optimum temperature changes with different humidities.

A general survey of the literature shows that the variability among fungi in their response to temperature and humidity is great, indicating that there is no simple temperature-humidity relationship for fungi.

DEPARTMENT OF BIOLOGY,
PRINCETON UNIVERSITY,
PRINCETON, NEW JERSEY

### LITERATURE CITED

- Dow Chemical Company. 1945. Personal communication.
- Galloway, L. D. 1935. The moisture requirements of mould fungi with special reference to mildew to textiles. Journ. Text. Inst., Manchester 26: T123-T129.
- Groom, P., and Paniset, T. 1933. Studies on Penicillium chrysogenum
  Thom in relation to temperature and relative humidity of the air.
  Ann. Appl. Biol. 20: 633-660.
- Harvey, S. C. 1910. The quantitative determination of the chlorids in the urine. Arch. Int. Med. 6: 12-18.
- Longrée, Karla. 1939. The effect of temperature and relative humidity on the powdery mildew of roses. Cornell Univ. Agr. Exper. Sta. Mem. 223, 43 pp.
- Shelford, V. E. 1929. Laboratory and field ecology. Baillière, Tindall, and Cox, London.
- Tomkins, R. G. 1929. Studies on the growth of molds. Proc. Roy. Soc. B105: 375-401.
- Wellman, R. H., and McCallan, S. E. A. 1942. An analysis of factors causing variation in spore germination tests of fungicides. IV. Time and temperature. Contrib. Boyce Thompson Inst. 12: 431-450.

### **EXPLANATION OF FIGURES**

- Fig. 2. Graph showing the effect of temperature (ordinate) and relative humidity (abscissa) on the germination time (family of curves) of Aspergillus niger.
- Fig. 3. Photomicrographs showing the effect of different combinations of temperature and humidity on the morphology of the germinating spore of Aspergillus niger. A.  $(300 \times)$  Spores germinating at high temperature  $(45^{\circ}$  C.) and high relative humidity (100%). B.  $(300 \times)$  Spores germinating under optimal conditions  $(40^{\circ}$  C., 93% RH). C, D.  $(300 \times)$  Spores germinating at low relative humidities (70-78%) and optimal temperatures  $(30-40^{\circ}$  C.). E, F.  $(75 \times)$  Growth obtained if the fungus is kept for an extended time (3-5 weeks) under dry conditions (70-78%) RH) at an optimal temperature  $(30^{\circ}$  C.). Note how the hyphae spread out into fanlike processes and that in F normal sporangiophores can be seen.

# TWO SPECIES OF COPRINUS WITH NOTES ON THEIR CULTURAL CHARACTERS

fungi

Inst.,

genum le air.

lorids

ımid-

xper.

Tin-

Roy.

ctors

IV. 12:

ative

sper-

is of

e of

ture

inat-

ores

ures

an

an

fan-

MORTEN LANGE 1

(WITH 4 FIGURES)

The present paper deals with two species of *Coprinus*; the one being identified with a previously described but little known species, the other believed to be previously undescribed. Both have been grown in culture and notes are given about their cultural characters. The descriptions, however, are drawn from material grown under natural conditions, if not otherwise stated.

COPRINUS COTHURNATUS Godey ap. Gillet 1878. (Fig. II, III)

Pileus 1.2-2.0 cm. high before expanding, ovate, subovate, expanding through campanulate to almost flat, slightly umbonate, edge recurving and splitting irregularly; whitish then grayish, plicate-striate nearly to the centre, disc not sharply delimited. Veil rather prominent on young specimens, of a dense, granulose to somewhat filamentous covering (especially towards the margin, which on young buds is connected with the stipe by fibrils), dingy whitish, in places with a reddish avellaneous tinge (conf. below), on mature specimens left as a few scurfy, brownish patches on top Flesh very thin, watery white. Lamellae free, narrow, rather crowded, with a varying number of smaller ones between; white, turning black over all (through pale pinkish cinnamon); spores ripening simultaneously, edge floccose, remaining white until deliquescing. Stipe 5-6 cm. long, 3-5 mm. thick, terete or slightly compressed, attenuated upwards, white, scabrose-tomentose from short fibrils (very much so towards the base which is also covered with distinct, granulose veil-remnants and tinged reddish avellaneous), distinctly hollow, not fragile. Odor and taste nauseating, somewhat like Coprinus narcoticus, rather pronounced. Sporeprint black.

<sup>1</sup> University of Copenhagen, Botanical Laboratory. The main part of this work was carried out while the author was studying in the laboratory of Dr. A. H. Smith, University of Michigan Herbarium, Ann Arbor, Michigan, United States of America.

Spores  $11.6-14.2 \times 7.6-9.5 \times 8.3-10 \,\mu$ , mostly  $12.5-13 \times 8-9 \,\mu$ , broadly ovate, usually slightly flattened, some of them obscurely angular in face view, dark umber in H<sub>2</sub>O (coal black in KOH), truncate, germ pore conspicuous, slightly eccentric, about 2.5 µ broad. Basidia 4-spored, more or less dimorphic, 30-45 × 10-11 μ, sterigmata 2-3 μ long. Paraphyses pear-shaped. Pleurocystidia  $18-30(50) \times 15-40 \,\mu$ , scattered to rare, vesiculose and soon collapsing. (All material raised in culture has been absolutely devoid of pleurocystidia). Edge of gills made up of pearshaped, subglobose or globose cells, 15-35 µ broad, borne on thornlike projections on cylindric filaments 3-8 µ broad. Gill trama interwoven, cells about 3-5 u broad. Veil of cap similar to the tissue of the gill edge but inflated cells averaging somewhat larger (15-80 µ), slightly incrusted, readily collapsing, often with brownish amorphous material between. Clamp connections demonstrated on mycelium (see below) but not in fruitbody.

Habit, habitat and distribution. Fasciculate, in large clusters, on old, decaying haystack. May 21–25, 1947, Arboretum, Ann Arbor, Mich., U. S. A., leg A. H. Smith and M. Lange (Lange 718, 719). Previously recorded from France, Switzerland and England.

Observations. The specimens grown in vitro show a very prominent veil, which on young buds forms conspicuous warts. Stem and veil both turn brick red a few seconds after being touched and after some minutes fade to vinaceous brown or avellaneous. These characters are more difficult to ascertain on specimens found in nature, but are very striking on those produced in culture.

The peculiar color reaction of the veil and stem gives *C. cothurnatus* some similarity to *C. dilectus* and *C. roseotinctus*, but I do not believe there is any close relationship since these species have another type of veil. *C. cothurnatus* evidently belongs in the group of *C. niveus* and *C. semilanatus* by virtue of similar veil and spore characters, and appears to be very close to the latter, from which the changing color of the veil and differences in shape and size of spore distinguish it. It is not without some doubt that I give this interpretation of *C. cothurnatus*, a species badly treated or ignored in modern literature, but the original description, as given by Saccardo, fits my plant well, except for the mention of the gills turning black through "flesh color." No detailed description of the microscopic characters of *C. cothurnatus* is known to me. Al-

though the plant originally was recorded from cow dung, Martin (1904–05) finds it on humus, in the spring, thus better corresponding—if correctly interpreted—to the present find. I have had no access to Gillet's work (l. c.) but have seen a copy of the illustra-

3-9 µ,

urely

OH), 2.5 μ

< 10euroand

absopearhornrama o the arger own-

rated

Ann ange and

Stem l and These

n na-

I do have roup pore

ze of


this ored

Sac-

urn-

the

Al-



Figs. I-II. Microscopic characters of Coprinus cothurnatus and C. myceliocephalus.

tion (kindly made for me by Dr. J. Favre in Geneva)—it very much recalled the present plant.

Cultural characters. Spores spread on horse dung agar germinate readily in less than 24 hours at 18° C. (almost 100%). Poly-

spored mycelium is provided with clamp connections. Single-spore mycelia were isolated and mated together (TAB. 1). The species, accordingly, is heterothallic and bipolar. Fruitbodies form rather readily about a month after the inoculation both on agar in test tubes and on sterilized horse dung.

#### TABLE 1

Coprinus cothurnatus God. RESULT OF PAIRING OF 10 HAPLOID MYCELIA. + indicates formation of clamp connections and spore producing fruitbodies.

	A	C	D	H	F	E'	G	В	I	J
A		-	-	_	+	+	+	+	+	+
C	-		_	_	+	+	+	+	+	+
D	_	_		_	+	+	+	+		
H		-	-		+	+	+	+		
HFEG	+	+	+	+		_	_	_		
E	+	+	+	+	_		_	-		
G	+	+	+	+	-	_		_		
B	+	+	+	+	-	-	-		_	_

### Coprinus myceliocephalus sp. nov. (Fig. I, IV)

Pileus 0.4–0.8 (2.0) cm. altus, initio globosus, postea ex ovato expansus, velo universali valido, albo, in squamas adpressas lacerato obtectus, sub velo striatus, ex pallide cinnamomeo nigricans. Lamellae liberae, subconfertae, ventricosae, initio albae, dein cinnamomeae, tandem nigrae, marginibus flocculosis, albis. Stipes  $4\times0.2$  ( $12\times0.3$ ) cm., flocculosus, cavus, albus. Odor et sapor nulli. Sporae (12.5)  $14-17\times10-12.5$  (14)  $\times$  9–9.5  $\mu$ , subcordatae, subangulatae. Pleurocystidia  $140-175\times25-40$   $\mu$ , cylindrica, saccata. Chellocystidia  $50-150\times25-40$   $\mu$ . Basidia tetraspora. Velum universale ex hyphis mycelioideis, 2-8  $\mu$  latis contextum. **Typus** (M. Lange C. 88) in Mus. Bot. Hauniensi et in Herb. Univ. Mich. depositum, e fimo vaccinum ex Brewster Co. Texas Americae borealis habito natum aluit M. Lange, 5-5-48.

Pileus 0.4–0.8 cm. high before expanding (in culture up to 2.0 cm. high), nearly globose at first, then ovate, expanding to ± umbonate, with edge recurving and somewhat splitting at maturity. Veil prominent, a thick, white, fibrillose coating, breaking up in adpressed patches during expansion and still conspicuous on deliquescent cap, the larger patches especially on and around the umbo. Surface of cap exposed between patches when expanding, somewhat viscid, watery cinnamon-brown, turning black through grayish brown. Edge of cap fibrillose. Flesh whitish, thin. Lamellae free, moderately crowded, 2–3.5 mm. broad, white at first, soon pale grayish brown, then blackening through ashy gray. Edge prominently white floccose at first, deliquescing from the edge. Spores ripening simultaneously on whole gill. Stipe 4 cm. long, 2 mm. thick (in culture up to 12 × 0.3 cm.), white; young stipe densely clad by a white filamentous coating, when old with

more scattered fibrils; cylindric, slightly tapering above, the base slightly bulbose, with remnants of the velum partiale; hollow, flesh whitish. Odor and taste faint. Sporeprint black.

Spores  $(12.5)14-17 \times 10-12.5(14) \times 9-9.5 \mu$ , distinctly flattened,  $\pm$  angular-subcordate in face view, elliptic in side view, impellucid, nearly black (coal black in KOH); germpore large, apical or nearly so; some few spores deformed, with two germpores. Basidia 4-spored, somewhat dimorphic, broadly clavate, short to long pedicellate,  $25-40 \times 12-14 \mu$  incl. sterigmata (about




Fig. III. Coprinus cothurnatus.

 $4\,\mu$ ). Pleurocystidia rather numerous,  $140\text{--}175\times25\text{--}40\,\mu$ , inflated cylindric, sac-shaped, apex broadly rounded to somewhat acuminate, thin-walled, readily collapsing. Cheilocystidia numerous, similar to pleurocystidia or considerably shorter, some being nearly globose,  $50\text{--}150\times25\text{--}45\,\mu$ . Paraphyses broadly pear-shaped. Gill trama not prominent, interwoven, of elongated, somewhat branched cells, generally  $5\text{--}10\,\mu$  broad, some few inflated, much broader. (A section of young bud shows a loosely interwoven tissue between gill edges and stem, of which the cheilocystidia form a part.) Veil on cap made up of interwoven hyphae

odies.

CELIA.

ingle-

e spe-

form

agar

ansus, o velo iertae, is floc-Odor datae, heiloiyphis . Bot. wster

o 2.0 umnrity. p in delthe ding, ough thin.

the cm. oung with of a mycelioid character,  $2-8~\mu$  broad. Outer layer of veil in young buds shows a considerable number of narrow but rather thick-walled hyphae. Nuclei are seen mostly in this part of the veil on fixed and stained material. On older caps the veil consists almost exclusively of somewhat wider, more thin-walled and irregularly branched hyphae. The filaments on the stem are made up of shorter hyphae of this latter type. No clamp connections seen in any part of fruitbody, but demonstrated in mycelium (conf. below).

Habit, habitat and distribution. Single, occasionally few together, on dung of cow and goat. Texas, Galapagos Islands.

Material studied. M. Lange C. 88-type, developed in vitro in laboratory on cow dung from SE. of Santiago Peak, Brewster Co., Texas, U.S.A. May 5, to 11, 1947 (the dung collected April 7, by R. McVaugh); M. Lange C. 36, Cultures obtained from goat dung collected on the Galapagos Islands were furnished through the courtesy of Professor G. W. Martin, Univ. of Iowa, Iowa City, Iowa.

Observations. The description is drawn from specimens which occurred with other Coprini on the moistened dung, and should be considered exactly corresponding to what will be met in nature. Of the Galapagos plant I have seen only specimens raised in pure culture, on sterilized substrate. These differ slightly in average size and shape of spores:  $16-18 \times 12.5-14.5(15) \times 9.5-10~\mu$ , some of them more distinctly angular, and many of them (in some mounts up to 20 per cent) with two germpores; these spores often collapse before maturity and very likely are sterile. I deem this character much too inferior to give the Galapagos plant the rank of a distinct species or variety.

The closest relative previously described in literature seems to be C. vermiculifer Joss. (Josserand, 1944). It has the same type of veil, especially in young stages, and the same type of gills with a floccose edge occasionally splitting. Microscopically the species are easily separated on account of the very different spores and the lack of clamp-connections in the veil tissue. Macroscopically the species must be very hard to distinguish. I have looked for adequate descriptions of similar species from the tropics and subtropics, but the generally very brief diagnoses have not permitted me to identify C. myceliocephalus with any of them.

CULTURAL CHARACTERS. The species fruits very readily on horse dung agar as well as on sterilized dung, sometimes as early as 11 days after the inoculation and after the development of a very vigorous white, aerial mycelium. The spores germinate in a very




Fig. IV. Coprinus myceliocephalus.

low percentage on horse dung agar, generally only 0.01–1 per cent. Several different ceatments, namely heating of the spores in juice of dung, addition to the agar of other types and concentrations of dung juice, and of various mineral nutrient, were tried with little or no effect. The germination of spores and development of one

in young k-walled on fixed most exregularly le up of s seen in below).

vitro in ster Co., oril 7, by oat dung

wa City,

few to-

ns which hould be nature. in pure average  $\mu$ , some in some res often eem this

the rank

ms to be type of s with a species and the cally the for adeand sub-

mycelium on an agar-plate seem to stimulate the germination of the neighboring spores, and some young mycelia will generally develop in the outskirts of the older one. The first hyphae grow close to the agar surface, but in a distance of about 2-3 mm. from the germinating spore aerial hyphae develop, some of which are provided with very prominent and well developed clamp-connections, two or three at the same transversal wall (FIG. Ie). As far as I know, this phenomenon, described for instance from Coniophora, has not previously been noticed for a species of Coprinus, in spite of the thorough study of the cultural characters of this genus. It has been observed as a regular feature of all mycelia studied of this species including the single-spore mycelia, but the clampconnections seem to occur with a varying frequency from plate to They are most constantly seen on the first aerial hyphae, and never observed on hyphae growing down in the agar. single-spore mycelia were isolated (from both strains). All produced normal fruitbodies, the spores of which gave rise to a second generation. From this it is evident that the species is truly homothallic.

Temperature: Mycelia were placed in constant temperature chambers at different temperatures. As one would expect, knowing the place of the origin of the cultures, the growth was very vigorous at 35° C., slightly slower at 22° C., and at 15° C. nearly zero. Cultures kept at 18° C. will sometimes develop normally, but sometimes stop growing after a while. The lower limit for normal growth thus appears to be about 20° C. and is considerably higher than for Coprini from cooler climates.

Light: The species requires light for normal fruiting. In the dark, long stipes develop with small brownish caps, the veil becomes very much reduced, and these caps never open. They collapse after a while without ripening spores. Some cultures developed no buds at all in the dark.

Acknowledgments. My work on the Coprini, to which this paper is a first contribution, has been suggested by Dr. A. H. Smith, who most kindly placed a considerable amount of material at my disposal, and gave valuable advice. To the Director of the University of Michigan Herbarium, Dr. E. B. Mains, and also to the directors of the Carlsberg Laboratorium and the Botanical

tion of lly degrow from ch are onnec-As far

Conionus, in genus. ied of lampate to

Ten prosectruly

phae,

ature nowvery early nally, t for rably

be-

this H. al at the o to iical

Laboratorium of the University in Copenhagen, where my culture work was carried out, I owe a debt of thanks for granting me the necessary facilities. The latinizing of the diagnosis is by Tyge Christensen.

#### LITERATURE CITED

Gillet, C. C. 1878-90. Les champignons qui croissent en France. Paris, 3 vols.

Josserand, M. 1944. Étude sur quelques Coprins. Bull. Soc. Myc. Fr. 60: 5-18.

Martin, Ch. Ed. 1904-05. Contribution à la flore mycologique suisse et plus spécialement genevoise. Bull. Trans. Soc. Bot. Genève 11: 110-130.

Massee, G. 1896. A revision of the genus Coprinus. Ann. Bot. 10: 125-184.

Saccardo, P. A. 1887. Sylloge Fungorum. Vol. V. Patavii.

### **EXPLANATION OF FIGURES**

I. Coprinus myceliocephalus. A, spores in face view;  $A_1$ , spore in side view; B, two basidia; C, two paraphyses; D, pleurocystidia;  $D_1$ , short cheilocystidium; E, hyphae from mycelium with triple and double clamp-connections; F, hyphae from veil of young cap;  $F_1$ , hyphae from flocci on stem;  $A_1 \times 1000$ ;  $A_2 \times 1000$ ;  $A_3 \times 1000$ ;  $A_4 \times 1000$ ;  $A_5 \times 1000$ ;

II. Coprinus cothurnatus God. A, elements from veil on cap; B, mycelium with clamp-connection; C, two basidia; D, two paraphyses; E, two spores in face view;  $E_{10}$  spore in side view. (A-D × 500; E and  $E_{1}$  × 1000.)

### MYCOLOGICAL NOTES. IX

C. L. SHEAR

### 37. THE GENUS VENTURIA AND A PROPOSED LECTOTYPE

De Notaris in his original publication of the genus *Venturia* (Atti Sci. Ital. **6**: 484. 1844) described two species, *V. rosae* and *V. dianthi*. It might naturally be expected that subsequent authors would have based their concept of the genus on one or the other of these species, but this was not the case, the type method of fixing the application of generic names having not yet been proposed. Cesati and de Notaris (Comment. Soc. Crittogam. Ital. 1: 225. [Reprint p. 51.] 1863) transferred other species to the genus, including *V. dickei* (Berk. & Br.), *V. eres* (Berk. & Br.), *V. macrotrichia* (Berk & Br.), and *V. chaetomium* (Cda.).

Saccardo (Sylloge Fungorum 1: 586. 1882) took the genus in the sense of Cesati and de Notaris (1. c. 1863) "pro minore parte" rather than as first set up by de Notaris. He noted that the two original species were setose Pleosporas for which at a later date (Syll. Fung. 2: 285. 1883) he established the genus *Pyrenophora*. In his compilation he included species referred to the genus by Fries, Cooke, Karsten, and others as well as a number added by himself to make up a heterogeneous group of forty species.

In 1886 Berlese and Saccardo (Ann. Soc. Ven. Trent. 10: 174) proposed a new genus, *Proventuria*, based on the original *Venturia rosae* de N. (*Pyrenophora rosae* [de N.] Sacc.), a study of authentic specimens having shown that the ascospores were uniseptate and brown rather than muriform. The other original species, *V. dianthi*, was renamed *Pyrenophora notarisii* Sacc. (Syll. Fung. 2: 285. 1883) as already noted.

Since the publication of the Sylloge the general usage of mycologists and phytopathologists has been that of Saccardo. However, even a casual study of the various species included by Saccardo shows that several genera are represented, so that in order to apply the name in a definite manner and to avoid further confusion, the

name should be conserved and typified by a species representing the most general current usage. The best known group of species is that including V. inaequalis (Cke.) Wint. and V. pyrina Aderh. On account of their great economic importance as the cause of apple and pear scab they have been thoroughly investigated. Their conidial form is Fusicladium. Until we have more complete information in regard to many of the other species now included in the genus no satisfactory, complete segregation can be made.

Clements and Shear (The Genera of Fungi, p. 267. 1931) proposed as the type *V. chlorospora* (Ces.) Karst. which Ellis and Everhart (No. Amer. Pyren. p. 138. 1892) noted as a synonym of *V. inaequalis* (Cke.) Wint., although they are now regarded as distinct species. But it is perhaps better to take *V. inaequalis* as the type since it is the more familiar species. Bisby and Mason (Trans. Brit. Myc. Soc. 24: 172. 1940) state that *Venturia* should be conserved and included in their list of species, *V. aucupariae*, *V. chlorospora*, *V. inaequalis*, and *V. pirina*.

In 1923 Sydow (Ann. Myc. 21: 171) protested the action of Saccardo and Berlese, as mentioned above, in transferring the two original Venturia species of de Notaris to other genera and using the name for another group of heterogeneous species. He proposed a new genus, Spilosticta, typified by V. rumicis (Desm.) Cke. and added a new species, S. bistortae. Species of this genus were characterized by buried foliicolous perithecia and an Ovularia conidial stage. For the leaf inhabiting species with a Fusicladium conidial stage he proposed the genus Endostigme to include the following species placed in Venturia by Saccardo: E. inaequalis (Cke.) (type), E. ditricha (Fr.), E. tremulae (Aderh.), E. chlorospora (Ces.), E. fraxini (Aderh.), E. pirina (Aderh.), and E. crataegi (Aderh.). Endostigme would then be an exact synonym of Venturia as proposed for conservation and typified by Clements and Shear and Bisby and Mason.

Petrak (Ann. Myc. 38: 193. 1940) considered Sydow's two genera synonymous and made the new combination Spilosticta inaequalis (Cke.) Petr. for the apple scab fungus. Jørstad (Nyt. Mag. Nat. 84: 251-3. 1943) accepted Sydow's genera Spilosticta and Endostigme as distinct because of their different conidial stages. For the type species of the latter, however, he proposed to substi-

E nturia

thors other fixing posed. 225. s, in-

arte"
two
date
hora.
s by
d by

us in

turia henotate , V.

olover, ardo oply the tute Sphaerella cinerascens Fleischhack for E. inaequalis, making the new combination E. cinerascens (Fleischhack) Jørstad, since he considered cinerascens the oldest valid specific name for the apple scab fungus. This action does not appear to be justified for the following reasons:

Jørstad's decision was based on Rabenhorst's use of the name in 1865 for a fungus issued as No. 845 of his exsiccati, the label reading as follows:

Rabenhorst, Fungi europaei. 845. SPHAERELLA CINERASCENS Fckl. Fung. Rhenan. N. 824.

Paraphysibus nullis, ascis tubulosis octosporis, sporis uniseriatis, pulchre chlorinis, uniseptatis, 1/76 mm longis a Sphaerella ditricha Fr., et Sphaerella chlorospora Cesati vix diversa.

In foliis Sorbi ariae legit Arnstadiae (Thuringiae) Majo 1865.

Dr. Fleischhack.

There is nothing here to justify the conclusion that Fleischhack was the author of the description or the name, since Rabenhorst clearly cites Fuckel as the author of the species and Fleischhack as the collector. This error in ascribing the authorship of Sphaerella cinerascens to Fleischhack apparently originated with M. C. Cooke (Jour. Bot. 4: 248. 1866), who included it as a synonym in describing Sphaerella inaequalis as a new species common in England on dead leaves of ash, hawthorn, pear, and apple: "Sphaerella cinerascens, Fleisch. Rab. Fung. Eur. no. 845 (not S. cinerascens, Fuckel, Fung. Rhen. no. 824." This synonymy is accepted on the basis of Cooke's statement since we have been unable to find the fungus itself on the specimen of Rabenhorst, No 845, in the Mycological Collections of the Bureau of Plant Industry. It will be noted that the label on this specimen as cited above credits the binomial Sphaerella cinerascens to Fuckel on the basis of a specimen issued in the latter's Fungi Rhenani, No. 824. However, the Fuckel specimen was issued under the name Sphaeria cinerascens so that Rabenhorst's usage constitutes a new combination, although it was again cited by Fuckel in 1869 (Symb. Myc. 103) as Sphaerella cinerascens Fckl. Both Cooke (l. c.) and Fuckel (l. c.) state that Sphaerella cinerascens Fckl. in Rab. Fung. Eur. no. 845 and Sphaerella (Sphaeria) cinerascens Fckl. in Fung. Rhenani no.

824 are different fungi, but this is irrelevant to the present discussion.

After all the specific epithet cinerascens is invalid for the apple scab fungus since Sphaeria cinerascens Fckl. is a later homonym of Sphaeria cinerascens Schw. (Syn. Fung. Am. Bot. p. 225. 1832) described on leaves of Asclepias. Cooke transferred this species to Sphaerella (Jour. Bot. 21: 130. 1883) and Saccardo (Svll. Fung. 1: 31. 1883) considered it a Laestadia. With the elimination of cinerascens, inaequalis, the epithet most generally used in the extensive literature on the apple scab fungus, remains acceptable. Winter appears to have been the first to make the transfer from Sphaerella of Cooke to Venturia in 1875, or at least it is credited to him by de Thuemen on the label of his specimen No. 261, Mycotheca universalis. Aderhold (Hedwigia 36: 81-2. 1897) in his careful studies of a related group of Venturia species restricted the name V. inaequalis to the fungus on Malus and Pyrus (other than P. communis), drew up an emended description, and credited the combination to himself. A correct citation of the species would appear to be Venturia inaequalis (Cke.) Wint. emend. Aderh., as noted by Jørstad (l. c.), which for ordinary usage becomes

VENTURIA INAEQUALIS (Cke.) Wint. apud de Thuemen Myc. Univ. Exsic. no. 261. 1875.

Sphaeria cinerascens Fckl. Fungi Rhen. no. 824. 1863. (non S. cinerascens Schw. Syn. Fung. Am. Bot. 225, 1832.)

Sphaerella cinerascens Fckl. apud Rab. Fungi Eur. no. 845. 1865. (non S. cinerascens Cke. Jour. Bot. 21: 130. 1883.)

Sphaerella inaequalis Cke. Jour. Bot. 4: 248. 1866,

Venturia inaequalis Aderhold Hedwigia 36: 81-2. 1897.

Didymosphaeria inaequalis Niessl in Rabh. Fung. Eur. Exsic. no. 2663.

Endostigme inaequalis Syd. Ann. Myc. 21: 171. 1923.

Spilosticta inaequalis Petr. Ann. Myc. 38: 193. 1940.

Endostigme cinerascens Jørstad Nytt. Mag. Nat. 84: 252. 1944.

#### 38. VENTURIA ON ERICACEAE

There is a group of closely related species usually referred to Venturia, which are found on leaves of ericaceous plants. They

ame in l read-

making

, since for the

ied for

pulchre aerella

nack.

k was learly as the erella Cooke

n degland
erella
scens,
n the

d the Iycoill be

ie biimen uckel

that

rella state and

no.

differ from the type species, *V. inaequalis*, in having superficial perithecia and no known conidia. They appear to be nearer to *Coleroa* Rab. as typified by *C. chaetomium* (Kze.) Rab., which occurs on *Rubus*. Paraphyses are usually uncertain or wanting. *Gibbera*, as typified by *G. vaccinii* Fr., has *Helminthosporium* conidia and paraphyses and does not seem congeneric. The species to be considered are three.

Venturia arctostaphyli Cke. & Hark. Grev. 13: 20. 1884.

This species was originally described from dead leaves of *Arctostaphylos* collected in California. An examination of part of the type material collected by Harkness on leaves of *A. pumila* shows small superficial perithecia scattered over spots on dead leaves. The surface of the spots has the grayish-white, somewhat glistening appearance common to this group of species. They lose something of this characteristic after infected leaves have fallen and become faded. We find mature spores  $15-18\times 5~\mu$ , somewhat larger than the measurements recorded in the description  $(12-15\times 5~\mu)$ . Setae around the summit of the perithecia are from  $50-75~\mu$  long. This does not seem to be specifically distinct from the fungus found on *A. uva-ursi* in Massachusetts, although in the eastern form, the setae on the perithecia are only  $40-45~\mu$  long and not so numerous. This slight difference, however, could scarcely be considered specific.

Venturia cassandrae Pk. Report N. Y. State Bot. 38: 104. 1885.

Peck noted that this species caused reddish-brown or brownish spots, sometimes with a grayish center and that the perithecia were minute, broad, black, with a few short, straight, diverging black setae above. He found the fungus on living leaves of Cassandra calyculata in New York State. He adds further: "the perithecia sometimes occur on the upper surface of the leaf, but oftener on the lower. They are so small that they are scarcely visible to the naked eye. Sometimes they emerge from beneath the scales of the leaf, and then they appear erumpent, although in reality they are superficial."

Venturia gaultheriae Ell. & Ev. Jour. Myc. 1: 153. 1885.

eriroa

on

ra.

ind

on-

to-

the

WS

res.

ing

ing

me

nan

 $\mu$ ).

ng.

ınd

the

us.

red

85.

ish

ere ack

dra

cia

on

the

the

are

The description of this species as taken from Ellis and Everhart's North American Pyrenomycetes follows:

"On orbicular, dark brown,  $\frac{1}{3}$  mm. spots, which are mostly of a lighter color (gray) in the center. Perithecia scattered, orbicular (75  $\mu$ ), membranous and rather coarsely cellular, with a few black, continuous, straight, spreading,  $35 \times 3 \mu$  bristles above. Asci ovate-oblong,  $30\text{--}35 \times 8\text{--}11 \mu$ , broader and slightly curved below, sessile, without paraphyses. Sporidia biseriate, subhyaline (with a greenish-yellow tint), ovate-oblong, 3–4-nucleate, uniseptate and slightly constricted at the septum,  $11\text{--}14 \times 3 \mu$ .

"On living leaves of Gaultheria procumbens, Newfield, N. J."

From examination and comparison of type material and other authentic specimens we conclude that the three species described above are the same for which the oldest name V. arctostaphyli Cke. & Hark. may be used. We have found this species also on  $Vaccinium\ macrocarpon$ , the cranberry. Other species closely related and apparently congeneric are V. pulchella Cke. & Pk. and V. kalmiae Pk., differing from V. arctostaphyli in the size of the ascospores, which are  $7-10\times 3\,\mu$ . Further study of these and other species such as V. dickei (Berk. & Br.) Ces. & de N. and V. myrtilli Cke. must be made in order to determine whether they deserve generic segregation.

# 39. VENTURIA CINCINNATA (Fr.) Fr.

Fries (Syst. Myc. 2: 451. 1823) described *Sphaeria cincinnata*, occurring very rarely on leaves of *Vaccinium oxycoccum*. He later (Summa Veg. Scand. p. 405. 1849) transferred the species to *Venturia*.

Schweinitz (Syn. Fung. Am. Bor. p. 22. 1832) reported the species as occurring on leaves of *Oxycoccus macrocarpus* at Pocono, Pennsylvania. An examination of his specimen, however, indicates that it is not Fries' species, but what was later named by Peck (Rept. N. Y. State Bot. 25: 106. 1873) as *Venturia compacta* and transferred erroneously by the writer (U. S. Dept. Agric. Tech. Bull. 258: 13. 1931) to *Gibbera* which has a conidial stage. The above species has none.

Just what species Fries actually had is uncertain. We were unable to find any specimen under this name in his herbarium. Later reports of the species appear to be limited to three: Ellis and Everhart (North Amer. Pyren. p. 142. 1892) record it from Greenland on Vaccinium palustre; Rostrup (Bot. Tids. 27: 35 F. 1906) also reports it from Greenland and on the same host; and there is a specimen in the Mycological Collections of the Bureau of Plant Industry from Bresadola. This is labelled Venturia cincinnata (Fr.) Rost. in his own handwriting and with no other data than the statement "Pyrenomycète sur les feuilles d' Oxycoccos palustris avec un croquis." The specimen is presumably of European origin. It consists of a number of dead leaves bearing on their lower surfaces perithecia typical of Acanthorhynchus vaccinii Shear. The sketch referred to on the label shows the typical short beak of this species with the stiff setae as well as the asci, ascospores and paraphyses characteristic of the species. Measurements given for the ascospores are  $30-40 \times 12-15 \,\mu$ . This is clearly not the fungus described by Fries who states that the perithecia of his fungus are entirely superficial, small, subparabolic and sometimes surrounded by curly hairs at the base. In the Bresadola specimen the perithecia are large, subglobose, and completely embedded in the leaf tissue.

No specimens have been seen on Oxycoccus or Vaccinium which agree with Fries' description. His fungus may be related to the Venturia arctostaphyli group, although we have never seen any specimens of this group with perithecia having curly basal hairs nor with conical-cylindrical perithecia. It may really be "sui generis" as Fries says.

# 40. Notes on Antennaria, Antennularia, and Niesslia

Antennaria as described by Link (In Schrader, Neu. Jour. Bot. 3: 16, 1809, not Gaert., 1791) was typified by A. ericophila. Link described and illustrated only the conidial form of the fungus. Nearly a century later Neger (Centralbl. Bakt. Par. 20 (2): 94. 1907) collected Link's fungus, which originally came from Portugal, in Andalusia on Erica arborea. He found perithecia as well as the typical Antennaria form of conidiophores described

and illustrated by Link. At an altitude of 200 to 400 meters there was but small growth of subiculum on the smallest branches of the host plant, and especially on the leaf axils and nerves of fallen leaves. This subiculum was small (about the size of a pin-head) and bore perithecia particularly in the leaf axils. The asci were 35 F.  $40-50 \times 15-20 \,\mu$ , united in a mass which is typical of some Venturia species; the ascospores were 2-celled, greenish, and 18-20 × 7 μ. At high altitudes, the mycelium of the fungus developed around the axes of the whole plant in dense black mycelial balls, in some instances reaching the size of one's fist. These were easily reoccos moved. They absorbed water like a sponge. At about 700 meters these masses developed a peculiar growth of erect Antennaria conidiophores 1-2 mm. high, bearing a head of 4-celled dark conidia. The column (saule) of conidiophores was composed of necklacelike hyphae and the interior hyphae bore club-shaped brown conascoidia, 40-50 µ long. No perithecia were found in these mycelial balls. This, he thinks, was probably due to the greater humidity and rainfall, as well as the colder climate at this altitude. The fungus was not parasitic, he says, but its exclusion of the light and air from the host frequently weakened and killed the plants.

> Von Hoehnel (Frag. Myk. no. 379. 1909) discusses Antennaria Link and states that nothing certain was known about its identity and relationships until Neger described it as discussed above. After an examination of Neger's specimens von Hoehnel said that he was convinced Antennaria ericophila Lk. was identical with Coleroa straussii (Sacc. & Roum.) Hoehn. (Venturia straussii Sacc. & Roum.), but that no Antennaria form of C. straussii was known. Further investigation, he said, showed that when mature the perithecia had a distinct ostiole and that conidia were found.

> Venturia straussii Sacc. & Roum. (Rev. Myc. 6: 95. 1884) was based on material collected in 1884 by M. Merlet on Erica scoparia near Bordeaux, France, and sent by him to Roumeguère. Saccardo, who received the specimen for study, considered it identical with the fungus described and illustrated by Strauss (in Sturm, Deutschland Flora, Abt. III, Heft 34: 29-30, Tab. 3, Figs. A-G. 1853) under the name Chaetomium pusillum Fr. Strauss' identification, however, was in error, as was indicated by Saccardo's action in setting up a new name for the Merlet specimen.

were rium. s and from

; and au of cindata

Lurog on ccinii short

nents y not of his times

imen ed in

hich the any nor eris"

IA Bot. hila. gus.

94. Pora as ibed

Ho

Ex

31

op

M

spe

lea

to

an

en

on H

K

an

yr

A

F

A

The original Friesian fungus was referred to *Niesslia* by Schroeter and also named de novo by Albertini and Schweinitz and Corda as the following partial synonymy will show. This species occurring on pine needles should not be confused with the two species on Ericaceous hosts discussed hereafter.

Niesslia pusilla (Fr.) Schroet., in Cohn, Krypt. Fl. Schl. 3: 294. 1893.

Sphaeria exilis Alb. & Schw. Conspect. Fung. 44. 1805. Chaetomium pusillum Fr. Syst. Myc. 3: 255. 1829.

Sphaeria chaetomium Corda Icones II, 29. 1838.

Venturia chaetomium Ces. & de N. Schema Class. Sfer. Ital. 225. 1863.
Niesslia chaetomium Auers. in Gonnerman Rab. Mycol. Eur. Heft V & VI 30. 1869.

Niesslia exilis Wint. in Rab. Krypt.-Fl. 2 Aufl., 1 Bd., II Abt. 196. 1885.

The fungus described and illustrated by Strauss (1. c.) was found on living leaves of *Erica carnea* near Munich. No type material is known. There is little doubt, however, that *Venturia straussii* Sacc. and Roum. is the same and that as von Hoehnel states it is the same as *Antennaria ericophila* (Lk.) Neger.

The name Antennaria Link being untenable because of an earlier use of the name for a phanerogamic plant by Gaertner (1791), Reichenbach in 1828 (Consp. Reg. Veg. p. 5) proposed Antennularia as a substitute. This was also invalid for an ascogenous fungus until the perithecial form was found and described by Neger and adopted by von Hoehnel as discussed above. We regard the genus Antennularia (Reich.) Hoehn. as having been validated by von Hoehnel (Frag. Myk. 379. 1909) when he described Neger's specimens of Antennaria ericophila Lk., which showed conidia, perithecia, and ascospores, and adopted Reichenbach's name for the fungus.

He added three other species to the genus in addition to the type, A. ericophila (Lk.) Hoehn., among them being A. salisburgensis (Niessl) Hoehn. The original material upon which the description of this latter species was based was distributed in 1886 as no. 3550 in Rabenhorst—Winter's Fungi Europaei Exsiccati. After comparing specimens of this number with the original material of Venturia straussii Sacc. & Roum. (Antennularia ericophila (Lk.)

a by

and

ecies

two

294.

63.

& VI

1885.

und

al is

issii it is

lier

91),

ten-

ous

ger the

by

er's

erithe

pe,

sis

ip-

no. ter

of

(.)

Hoehn.) issued by Roumeguère as no. 2828 of his Fungi Gallici Exsiccati and other specimens of the same collection issued as no. 3142 of Rabenhorst—Winter's Fungi Europaei, we are of the opinion that the two are different species. Von Hoehnel (Frag. Myk. no. 115. 1907) came to the same conclusion after studying specimens of the same numbers. The chief difference is that in *A. ericophila* the perithecia form dense masses in the axils of the leaves and on the stems whereas in the second species they appear to be confined to the leaves. The ascospores in the first are shorter and broader  $(15-18 \times 6-8 \mu)$  than in the second  $(18-21 \times 5-6 \mu)$ .

Von Hoehnel has added two other species to this genus, A. engleriana (P. Henn.) Hoehn. (Frag. Myk. no. 356. 1909) based on Dimerosporiopsis englerianus P. Henn. and A. rhododendri Hoehn. based on the ascogenous form of Torula rhododendri Kunze. We have seen no authentic specimens of either of these and therefore can express no opinion regarding them. The synonymy of the two species which we have studied is as follows:

Antennularia ericophila (Neger) Hoehn. Sitzungsb. Akad. Wiss. Wien. 118, abt. 1: 1198. 1909.

Antennaria ericophila Lk. in Schrader Neues Jour. Bot. 3: 16. 1809.

Venturia straussii Sacc. & Roum. Rev. Myc. 6: 95. 1884.

Antennaria ericophila Neger, Centralbl. Bakt. 20 (2): 94. 1897.

Coleroa straussii Hoehn. Sitzungsb. Akad. Wiss. Wien 116, Abt. 1: 115. 1907.

Exsiccati specimens examined: Roumeguère, Fungi Gallici exs., no. 2828, as Venturia straussii Sacc. & Roum.; Rabenhorst—Winter Fungi Europaei, no. 3142, as V. Straussii Sacc. & Roum.

Antennularia salisburgensis (Niesel) Hoehn. Oesterr. Bot. Zeit. 63: 233. 1913.

Chaetomium pusillum Strauss in Sturm, Deutschland Flora, Abt. III, Heft 34: 29-30. 1853, non Fr. 1829.

Gibbera salisburgensis Niessl Hedw. 26: 33. 1887.

Eriosphaeria salisburgensis Neger Ber. Deutsch. Bot. Gesell. 19: 471. 1901.

Coleroa salisburgensis Hoehn. Sitzungsb. Akad. Wiss. Wien 116, Abt. 1: 115. 1907.

Gibbera straussii Zahlb. Krypt. Exs. Mus. Pal. Vind., no. 824.

Exsiccati specimens examined: Jaap, Fungi Sel. Exs., no. 614, as Antennaria salisburgensis (Niessl) Hoehn.; Zahlbruckner, Krypt. Exs. Mus. Pal. Vind., no. 824, as Gibbera straussii Zahlb.; Rabenhorst-Winter, Fungi Europ. Exs., no. 3550, as Gibbera salisburgensis (Niessl) Hoehn.; Rehm, Ascom. Exs., no. 1939, as Coleroa salisburgensis (Niessl) Hoehn.

The fungus issued by Jaap (Fungi Sel. Exsic. no. 657) as Antennaria salisburgensis (Niessl) Hoehn, associated with the aphid, Eriococcus ericae, on Erica tetralix is doubtfully the same as his no. 614. It has perithecia with the short, stiff setae and the colored ascospores of the same size and shape as the other specimen, but the perithecia occur on the stems as well as on the leaves and show more of the long basal hyphae, which seem to have grown over some of the aphids. It will be necessary to learn more in regard to the relation between the aphids and this and related species of these fungi, especially Antennularia ericophila, before a final decision is possible.

A further specimen found in the Mycological Collections of the Bureau of Plant Industry should be considered here. It was issued by C. Torrend as no. 146 of his Fungi Selecti Exsiccati and is labelled "Gibbera salisburgensis Niessl, ad folia Ericae arboreae, Madère, C. de Menzes, VI–1912." It consists of a few leaves bearing on the under side small dense groups of perithecia without subiculum, appendages, or setae. The perithecia are slightly rough or minutely verrucose. The perithecial wall is much thicker than in the other species mentioned, but paraphyses, asci, and ascospores are typical. The spores are  $15-20\times 6-8\,\mu$ . This is not Niessl's species. It would seem to indicate that there may be another group of species without subiculum or setae, but otherwise alike.

Neger (Ber. Deutsch. Bot. Gesell. 19: 471. 1901) after examining the Rabenhorst-Winter specimen (no. 3550) of Gibbera salisburgensis collected the fungus himself on Erica carnea. His material agreed exactly with the other. He described it in detail and showed a leaf section which indicated that a dense mass of vegetative hyphae developed beneath the thin epidermis of the upper leaf surface from which the superficial subiculum and perithecia arose. Infection took place in the summer and perithecia formed the following year. He maintained that the fungus was not a

Gibbera and transferred it to Eriosphaeria, as E. salisburgensis (Niessl) Neger.

614.

kner,

hlb.;

bbera 19, as

An-

phid.

s his lored

show over gard es of

the sued s lareae, earsubih or n in ores ssl's ther like. ambera His etail egeper ecia med t a

There has just come to our attention Petrak's paper "Über Gibbera Fr. und verwandte Gattungen" (Sydowia 1: 169–201. 1947) in which he revises the genus Gibbera with G. vaccinii Fr. as the type and includes Antennularia as a subgenus. He does not include the production of conidia as a generic character. We regard this as a character of sufficient importance to exclude such species as Venturia compacta and other superficial species having no conidia.

# MYCOLOGICAL SOCIETY MEMBERSHIP LIST \*

E

- AHMAD, SULTAN, Botany Dept., Government College, Lahore, Pakistan, India.
- AINSWORTH, G. C., London School of Hygiene and Tropical Medicine, Keppel St., London W.C., England.
- AJELLO, LIBERO, Communicable Disease Center, U. S. Public Health Service, Atlanta, Georgia.
- ALEXOPOULOS, CONST. J., Botany Dept., Michigan State College, East Lansing, Michigan.
- ALVAREZ, CANUTO C., Facultad Nal. de Agronomia, Medellin, Columbia, South America.
- AMES, LAWRENCE M., Materials Branch, The Engineer Board, Fort Belvoir, Virginia.
- AMES, RALPH W., 807 West Nevada, Urbana, Illinois.
- Anderson, (A.) Kermit W., Washington High School, Litchfield, Minnesota.
- Andrus, C. F., Div. of Fruit & Vegetable Crops, U. S. Dept. of Agric., Box 844, Charleston, S. C.
- ANTIKAJIAN, GRACE, 2741 Sedwick Avenue, New York 63, New York.
- APPEL, BERNARD, 281 Ocean St., Lynn, Massachusetts. ARDE, WALKER R., 4324 Spruce St., Philadelphia 4, Pa.
- ATKINSON, ROBERT G., Dominion Laboratory of Plant Path., Box 87, St. Catharines, Ontario, Canada.
- ATWELL, ERNEST A., Forest Products Lab. Canada, Isabella & Metcalfe Sts., Ottawa, Ontario, Canada.
- AYERS, T. T., Horticultural Research, Agricultural Expt. Station, Louisiana State University, Baton Rouge, La.
- BABCOCK, S. H., JR., Lederle Laboratories, American Cyanamide Co., Pearl River, New York.
- BACHE-WIIG, SARA, 36 Prospect Avenue, Northampton, Massachusetts.
- BACKUS, EDWARD J., Lederle Laboratories, Pearl River, New York.
- BACKUS, MYRON P., Biology Bldg., University of Wisconsin, Madison, Wisconsin.
- BAGDON, VINCENT J., 703 Timber Branch Drive, Alexandria, Virginia.
- BAKER, GLADYS E., Dept. of Plant Sciences, Vassar College, Poughkeepsie, New York.
- BAKER, KENNETH, Div. Plant Path., U. of Calif. L. A., Los Angeles, California.
- BARGHOORN, ELSO S., Jr., Biological Laboratories, 16 Divinity Avenue, Cambridge 38, Mass.
- \*A double asterisk indicates life membership; a single asterisk, associate membership.

- BARNETT, HORACE L., Botany Dept., West Virginia University, Morgantown, West Virginia.
- BARRETT, J. T., 107 Hilgard Hall, University of California, Berkeley, California.
- BARRUS, M. F., N. Y. State College of Agriculture, Cornell University, Ithaca, New York.
- BARTSCH, ALFRED F., State Board of Health, State Office Building, Madison, Wisconsin.
- Batista, Augusto C., Instituto de Pesquisas Agronomicas, Dois Irmaos, Recife, Brazil, S. A.
- BAXTER, Dow V., School of Forestry & Conservation, University of Michigan, Ann Arbor, Michigan.
- BECHTEL, A. R., Dept. of Botany, Wabash College, Crawfordsville, Indiana.
- Benedek, Tibor, 1706 E. 56th St., Chicago 37, Illinois.

  Beneke, Everett S., Dept. Botany and Plant Path., Michigan State College,
  East Lansing, Michigan.
- Benham, Rhoda Williams, College of Physicians & Surgeons, Columbia University, 630 W. 168th St., New York City.
- BENJAMIN, RICH. K., Rm. 308, Nat. History Bldg., University of Illinois, Urbana, Illinois.
- BENNETT, RALPH E., 724 S. 25th St., Terre Haute, Indiana.

P

kistan,

, Kep-

Serv-

Lans-

ımbia,

elvoir,

finne-

gric.,

7, St.

Sts.,

siana

Pearl

lison,

epsie,

Cali-

enue,

ciate

S.

- Bergman, H. F., Bureau of Plant Industry, East Wareham, Massachusetts. Berry, Edward C., 1008 Marshall St., Richland, Washington.
- Bessey, Ernst A., Botany Dept., Michigan State College, East Lansing, Michigan.
- BINGHAM, R. T., 618 Realty Bldg., Spokane 8, Washington.
- BISHOP, HARLOW, Cottage Avenue, Nanuet, New York.
- BITANCOURT, AGESILAU A., Instituto Biologico, Caixa Postal 119A, São Paulo, Brazil, South America.
- BLACKWELL, ELIZABETH, Royal Holloway College (Univ. of London), Englefield Green, Surrey, England.
- BLISS, DONALD E., University of California, Citrus Expt. Station, Riverside, California.
- BODMAN, SISTER MARY CECILIA, 116 E. Jefferson St., Iowa City, Iowa.
- Bonar, Lee, Dept. of Botany, University of California, Berkeley, California. Bonner, John T., Dept. of Biology, Princeton University, Princeton, New Jersey.
- BOONE, DONALD M., 1116 Mount St., Madison 5, Wisconsin.
- BOOTHROYD, CARL W., Dept. Plant Path., Cornell University, Ithaca, New York.
- Botts, William, 315 Summit Lane, Fort Mitchell, Kentucky.
- BOURNE, B. A., United States Sugar Corp., Clewiston, Florida.
- BOUTHILET, ROBERT J., 3616 46th Avenue S., Minneapolis 6, Minnesota.
- Boyce, J. S., Osborn Botanical Labs., Yale University, New Haven, Con-
- Brasfield, T. W., 521 Lexington St., Homden, Connecticut.
- Braun, Alvin J., N. Y. State Agric. Station, Drawer 353, Geneva, New York.
- Brian, Percy W., Imperial Chem. Industries, Ltd., Butterwick Res. Labs., The Frythe, Welwyn, Herts, England.

BRODIE, HAROLD J., Dept. Botany, Indiana Univ., Bloomington, Indiana.

Brooks, Travis E., Botany Dept., Kansas State College, Manhattan, Kansas. Brown, Archibald W., Dominion Lab. Plant Pathology, University of Manitoba, Fort Garry, Winnipeg, Manitoba, Canada.

Brown, J. G., Dept. of Plant Pathology, University of Arizona, Agric. Experiment Station, Tucson, Arizona.

Brunel, Jules, Institut Botanique de L'Universite, 4101 est. rue Sherbrooke, Montreal, Canada.

BUDDIN, WALTER, 1 Talfourd Avenue, Reading, England.

BURK, RICHARD J., JR., 130 Mary St., Lawrenceburg, Indiana.

BURLINGHAM, GERTRUDE S., 818 Antoinette Ave., Winter Park, Florida.

BUTLER, EDWARD E., 412 Abbot Apts., East Lansing, Michigan.

CAIN, ROY F., Dept. of Botany, University of Toronto, Toronto 5, Canada. CALDWELL, GWENDOLYN BURTON, 662 Milledge Ave., Athens, Georgia.

CAMPBELL, T. H., Rm. 102 Botany and Zoology Bldg., Ohio State University, Columbus 10, Ohio.

CAMPBELL, W. A., School of Forestry, University of Georgia, Athens, Georgia.

CARRION, A. L., Dept. of Mycology, School of Tropical Medicine, San Juan, Puerto Rico.

CARVAJAL, FERNANDO, 303 Diehl Drive, Lawrenceburg, Indiana.

Cash, E. K., Bureau of Plant Industry, U. S. Dept. of Agriculture, Beltsville, Maryland.

CHARLES, VERA K., 2039 N. H Avenue, Washington, D. C.

Chen, S. M., c/o K. W. Wegner, 105 Winona St., Northfield, Minnesota. Chilton, St. John P., Dept. of Botany, Bact and Plant Pathology, Louisiana State University, Baton Rouge, La.

Chiu, Wei-Fan, Plant Pathology Dept., College of Agriculture, National Tsing Hua University, Peiping, China.

CHRISTENBERRY, G. A., Furman University, Greenville, South Carolina.

CHRISTENSEN, CLYDE, University Farm, St. Paul, Minnesota. CHRISTENSEN, J. J., University Farm, St. Paul, Minnesota.

CHRISTISON, ISABEL B., American Type Culture College, Georgetown School of Medicine, Washington 7, D. C.

Chupp, Charles D., Dept. of Plant Path., Cornell University, Ithaca, New York.

CICCARONE, ANTONIO, Departmento de Fitopathologia, Apartado 669, Calle Rivas 74, Maracay, Venezuela, S. A.

CLARK, ELIZABETH S., Dept. Plant Pathology, N. J. Agric. Experiment Station, New Brunswick, New Jersey.

CLARK, LILY H., Plant Path. Division, University of California, Los Angeles 24, California.

CLARK, MARIE B., Dept. of Botany, Howard University, Washington, D. C. CLUM, FLOYD M., 527 Watson St., Ripon, Wisconsin.

COHEN, CYRIL, Botany Department, Iowa State College, Ames, Iowa.

Сокев, W. C., Dept. of Botany, University of North Carolina, Chapel Hill, North Carolina.

COLLINS, DONALD C., 7046 Hollywood Blvd., Hollywood 28, California. Conant, Geo. H., Triarch Botanical Products, Ripon, Wisconsin.

- CONANT, NORMAN F., Duke Hospital, Duke Medical School, Durham, North Carolina.
- CONNERS, I. L., Central Experimental Farm, Ottawa, Canada.
- COOK, HAROLD THURSTON, Virginia Truck Exp. Station, Norfolk, Virginia.
- COOK, MELVILLE T., Botany Dept., L. S. U., Baton Rouge, Louisiana. COOK, W. R. IVIMEY, University College, South Wales and Monmouthshire,
  - Newport Rd., Cardiff, Wales.
- COOKE, WM. BRIDGE, 1718 A Street, Pullman, Washington.
- CORNER, E. J. H., Unesco Office, c/o Museum Nacional, Rio de Janeiro, Brazil, South America.
- COTNER, FRANK B., Botany Dept., Montana State College, Bozeman, Mon-
- COTTER, SHIRLEY L., 20 Langford Park, St. Paul 8, Minnesota.
- COUCH, JOHN N., Dept. of Botany, University of North Carolina, Chapel Hill. North Carolina.
- COVERT, Scott, Southampton, Pennsylvania.

ansas.

Agric.

Sher-

la.

ınada.

niver-

thens,

Juan,

Belts-

esota.

ouisi-

tional

chool

New

Calle

iment

An-

D. C.

Hill,

- COWAN, R. S., Dept. of Botany, University of Hawaii, P.O. Box 18, Honolulu, T. H.
- CRANDALL, BOWEN S., Centro Nacional de Agronomia, c/o American Embassy, San Salvador, El Salvador, C. A.
- CUMMINS, GEORGE B., Botany Dept., Agricultural Experiment Station, Lafayette, Indiana.
- CUNNINGHAM, H. S., Long Island Vegetable Res. Farm, Riverhead, New York.
- CURRY, D. P., Route 3, Columbia, South Carolina.
- CUTTER, V. M., JR., Dept. of Botany, Yale University, New Haven, Conn. Damon, Samuel C., II, Dept. Botany, Brown University, Providence,
- Rhode Island.
- DARBY, RICHARD T., 5029 Hazel Avenue, Philadelphia 43, Pa.
- DARKER, GRANT D., Ben Venue Lab., Inc., Bedford, Ohio.
- DARLEY, ELLIS F., c/o American Embassy, Guatemala City, Guatemala, Central America.
- DAVIDSON, Ross W., Plant Industry Station, Beltsville, Maryland.
- DAVIS, BENJAMIN HAROLD, N. J. Agric. Expt. Station, New Brunswick, New Jersey.
  - DEARDEN, ELIZABETH R., Dept. of Botany, University of Toronto, Toronto, Ontario, Canada.
- DEARNESS, JOHN, University of Western Ontario, 30 Marley Place, London, Ontario, Canada.
- DE L'AMATER, EDWARD D., Lab. of Mycology, Mayo Clinic, Rochester, Minnesota.
- DENNISON, MRS. RAYMOND A., 1166 W. Cypress St., Gainesville, Florida.
- Dick, Esther A., Botany Dept., Brown University, Providence, Rhode Island.
- DIEHL, W. W., U. S. Dept. of Agriculture, B.P.I., Washington, D. C.
- DIGILIO, ANTONIO PEDRO LUIS, Seccion Criptogamia, Instituto Miguel Lillo, Calle Miguel Lillo 205, Tucuman, Argentina, S. A.
- Dodge, B. O., New York Botanical Garden, Bronx Park, New York.

Donge, C. W., Missouri Botanical Garden, 2315 Tower Grove Ave., St. Louis, Missouri.

DOSDALL, LOUISE, University Farm, St. Paul, Minnesota.

DOUBLES, JAMES A., JR., 1009 Courtland St., Greensboro, North Carolina. DOWNING, MARY H., Biological Labs., Quartermaster Depot, 2800 S. 20th St., Philadelphia 45, Pa.

DRAKE, CHARLES H., Dept. of Bacteriology, Washington State College, Pullman. Washington.

DRAYTON, F. L., Div. of Botany, Central Experimental Farm, Ottawa, Canada.

DRECHSLER, CHARLES, B.P.I., U.S.D.A., Washington, D. C.

DRIVER, CHARLES H., Fulton County Botanical Garden, Route 8, Atlanta, Ga. Du Charme, E. P., Citrus Expt. Station, Lake Alfred, Florida.

DUDDINGTON, C. L., The Polytechnic Dept. of Biology, 309 Regent St., London W. 1, England.

DULANEY, E. L., Microbiological Research, Merck & Co., Rahway, New Jersey.

EDGERTON, C. W., Dept. of Botany, Louisiana State University, Baton Rouge 3, La.

EHRLICH, HOWARD G., Div. Plant Path., Univ. Minn. Univ. Farm, St. Paul. Minnesota.

EHRLICH, JOHN, Research Lab., Parke, Davis and Co., Detroit 32, Michigan. Elliot, E. W., c/o A. Jackson, P.O. Box 121, Buffalo, Montana.

Ellis, Don E., Plant Path. Section, Box 5397, North Carolina State College, Raleigh, North Carolina.

ELLISON, BERNARD, Dept. of Botany, Univ. of Michigan, Ann Arbor, Mich. Emerson, Ralph, Dept. of Botany, University of California, Berkeley, California.

EMMONS, CHESTER W., National Institute of Health, Washington, D. C. ESPENSHADE, MARLIN A., George Washington University, School of Medicine, 1335 H St., N.W., Washington, D. C.

EZEKIEL, WALTER N., Bureau of Ordnance, Navy Department, Washington, D. C.

FAULL, J. H., 72 Fresh Pond Ave., Cambridge 38, Massachusetts.

FAWCETT, H. S., Citrus Expt. Station, University of California, Riverside, California.

FAY, DOLORES J., 115 S. 11th St., Newark 7, New Jersey.

Fennell, Dorothy J., Northern Regional Research Lab., 825 N. University, Peoria, Illinois.

Fergus, Charles L., Botany Dept., Pennsylvania State College, State College, Pa.

FISHER, FRANCINE E., Citrus Expt. Station, Lake Alfred, Florida.

FISHER, GEORGE W., Agricultural Experiment Station, Pullman, Washington. FITZPATRICK, HARRY M., Dept. of Plant Pathology, Cornell University, Ithaca, New York.

FORD, AUSTIN J., 310 Lane Street, Jessup, Pennsylvania.

FOWLER, MARVIN E., Div. of Forest Pathology, Department of Agriculture, Bureau of Plant Industry, Washington, D. C.

FRIEDMAN, BERNARD A., Rm. 1022, 641 Washington St., New York 14, New York. Fritz, Clara W., Forest Products Labs. of Canada, Metcalfe & Isabella Sts., Ottawa, Canada.

FULMER, W. L., 505 Boylston North, Seattle, Washington.

ve., St.

olina.

S. 20th

College,

)ttawa,

ta, Ga.

nt St.,

New

Baton

Paul.

higan.

· Col-

Mich.

keley,

C.

Medi-

igton,

rside,

rsity,

Col-

gton.

rsity,

ture.

New

GAINES, J. G., Coastal Plain Expt. Station, Tifton, Georgia.

GALLAGHER, ROBERT S., 345 East Main St., Moorestown, N. J.

Gammel, John A., Dept. Dermatology, School of Medicine, Western Reserve Univ., Cleveland, Ohio.

GARCES, CARLOS O., Palace 58-59, Medellin, Columbia, S. A.

GARDNER, MAX W., University of California, 107 Hilgard Hall, Berkeley, California.

GARDNER, PHYLLIS, Dept. of Botany, University of California, Berkeley 4, California.

GAUVEY, MARY P., 705 Grand Avenue, Dayton, Ohio.

GEORG, LUCILLE K., 72 W. 109th St., New York 25, New York.

GILBERT, E. M., Biology Bldg., University of Wisconsin, Madison, Wisconsin.

GILBERT, E. J., 4 Rue de Musset, Paris 16, France.

GILBERT, FRANK A., 2262 Yorkshire Road, Columbus 8, Ohio.

GILKEY, HELEN M., Herbarium, Oregon State College, Corvallis, Oregon.

GILMAN, JOSEPH C., Botany Dept., Iowa State College, Ames, Iowa. GOHEEN, AUSTIN C., (A) Dept. Plant Pathology, State College of Washington, Pullman, Washington.

GOLDSWORTHY, MARION C., Div. of Fruit and Veg. Crops, Plant Ind. Station, Beltsville, Maryland.

GOODDING, LESLIE N., Box 303, Jerome, Arizona.

GORDON, FRANK M., Farm Bureau Office, Old Court House Annex, Mineola, New York.

GORDON, W. L., Dominion Rust Research Lab., Winnipeg, Manitoba, Canada. GOSSELIN, ROGER, 3 Belleau Avenue, Levis, P. Q., Canada.

Goto, Shosuke, Dept. Plant Path., Univ. Farm, St. Paul, Minnesota. Gravatt, Annie R., Bureau of Plant Industry, Beltsville, Maryland.

GRAVATT, ANNIE R., Bureau of Flant Industry, Bensvine, Marylan Graves, Arthur H., 255 S. Main Street, Wallingford, Connecticut.

GRAY, WILLIAM D., Ohio State University, Columbus, Ohio.

GRAY, WILLIAM D., Onto State University, Columbus, Onto.

GREATHOUSE, GLENN A., National Academy of Science Bldg., 2101 Constitution Ave., Washington, D. C.

GREENE, H. C., Dept. of Botany, Biology Bldg., University of Wisconsin, Madison, Wisconsin.

GROVES, J. W., Division of Botany, Central Experimental Farm, Ottawa, Canada.

GRUBER, WM. B., 6476 S.W. Burlingame Place, Portland, Oregon.

GRUNBERG, EMANUEL, 2517 West Burnett Avenue, Louisville, Kentucky.

GUISCAFRÉ-ARRILLAGA, J., Dept. of Botany, Louisiana State University, Baton Rouge, La.

HAENSELER, C. M., Agricultural Expt. Station, New Brunswick, New Jersey. HAHN, GLENN G., Marsh Hall, Yale University, 360 Prospect St., New Haven, Connecticut.

HANNA, W. F., Dominion Rust Research Lab., Winnipeg, Manitoba, Canada. HANSBROUGH, JOHN R., Marsh Hall, Yale University, 360 Prospect St., New Haven, Connecticut.

HANSEN, H. N., 211 Hilgard Hall, University of California, Berkeley 4, California. HANSON, ANNE M., 175 West 76th St., New York 23, New York.

HARDING, PAUL R., JR., Bowling Green State Univ., Bowling Green, Ohio.

HARDISON, J. R., Agricultural Expt. Station, Corvallis, Oregon.

HARRAR, J. G., Rockefeller Foundation, Viena 26, Mexico City, D. F. Mexico.

HARRISON, K. A., Lab. of Plant Pathology, Kentville, Nova Scotia, Canada.
HARVEY, J. V., Biological Lab., Q. M. Depot, 2800 S. 20th St., Philadelphia 45, Pa.

HASKINS, REGINALD H., Prairie Regional Lab., Natl. Res. Council, Saskatoon, Saskatchewan, Canada.

HATCH, WINSLOW R., Dept. of Botany, State College of Washington, Pullman, Washington.

HAZEN, ELIZABETH L., 339 E. 25th St., New York, N. Y.

HENRY, L. K., Asst. Curator of Botany, Carnegie Museum, Pittsburgh, Pennsylvania.

HESLER, L. R., Dept. of Botany, University of Tennessee, Knoxville, Tenn. HESSELTINE, C. W., Lederle Labs., Inc., Pearl River, New York.

HILBORN, M. T., Agricultural Expt. Station, Orono, Maine.

HILLEGAS, ARTHUR B., Research Labs., Parke Davis & Co., Detroit 32, Michigan.

HIRSCHHORN, ELISA, Estacion Experimental Central, Castelar, F.C.O., Argentina, S. A.

HONEY, EDWIN E., Dept. Plant Path., Pennsylvania State College, State College, Pennsylvania.

Hopkins, J. G., Dept. of Medical Mycology, 630 W. 168th St., New York, New York.

HOPPE, MARDEL E., R. R. 2, Box 281, Waukesha, Wisconsin.

HORNER, RUTH M., Div. of Botany and Plant Path., Central Experimental Farm, Ottawa, Ontario, Canada.

Horson, Hugh H., 3949 15th St., N.E., Seattle 5, Washington.

House, Homer D., New York State Museum, Albany, New York.

HOWARD, FRANK L., Dept. of Botany, Rhode Island State College, Kingston, Rhode Island.

HOWARD, GRACE E., Dept. of Botany, Wellesley College, Wellesley, Mass. HOWELL, ARDEN, JR., T. B. Control Division, U.S.P.H.S., Box 3536, Duke Hospital, Durham, North Carolina.

HUGHES, STANLEY J., Imperial Mycological Institute, Ferry Lane, Kew, Surrey, England.

HUNT, N. REX, Felton, California.

HUTCHINSON, WESLEY G., Botany Dept., University of Pennsylvania, Philadelphia 4, Pa.

Hwang, Shuh-Wei, c/o K. T. Tom, Bank of China, 40 Wall St., New York 5, New York.

ILLMAN, WILLIAM IRWIN, Div. Applied Biology, National Research Council, Sussex St., Ottawa, Ontario, Canada.

IMLE, ERNEST P., U. S. Dept. of Agriculture, Rubber Station, Turrialba, Costa Rica.

IMSHAUG, HENRY A., Dept. of Botany, University of Michigan, Ann Arbor, Michigan.

JACKSON, H. A. C., 35 Campbell Ave., Montreal West, Quebec, Canada.

- JACKSON, H. S., Dept. of Botany, University of Toronto, Toronto 5, Ontario, Canada.
- JACQUES, J. E., Montreal Botanic Garden, Montreal, Canada.
- JENKINS, ANNA E., U. S. Dept. of Agriculture, Beltsville, Maryland.
- JENNINGS, OTTO E., Carnegie Museum, Pittsburgh, Pa.

Ohio.

, D. F.

Canada. Philadel-

Saska-

n, Pull-

sburgh,

, Tenn.

oit 32,

F.C.O.,

te Col-

York,

mental

gston,

ass.

Duke

Kew,

Phila-

New

ıncil,

alba,

bor,

- Jennison, Marshall W., Dept. Plant Sciences, Syracuse Univ., Syracuse 10. New York.
- JOHANSON, ALFHILD E., 730 Cranford Avenue, New York 66, N. Y.
- JOHNSON, AARON G., Bureau of Plant Industry, Beltsville, Maryland.
- JOHNSON, GEORGE T., Dept. of Biology, Mass. Inst. of Technology, Cambridge 39, Mass.
- JOHNSON, HOWARD W., Delta Branch Expt. Station, Stoneville, Miss.
- Jones, Fred R., 214 Hort. Bldg., University of Wisconsin, Madison, Wis-
- JONES, JOYCE HEDRICK, Museums Bldg., University of Michigan, Ann Arbor, Michigan.
- JONES, RICHARD C., Dept. Biology, State Teachers College, New Paltz, New York
- Jones, Walter, Plant Pathology Lab., Saanichton, British Columbia, Canada.
- JORSTAD, IVAR, Botanical Museum, Oslo, Norway.
- Josserand, Marcel, 24 Rue de la Part-Dieu, Lyons, France.
- Jump, John A., Dept. of Biology, University of Notre Dame, Notre Dame, Indiana.
- JUST, THEODOR K., Chicago Natural History Museum, Chicago 5, Illinois. KANOUSE, BESSIE B., 406 South 5th Ave., Ann Arbor, Michigan.
- KARLING, J. S., Dept. of Botany, Purdue University, W. Lafayette, Indiana. KARPLUK, Mrs. Frances E., N. Y. Skin and Cancer Unit, 330 Second Avenue, New York 3, New York.
- KEEPING, E. SILVER, Provincial Lab., University of Alberta, Edmonton, Alberta, Canada.
- Keil, Harry L., Biological Labs., Rohm & Haas Co., Bristol, Pa.
- KEITT, G. W., Hort. Bldg., University of Wisconsin, Madison, Wisconsin. KERN, FRANK D., Dept. of Botany, Pennsylvania State College, State Col-
- lege, Pa.
  KEVORKIAN, ARTHUR G., Atkins Gardens and Research Lab., Apartado 251,
- Cienfuegos, Cuba.

  Khan, Abdul H., 107 Hilgard Hall, University of California, Berkeley, California.
- KIENHOLZ, JESS R., U. S. Fruit Disease Lab., Hood River, Oregon.
- King, Thomas H., Dept. of Plant Pathology, University Farm, St. Paul,
- KOLK, LAURA A., Brooklyn College, Brooklyn, New York.
- KOPF, KENNETH, Hawaiian Pineapple Co., Honolulu 1, T. H.
- KORF, RICHARD P., Dept. Plant Path., Cornell University, Ithaca, New York.
- LAMBERT, E. B., Division Fruit and Vegetable Crops and Diseases, Plant Industry Station, Beltsville, Maryland.
- LANGE, MORTEN, Kongensgade 33, Copenhagen, Denmark.
- Ledingham, G. A., Prairie Regional Research Lab., National Research Council, Saskatoon, Sask., Canada.

LENTZ, PAUL L., Mycology and Disease Survey, Plant Industry Station, Beltsville, Maryland.

LEFEBURE, C. L., U. S. Hort. Field Station, U.S.D.A., Beltsville, Maryland. LEVINE, MICHAEL, 100 Halsted Avenue, Yonkers 4, New York.

Lewis, George M., Cornell Univ. Medical School, 121 E. 60th St., New York, N. Y.

LILLY, VIRGIL G., Science Hall, West Va. University, Morgantown, West Virginia.

LIMBER, DONALD P., 3 Mayfair Road, Morris Plains, New Jersey.

\*\*LINFORD, M. B., P.O. Box 3166, Honolulu 2, Hawaii.

Ling, Lee, Agric. Div. Food-Agric., Organization U.N., Longfellow Bldg., Connecticut Ave., Washington 6, D. C.

LOCKWOOD, LEWIS B., Northern Regional Research Lab., Bureau Agric. Chem. and Engr., Peoria, Illinois.

LOHMAN, MARION L., Botany Dept., University of Hawaii, P.O. Box 18, Honolulu 10, T. H.

LOMBARD, MRS. FRANCES F., 5301 38th Avenue, Hyattsville, Maryland. Lowe, Josiah L., College of Forestry, Syracuse University, Syracuse, New York.

Lucas, G. B., U.S.D.A. Tobacco Lab., Oxford, North Carolina.

LURIE, H. I., S. A. Inst. for Medical Res., Hospital St., Johannesburg, South Africa.

LUTJEHARMS, W. J., Dept. of Microbiology, N. J. Agr. Expt. Station, Rutgers University, New Brunswick, New Jersey.

LUTTRELL, EVERETT S., Botany Dept., University of Missouri, Columbia, Missouri.

McColloch, L. P., B.P.I., U.S.D.A., Beltsville, Maryland.

McCrea, Adelia, R.F.D. 1, Roscommon, Michigan.

McDonough, E. S., Marquette University, Dept. of Biology, 1217 W. Wisconsin Avenue, Milwaukee, Wisconsin.

McFarland, Frank T., Dept. of Botany, University of Kentucky, Lexington, Kentucky.

McGuire, J. M., Biological Div., Lilly Research Labs., Indianapolis, Indiana. McKeen, Constance G., Botany and Plant Path., Central Exper. Farm, Ottawa, Ontario, Canada.

McKenzie, Malcolm A., University of Massachusetts, Clark Hall, Amherst, Massachusetts.

MACEO, JOSEPHA V., University of Puerto Rico, Rio Piedras, P. R.

MACRAE, RUTH, Division of Botany, Central Experimental Farm, Ottawa, Canada.

\*\*Mains, E. B., University Herbarium, Univ. Museums, University of Michigan, Ann Arbor, Michigan.

MANEVAL, W. E., 305 Westmont Ave., Columbia, Missouri.

MARSHALL, RUSH P., Bartlett Tree Res. Lab., Stamford, Connecticut. MARTIN, G. W., Dept. of Botany, University of Iowa, Iowa City, Iowa.

Mason, E. W., Imperial Institute of Mycology, Ferry Lane, Kew, Surrey, England.

MASSEY, L. M., Dept. of Plant Pathology, Cornell University, Ithaca, New York.

MATTHEWS, VELMA D., Coker College, Hartsville, South Carolina.

MEINECKE, E. P., 3157 Jackson St., San Francisco 15, California.

tation,

and.

New

West

Bldg.,

Agric.

x 18,

New

burg,

ation,

mbia.

W.

xing-

liana.

arm,

Am-

awa,

y of

rrey,

New

MEYER, ESTHER, University of Illinois, College of Medicine, 1853 W. Polk St., Chicago 12, Illinois.

MIELKE, JAMES L., Forest Pathology, P.O. Box 523, Albuquerque, New Mexico.

MILLER, JULIAN H., Dept. of Plant Breeding and Plant Pathology, University of Georgia, Athens, Georgia.

MILLER, L. W., Dept. of Biological Science, Ill. State Normal University, Normal, Illinois.

MILLER, PIERRE A., 405 Hilgard Avenue, Los Angeles 24, California.

MIX, A. J., Dept. of Botany, University of Kansas, Lawrence, Kansas.

MOLINA-LLARDEN, MARIO, Finca Santa Teresa, Tucuru, Alta Verapaz, Guatemala, Central America.

Montgomery, Royal M., 57 W. 57th St., New York, New York.

Moore, George T., Missouri Botanical Garden, St. Louis, Missouri.

Moore, Morris, Barnard Skin and Cancer Hospital, Washington and Theresa Sts., St. Louis, Missouri.

Morgan-Jones, J. F., Dept. of Biology, Mount Allison University, Sackville, N. B., Canada.

MORROW, MARI BETZNER, Dept. of Botany and Bacteriology, University Station, Austin, Texas.

Morse, Miss E. E., Hotel Claremont, Berkeley, California.

Moss, E. H., University of Alberta, Edmonton, Canada.

MRAK, EMIL M., 339 Hilgard Hall, University of California, Berkeley, California.

MUIR, ROBERT D., Bristol Labs. Inc., Syracuse, New York.

MULLER, ALBERT S., Escuela de Agricultura Nacional, Villa Nueva, Barcena, Guatemala, C. A.

MUNDKUR, B. B., Imperial Agricultural Research Institute, New Delhi, India.

Nannfeldt, Jan A., Uppsala Univ. Inst., Systematisk Botanik, Uppsala, Sweden.

Nauss, R. W., 1303 York Avenue, Apt. 2H, New York 21, New York. Nickerson, Walter J., c/o Dr. G. A. Edwards, Dept. of Biology, Tufts College, Medford, Mass.

NIEDERHAUSER, J. S., c/o Rockefeller Foundation, Calle Viena 26, Mexico D. F., Mexico.

Nobles, Mildred K., Div. of Botany, Central Experimental Farm, Ottawa, Canada.

NORDIN, V. J., Dominion Lab. of Forest Pathology, Dept. Botany, Univ. of Toronto, Toronto, Canada.

NYLAND, GEORGE, Dept. of Plant Path., Univ. Calif., Davis, California.

Ochoa, A. Gonzalez, Laboratorio de Micologia, Inst. de Salubridad-Y, Enfermedades Trop., Mexico D. F., Mexico.

OLIVE, LINDSAY, Dept. of Botany, L. S. U., Baton Rouge, Louisiana.

ORJUELA, JUAN N., Apartado Nal. 2645, Bogota, Colombia, South America.

ORONOZ, MANUEL RUIZ, Calle de Reynosa 56, Tacubaya, Mexico D. F.,

Mexico.

ORTON, C. R., West Va. University, Morgantown, West Virginia.

OVERHOLTS, MRS. L. O., 111 La Crosse St., Edgewood, Pittsburgh 18, Pa. Pady, S. M., Dept. of Botany, McGill University, Montreal, Quebec, Canada. Page, Robert M., School of Biological Sciences, Stanford University, California.

PARKER, B. W., Dept. of Biology, Lehigh University, Bethlehem, Penn.
PARKER, CHARLES S., Dept. of Botany, Howard University, Washington,
D. C.

PARKS, HAROLD E., Spruce Grove, Trinidad, California.

PAULSEL, DONALD Q., Cosmic Ray Project, Bedford A. A. Field, Bedford, Mass.

PAYETTE, ALBERT, St. Anne-de-la-Pocatiere, Quebec, Canada.

Pearson, A. A., Nutcombe House, Hindhead Road, Hindhead, Surrey, England.

Pease, Dorothy, Brooklyn College, Biology Dept., Bedford and Ave. H, Brooklyn 10, New York. Pelletier, Réal L., Department of Plant Pathology, University of Wis-

consin, Madison, Wisconsin.

\*\*Pereira-Filho, Manuel Jose, Caix 658, Porto Alegre, Estado do Rio Grande do Sul, Brazil, South America.

PETERSEN, GRACE A., 8511 110th St., Richmond Hill, New York. PETTY, MILTON, JR., Lederle Labs., Inc., Pearl River, New York.

PLAKIDAS, A. G., Dept. of Botany, Louisiana State University, Baton Rouge, La.

PLUNKETT, ORDA A., U.C.L.A., 405 Hilgard Ave., Los Angeles, California. POTTRAS, A. W., Botany Dept., University of Illinois, Urbana, Illinois.

POLLACK, FLORA G., 7209 Garland Avenue, Takoma Park, Maryland.

POMERLEAU, RENE, Dept. of Lands and Forests, Quebec City, Canada.

POOLE, ROBT. F., Clemson College, Clemson, South Carolina.

POPE, SETH, Wyeth Inc., 111 N. High St., West Chester, Pennsylvania.

PORTER, CHARLES L., 924 N. Main St., West Lafayette, Indiana.

POTTER, VICTOR C., 221 N. Elm St., Ithaca, Michigan.

POUND, ROSCOE, Langdell Hall, Harvard University, Cambridge, Mass.

POVAH, ALFRED H., 143 Elmhurst Avenue, Detroit 3, Michigan.

Prasad, Nirankar, Div. of Plant Pathology, University of California, Berkeley, California.

PRESLEY, JOHN T., Dept. of Plant Path., Agricultural Expt. Station, State College, Miss.

Preston, Dudley A., Dept. Botany and Plant Pathology, Oklahoma A. & M. College, Stillwater, Oklahoma.

PRINCE, ALTON E., Dept. of Botany, University of Maine, Orono, Maine. RADER, WILLIAM E., Shell Agric. Lab., Box 1531, Modesto, California.

RANZONI, F. V., Dept. of Botany, University of California, Berkeley 4, California.

RAPER, KENNETH B., Northern Regional Research Lab., U.S.D.A., Peoria, Illinois.

REBELL, G. C., Dept. Dermatology and Syphilol., School of Medicine, University of Pennsylvania, Philadelphia, Pa.

RAY, W. W., Dept. of Botany, Univ. of Nebraska, Lincoln 8, Nebraska. Reed, George M., 25 S. Linwood Avenue, Pittsburgh 5, Pa.

8, Pa. Canada. iversity,

enn. nington,

Bedford,

Surrey, ve. H,

f Wis-

do Rio

Baton fornia.

ania.

Berke-State

S.

A. & ine.

ley 4,

Uni-

REESE, E. T., Biological Lab., Phila. Q. M. Depot, Philadelphia 45, Pa. Reilly, Joseph J., Div. Plant Path., St. Paul Campus, Univ. Minn., St. Paul 1, Minnesota.

REISCHER, HELEN S., 251 W. 68th St., New York, New York.

REYNOLDS, ERNEST S., N. Campus, Rm. 209, University Station, Coral Gables 34, Florida.

RHOADS, ARTHUR S., William F. Clapp Laboratories, Inc., Duxbury, Mass. ROANE, MARTHA K., Box 585, Blacksburg, Virginia.

ROBERTS, JOHN M., Bureau of Laboratories, Michigan Dept. of Health, Lansing 4, Michigan.

ROBERTS, JOHN W., 116D, Park Ave. S., Winter Park, Florida.

RODRIGUEZ, V. ING. JOSE, Oficina de Estudios Especiales, Direc. Gral. de Agricultura, San Jacinto, D. F. Mexico.

ROGERS, DONALD P., New York Botanical Gardens, Bronx Park, New York, New York.

ROGERSON, CLARK T., Dept. Plant Pathology, Plant Science Bldg., Ithaca, New York.

Rosen, H. R., Arkansas Agr. Expt. Station, Fayetteville, Arkansas.

ROTH, LEWIS F., Dept. of Botany, Oregon State College, Corvallis, Oregon. ROUTIEN, JOHN B., Chas. Pfizer Co., 11 Bartlett St., Brooklyn 6, New York. RUDOLPH, BERT A., Univ. of California, Deciduous Fruit Station, 2675 Stevens Creek Road, San Jose, California.

RUEHLE, GEO. D., Subtropical Expt. Station, Homestead, Florida. RUSDEN, PHILIP L., 206 Federal Bldg., Central Square, Cambridge, Mass.

RYAN, SISTER MARY HILAIRE, Rosary College, River Forest, Illinois.
SAHA, JAGADISH CHANDRA, Osborn Bot. Lab., Yale Univ., New Haven, Connecticut.

SAKSENA, RAM KUMAR, Dept. of Botany, University of Allahabad, Allahabad, India.

Salvin, Samuel B., Div. Infect. Diseases, Nat. Inst. of Health, Bethesda 14, Maryland.

SANFORD, MRS. AUGUSTA LEUCHS, Round Hill Road, Greenwich, Connecticut. SANTACROCE, NUNZIA G., 525 Caswell Avenue, Staten Island 2, New York. SAUNDERS, LEILA J., 114 Charles St., East Lansing, Michigan.

SAVILE, DOUGLAS B. O., Div. of Botany and Plant Path., Central Experimental Farm, Ottawa, Ontario, Canada.

Scully, Francis J., 904 Medical Arts Bldg., Hot Springs, Arkansas. Seaver, Fred J., N. Y. Botanical Garden, Bronx Park, New York, New York. Seeler, Edgar V., Farlow Herbarium, Divinity Avenue, Cambridge, Mass. Seligmann, Edward B., Jr., 816 Chestnut Rd., Apt. B, East Lansing, Michigan.

SHANOR, LELAND, Botany Department, University of Illinois, Urbana, Illinois.
SHAW, CHARLES G., Dept. Plant Pathology, Washington State College,
Pullman, Washington.

SHEAR, C. L., 2253 N. Madison St., Arlington, Virginia.

SHEMA, BERNARD F., 316 E. McKinley St., Appleton, Wisconsin.

SHERBAKOFF, C. D., Agric. Experiment Station, University of Tennessee, Knoxville, Tennessee.

SHIMP, MRS. HELEN N., Walnut, Illinois.

Shoup, C. S., Dept. Biology, Box 77, Vanderbilt University, Nashville, Tennessee.

SILVEIRA, VERLANDE D., Rua Visconde de Piraja, Ipanema, Rio de Janeiro, Brazil, S. A.

SIMMONS, EMORY G., Dept. of Botany, University of Michigan, Ann Arbor, Michigan.

SIMSON, FRANK W., 427 Currie Road, Durban, Natal, South Africa.

SINDEN, JAMES W., Dept. of Botany, Pennsylvania State College, State College, Pennsylvania.

SINGER, ROLF, Seccion Criptogamia, Instituto Miguel Lillo, Calle Miguel Lillo, Tucuman, Argentina, S. A.

SKINNER, CHAS. E., Dept. of Bacteriology and Public Health, Washington State College, Pullman, Washington.

Skolko, A. J., Dominion Lab. Forest Pathology, University of New Brunswick, Fredericton, N. B., Canada.

SLAGHT, HELEN B., 4801 Connecticut Ave., N.W., Washington 8, D. C.

SLEETH, BAILEY, 1461 Seventh Ave., Yuma, Arizona.

SLIPP, A. W., School of Forestry, University of Idaho, Moscow, Idaho.
SMART, ROBERT F., Dept. of Biology, University of Richmond, Richmond, Virginia.

SMITH, ALEXANDER H., University Herbarium, Museums Building, University of Michigan, Ann Arbor, Mich.

SNELL, WALTER H., Botany Dept., Brown University, Providence, R. I.SNYDER, W. C., 107 Hilgard Hall, University of California, Berkeley, California.

SOLHEIM, W. G., Botany Dept., University of Wyoming, Laramie, Wyoming. Sparrow, F. K., Dept. of Botany, University of Michigan, Ann Arbor, Michigan.

Sprague, Roderick, Dept. Plant Pathology, Washington State College, Pullman, Washington.

Springer, Martha E., Dept. of Botany, Willamette University, Salem, Oregon.

STAKEMAN, E. C., University Farm, St. Paul, Minnesota.

STESSEL, GEO. J., Dept. Plant Path., Univ. Wisc., Madison, Wisconsin.

STEVENS, NEIL E., Dept. of Horticulture, University of Illinois, Urbana, Illinois.

STEVENS, RUSSELL B., Dept. of Botany, University of Tennessee, Knoxville, Tennessee.

STEVENSON, JOHN A., 4113 Emery Pl., N.W., Washington 16, D. C. STEWART, IRENE M., 3198 Travers Ave., Hollyburn, B. C., Canada.

STIFLER, MRS. CLOYD BURNLEY, 315 16th St., Bradenton, Florida.

STILLINGER, C. R., 418 College Avenue, Moscow, Idaho.

STOUFFER, DAVID J., Safford, Arizona.

Stowell, Elwell A., Dept. of Botany, University of Wisconsin, Milwaukee Extension Div., Milwaukee 3, Wisconsin.

STRONG, MRS. MIRIAM C., 1213 North Walnut St., Lansing, Michigan. STUNTZ, D. E., Botany Dept., University of Washington, Seattle, Wash. SUMSTINE, DAVID R., King Edward Apt., Pittsburgh, Pennsylvania.

Sussman, Alfred S., Biological Labs., Divinity Ave., Cambridge 38, Mass.

Vashville,

Janeiro,

n Arbor,

e, State

Miguel shington

Bruns-

D. C.

laho. chmond, Univer-

R. I. y, Cali-

yoming. Arbor,

College, Salem,

. Irbana,

Knox-

aukee

n.

sh. Mass. SUTTON, W. D., 313 Wharncliffe Road N., London, Ontario, Canada.

SWARTZ, DELBERT, Box 93, University Station, Fayetteville, Arkansas.

SWARTZ, JACOB H., 422 Beacon St., Boston, Massachusetts.

SWEET, HERMAN R., 155 Lincoln Road, Medford, Massachusetts. TEHON, LEO R., Illinois State Natural History Survey, Urbana, Illinois.

TERRIER, CHARLES A., Station Federale d'Essais Viticoles et Arboricoles, Lausanne, Switzerland.

TERVET, IAN, Camp Detrick, Maryland.

THIRUMALACHAR, M. J., 20th V. Main Road, Malleswaran, Bangalore, India. Thom, Charles, Port Jefferson, L. I., New York.

THOMPSON, GEORGE EDWARD, Dept. of Plant Pathology, College of Agriculture, University of Georgia, Athens, Georgia.

THURSTON, H. W., Dept. of Botany, Pa. State College, State College, Pennsylvania.

TIFFANY, Mrs. Lois HATTERY, Botany Hall, Iowa State College, Ames, Iowa.

TIFFNEY, WESLEY N., 35 Cottage St., Sharon, Massachusetts. TORO, RAFAEL A., 1674 Irving St., N.W., Washington, D. C.

TORREY, G. S., R.F.D. 2, Storrs, Connecticut.

TRESNER, H. D., 615 North Lake St., Madison, Wisconsin.

TUCKER, C. M., 200 Lefevre Hall, University of Missouri, Columbia, Missouri.

Tullis, E. C., P.O. Box 2967, Beaumont, Texas.

UMANZIO, CARL B., Kirksville College of Osteopathy & Surgery, Kirksville, Missouri.

VAN DER LAAN, H. F., 4909 Wharton St., Galveston, Texas.

Vanterpool, T. C., Dept. of Biology, University of Saskatchewan, Saskatoon, Saskatchewan.

VERMILLION, MONROE T., Earich Road, R. D. 3, Athens, Ohio.

WAGENER, W. W., 720 Appraiser's Bldg., San Francisco, California.

WAKSMAN, SELMAN A., New Jersey Agric. Expt. Station, New Brunswick, New Jersey.

WALKER, A. R., Dept. of Botany, University of Western Ontario, London, Ontario, Canada.

WALKER, LEVA B., 1325 N. 38th St., Lincoln 3, Nebraska.

WALTER, JAMES M., 2410 16th Ave., Bradenton, Florida.

WALTERS, MAURICE B., 14556 Superior Road, Cleveland Heights, Ohio.

WARREN, JOHN R., Botany Dept., Duke University, Durham, North Carolina. WATERMAN, ALMA M., Marsh Hall, 360 Prospect St., New Haven, Connecticut.

WATERSTON, J. M., Moor Plantation, Ibadan, Nigeria, B. W. Africa.

WATSON, MRS. ALICE J., Mycology and Disease Survey, Plant Industry Station, Beltsville, Maryland.

Weaver, Elmer A., 7135 Boyer St., Philadelphia 19, Pa.

Weber, George G., Experiment Station, University of Florida, Gainesville, Florida.

WEGRICH, O. G., Commercial Solvents Corp., Terre Haute, Indiana. WEHMEYER, LEWIS E., 381 Orchard Hill Drive, Ann Arbor, Michigan.

WEIDMAN, FRED D., Medical Lab., University of Pennsylvania, 368 Hamilton Walk, Philadelphia, Pa. Weiss, Freeman G., Plant Industry Station, Beltsville, Maryland.

Welch, D. S., N. Y. State College of Agriculture, Cornell University, Ithaca, New York.

Wellman, F. L., Office of Foreign Agric. Relations, U.S.D.A., Washington,

WERNHAM, C. C., Dept. of Botany, Pennsylvania State College, State College, Pa.

West, Erdman, 101 Newell Hall, University of Florida, Gainesville, Florida.

\*\*Weston, Wm. H., Jr., Biological Labs., Harvard University, Divinity Ave., Cambridge, Mass.

WHEELER, WILLIS H., 3800 18th St., N., Arlington, Virginia.

WHIFFEN, ALMA, Research Labs., Upjohn Co., Kalamazoo, Michigan.

White, W. Lawrence, Harvard Univ., 20 Divinity Ave., Cambridge 38, Massachusetts.

WHITEHEAD, MARVIN D., Dept. Plant Pathology, University of Wisconsin, Madison, Wisconsin.

WICKERHAM, L. J., 219 Vincent St., Peoria Heights, Illinois.

WILHELM, STEPHAN, Div. Plant Pathology, University of California, Berkeley 4, California.

WILSON, C. M., 121 Poplar Ave., Norfolk, Virginia.

WILSON, MALCOLM, Dept. of Mycology, Royal Botanic Garden, Edinburgh 4, Scotland.

WOLF, FREDERICK A., Dept. of Botany, Duke University, Durham, N. C. WOLF, FRED T., Botany Dept., Vanderbilt University, Nashville, Tenn.

WOLFE, LLOYD R., 476 Park Ave., Glencoe, Illinois.

WOLFF, EMILY T., 139 N. Highland Road, Springfield, Pennsylvania. WOOD, JOHN L., Penn. State Forest School, Mont Alto, Pennsylvania. WOOLLIAMS, G. E., Lab. Plant Pathology, Summerland, B. C., Canada.

YARWOOD, CECIL E., Div. of Plant Pathology, University of California, Berkeley, California.

YUSEF, HASSAN M., N. Y. Botanical Gardens, Bronx Park, New York 58, New York.

ZABEL, ROBERT A., N. Y. State College of Forestry, Syracuse University, Syracuse, New York.

ZELLER, S. M., Oregon Agric. Expt. Station, Oregon State College, Corvallis, Oregon.

ZIEGLER, ARTHUR W., 1606 Meridian Rd., Tallahassee, Florida.

ZIFFER, JACK, Division of Biochemistry, London School of Hygiene and Tropical Medicine, Keppel St. (Gower St.), W.C. 1, London, England. ZUCK, ROBERT K., Dept. of Botany, Drew University, Madison, New Jersey. ZUNDEL, GEORGE L., 207 Agric. Education Bldg., State College, Pa.

#### DECEASED

BEARDSLEE, HENRY CURTIS, Painesville, Ohio. LONG, WILLIAM HENRY, Albuquerque, New Mexico. REA, PAUL M., Santa Barbara, California. ZELLER, S. M., Corvallis, Oregon.

# **MYCOLOGIA**

iversity, hington,

, State

Florida.

Divinity

dge 38,

Berke-

nburgh I. C.

fornia, ork 58, ersity, evallis,

e and gland. ersey.

FINANCIAL STATEMENT

(July 1, 1947-June 30, 1948)

Unexpended reserve, July 1, 1947		\$ 4,924.05
Mycological Society (members' subscriptions)	\$1,864.00	
Subscriptions	3,477.80	
Sale of back sets (vol. 25 and later)	824.35	
Payments for excess paging	388.52	
Special funds:		
Sale of back sets (vols. 1-24) and index	415.80	
Interest on endowment	583.00	
	\$ 998.80	
Total receipts '		\$ 7,553.47
Total on hand  Cost of printing and distribution:  Printing, binding, mailing 6 issues  Engraving	\$4,787.44	\$12,477.52
	\$5,997.17	
Replacing exhausted issues	524.82	
Miscellaneous office expenses		
Total cost		\$ 6,796.83
Balance		\$ 5,680.69
Transferred to endowment fund		\$ 1,000.00
Unexpended reserve, June 30, 1948		\$ 4,680.69
Endowment fund		14,000.00
Total on hand		\$18,680.69

### INDEX TO VOLUME XL\*

#### INDEX TO AUTHORS AND TITLES

Benham, Rhoda Williams. Effect of nutrition on growth and morphology of the Dermatophytes. 1. Development of macroconidia in *Trichophyton rubrum* 232-240, 2 figures

Bonner, John Tyler. A study of the temperature and humidity requirements of Aspergillus niger 728-738, 3 figures

Brian, Percy W., Hemming, H. G. and Jefferys, E. G. Production of antibiotics by species of *Myrothecium* 363-368

Brodie, Harold J. Variation in fruit bodies of Cyathus stercoreus produced in culture 614-626, 3 figures

Burlingham, Gertrude S. Henry Curtis Beardslee 505-506, portrait Cain, Roy Franklin. Myriogonium, a new genus among simplified Asco-

mycetes 158-167, 56 figures

Cash, Edith Katherine. Six new Indian Discomycetes 724-727

Chiu. Wei Fan. The Boletes of Yunnan 199-231

Chupp, Charles (with Thirumalachar, Mandayam Jeersannidhi). Notes on some Cercosporae of India 352-362

Covert, Scott W. A cellulose-degrading Gliocladium 386-387

Cummins, George Baker (with Thirumalachar, Mandayam Jeersanni-dhi). Status of the rust genera Allopuccinia, Leucotelium, Edythea, and Ypsilospora 417-422, 5 figures

Darby, Richard T. (with White, W. Lawrence, Stechert, Gladys M., and Sanderson, Kathryn). Assay of cellulolytic activity of molds isolated from fabrics and related items exposed in the tropics 34-84, 3 figures

Dearden, E. R. (with Jackson, Herbert Spencer). Martensella corticii Thaxter and its distribution 168-176, 11 figures

DeLamater, Edward D. Basic fuchsin as a nuclear stain for fungi 423-429, 1 figure; The nuclear cytology of Blastomyces dermatitidis 430-444, 5 figures

Drechsler, Charles. Three Zoöpagaceae that subsist by capturing soil Amoebae 85-105, 4 figures

Fennell, Dorothy I. (with Raper, Kenneth B.). New species of Penicillium 507-546, 11 figures

Fischer, George W. Graminicolous Smuts of North America. Century I 502-503

Harris, Hubert A. Heterothallic antibiosis in Mucor racemosus 347-351
Hazen, Elizabeth L. (with Schatz, Albert). The distribution of soil microorganisms antagonistic to fungi pathogenic for man 461-477, 3 figures

<sup>\*</sup> This index prepared by Aurèle La Rocque.

- Hemming, H. G. (with Brian, P. W., and Jefferys, E. G.). Production of antibiotics by species of Myrothecium 363-368
- Jackson, Herbert Spencer and Dearden, E. R. Martensella corticii
  Thaxter and its distribution 168-176, 11 figures
- Jefferys, E. G. (with Brian, P. W. and Hemming, H. G.). Production of antibiotics by species of Myrothecium 363-368
- Jenkins, Anna Eliza. The Swingle Sphaceloma hand lens and early records of the pathogene of citrus scab 630-632, 1 figure
- Jillson, Otis F. and Nickerson, Walter John. Mutual antagonism between pathogenic fungi. Inhibition of dimorphism in Candida albicans 369-385, 2 figures
- Kanouse, Bessie Bernice. The genus *Plectania* and its segregates in North America 482-497, 12 figures
- Karling, John Sidney. Keratinophilic Chytrids. III. Rhizophydium nodulosum sp. nov. 328-335, 20 figures
- Kern, Frank Dunn. Lee Oras Overholts 1-5, portrait

rphol-

dia in

quire-

ion of

s pro-

Asco-

Notes

sanni-

ythea,

s M.,

molds

34-84,

orticii

423-

430-

g soil

Peni-

Cen-

7-351

soil

-477,

ait

- Lange, Morten. Two species of Coprinus with notes on their cultural characters 739-747, 4 figures
- Lefebvre, Camille Leon and Sherwin, Helen S. An undescribed species of Helminthosporium on Sudan grass and Sorghum 708-716,
- Long, William H. and Stouffer, David J. Studies in the Gasteromycetes: XVI. The Geastraceae of the southwestern United States 547-585, 32 figures
- Lurie, H. I. A common antigenic factor in different species of Sporotrichum 106-113
- Mains, Edwin B. Entomogenous fungi 402-416, 4 figures; Two unusual fungi from Glacier National Park, Montana 717-723, 1 figure
- Martin, George Willard. Hemitrichia vs. Hyporhamma 125-126 Morse, Elizabeth Eaton. Variation in Montagnites arenarius (DC.) 255-
- 261, 19 figures
- Murrill, William Alphonso. Russula tricolor Marr. 635
- Nickerson, Walter John (with Jillson, Otis F.). Mutual antagonism between pathogenic fungi. Inhibition of dimorphism in Candida albicans 369-385, 2 figures
- Nyland, George. Preliminary observations on the morphology and cytology of an undescribed heterobasidiomycete from Washington state 478-482, 1 figure
- Olive, Lindsay Shepherd. Taxonomic notes on Louisiana Fungi—I. 6-20, 3 figures; Taxonomic notes on Louisiana Fungi—II. Tremellales 586-604, 3 figures
- Pady, Stuart M. Teliospore discharge in Puccinia tumidipes Peck 21-33, 19 figures
- Pease, Dorothy. Sphaeronaemella fimicola (Marchal) emend.: some characteristics in culture 114-124, 19 figures
- Rader, William E. Helminthosporium portulação a new pathogen of Portulação oleração L. 342-346, 1 figure
- Raper, Kenneth B. and Fennell, Dorothy I. New species of Penicillium 507-546, 11 figures

- Roberts, John Maurice. Developmental studies of two species of Nowakowskiella Schroeter: N. ramosa Butler and N. profusa Karling 127-157, 2 figures
- Rogers, Donald Philip. The meaning of Article 57 of the International Rules 241-254; On Zygodesmus 633-635; Proposals for the amendment of Art. 64 of the International Rules of Botanical Nomenclature 635-637
- Routien, John B. Hyphal proliferation through clamp-formation in Polyporus cinnabarinus Fr. 194-198, 1 figure
- Sanderson, Kathryn (with White, W. Lawrence, Darby, Richard T., and Stechert, Gladys M.). Assay of cellulolytic activity of molds isolated from fabrics and related items exposed in the tropics 34-84, 3 figures
- Schatz, Albert and Hazen, Elizabeth L. The distribution of soil microorganisms antagonistic to fungi pathogenic for man 461-477, 3 figures
- Seaver, Fred Jay. Humaria and Lachnea 498-499
- Sharp, Aaron J. Some fungi common to the highlands of Mexico and Guatemala and Eastern United States 499-502
- Shear, Cornelius Lott. Mycological Notes. IX 748-759
- Sherwin, Helen S. (with Lefebvre, Camille Leon). An undescribed species of Helminthosporium on Sudan grass and Sorghum 708-716, 2 figures
- Singer, Rolf. New genera of Fungi-IV 262-264
- and Smith, Alexander Hanchett. Emendations to our proposals concerning the nomenclature of the gill fungi 627-629
- (with Smith, Alexander Hanchett). Notes on the genus Cystoderma 454-460, 1 figure
- Smith, Alexander Hanchett. Notice of foray 268, Paul Marshall Rea 388; Henry C. Beardslee 389; Studies in the dark-spored Agarics 669-707, 93 figures
- and Singer, Rolf. Notes on the genus Cystoderma 454-460, 1 figure
   (with Singer, Rolf). Emendations to our proposals concerning the nomenclature of the gill fungi 627-629
- Sparrow, Frederick Kroeber. Soil phycomycetes from Bikini, Eniwetok, Rongerik and Rongelap atolls 445-453, 19 figures
- Sprague, Roderick. Some leaf spot fungi on western Gramineae—II 177-193, 2 figures; Some leaf spot fungi on western Gramineae—III 295-313, 2 figures
- Stechert, Gladys M. (with White, W. Lawrence, Darby, Richard T., and Sanderson, Kathryn). Assay of cellulolytic activity of molds isolated from fabrics and related items exposed in the tropics 34-84, 3 figures
- Stevenson, John Albert. John Lewis Sheldon 387-388; William H. Long 389
- Stouffer, David J. (with Long, William H.). Studies in the Gasteromycetes: XVI. The Geastraceae of the southwestern United States 547-585, 32 figures
- Tehon, Leo Roy. Notes on the parasitic fungi of Illinois 314-327, 15 figures

Thirumalachar, Mandayam Jeersannidhi and Chupp, Charles. Notes on some Cercosporae of India 352-362

and Cummins, George Baker. Status of the rust genera Allopuccinia, Leucotelium, Edythea, and Ypsilospora 417-422, 5 figures

Warren, John R. An undescribed species of Papulospora parasitic on Rhizoctonia solani Kuhn 391-401, 3 figures

Wehmeyer, Lewis Edgar. The developmental pattern within the genus Pleospora Rab. 269-294, 56 figures

White, W. Lawrence, Darby, Richard T., Stechert, Gladys M., and Sanderson, Kathryn. Assay of cellulolytic activity of molds isolated from fabrics and related items exposed in the tropics 34-84, 3 figures

Wilson, J. K. An examination of the exudate and juice of certain fungi found in their native environment 605-613, 1 figure

Zeller, Sanford Myron. (Book Review) Smith, Alexander H., North American Species of Mycena 265-268; Notes on certain Gasteromycetes, including two new orders 639-668

Ziegler, Arthur William. A new species of Achlya 336-341, 10 figures

## INDEX TO GENERA AND SPECIES

New names, the final members of new combinations, and the page where the former description appears, are in bold face type. Mistakes in the spelling of scientific names encountered in the text are corrected in the index.

Absidia capillata 44, 76

of No-

Karling

national

amend-Nomen-

tion in

ard T.,

molds

s 34-84.

micro-

, 3 fig-

ico and

scribed

08-716,

posals

Cysto-

Il Rea

garics

figure

ng the

Eniwe-

ae—II

e—III

., and

ls iso-

34-84,

m H.

stero-

States

27, 15

Acanthorhynchus vaccinii 754 Acaulopage 87, 92, 95, 98, 103; baculispora 89, 91, 93, 94, 103– 104; ceratospora, 95, 98; cercospora 95, 98; gyrinodes 97, 98,

105; ischnospora 92; macrospora 92, 93; stenospora 92, 93 Achlya 336, 338, 340; abortiva 338; crenulata 336–338, 340–341; glomerata 338, 341; radiosa 338;

recurva 338, 341 Acladium 44, 80 Acremonium 44, 80

Actinomyces 476 Actinopelte 318, 320; dryina 318, 319, 326

Actinothyrium gloeosporioides 318 Acurtis 628, 629

Aecidiolum 242 Aecidium 241, 242, 246, 251

Aegerita 249; candida 249 Agaricus 251, 637; arenarius 256; campestris 693; melleus 250; polytrichophilus 703, 707; radiosus 256; rubriceps 263, 264;

trechisporus 628 Agrocybe 670

Aleurodiscus 168, 249; amorphus 168; candidus 500; subgiganteus 249

Allopuccinia 417; diluta 417, 422 Alternaria 80; citri 736; tenuis 44 Amanita caesaraea 501; chlori-nosma 501; muscaria 501; rubescens 501; verna 501

Amanitopsis vaginata var. fu 501; vaginata var. livida 501 fulva Anellaria 689; separata 689

Ansatospora acerina 326 Antennaria 754-756; ericophila

754-757; salisburgensis 758 Antennularia 754, 756, 759; engleriana 757; ericophila 756-758; rhododendri 757; salisburgensis

756, 757 Anthopeziza 483, 491, 474-497; floccosa 491, 492, 495, 496; Win-teri 491, 495

Anthurus 645 Aphanomyces laevis 332

Apiocarpella 298, 299; macrospora 298, 299, 313

Apiosporella macrospora 298

Apostemidium 483

Arcangeliella 640, 643; Beccarii 664; magna 643; volvaria 663 Areolaria columnaris 653 Armillaria 250; mellea 250 Armillariella 250; mellea 250 Arrhytidia involuta 602

Arthrosporium 44, 81 Ascochyta 182, 297, 301, 305; agropyrina 297; avenae 297; elymi 182; graminicola 182, 301; graminicola zur. brachypodii 190; hordei 297, 298, 313; phleina 181, 182; utahensis 295, 297, 301, 313

Ascochytella avenae 297 Ascochytula 297

Aserophallus 645
Aspergillus 54, 77, 84, 469; candida 77; Chevalieri 44, 45, 77; Chevalieri var. intermedius 45, 77; clavatus 77; Fischeri 45, 77, 164, 166; flavipes 45, 77; flavus 45, 46, 78; fumigatus 46, 77; glaucus 77; japonicus 46; luchuensis 41, 46, 47, 78, 84; nidulans 47, 77; niger 47–49, 78, 84, 728–730, 732–734, 736–738; ochraceus 49, 78; Oryzae 50, 78; repens 50, 77; Sydowi 50, 51, 78; Tamarii 52, 78; terreus 52, 78; unguis 52, 53, 77, 469; ustus 53, 77; ustus var. laevis 53, 77; versicolor 54, 74, 78; Wentii 54, 78

Asperisporium 322; Acori 321, 327; Caricae 322 Aspropaxillus 629; giganteus 629

Astraeus 547, 580, 582, 654; hygrometricus 548, 580, 585, 654; pteridis 654; stellatus 580
Astrosporina 628
Atylospora fuliginosa 697
Auricularia auricularis 587

Batarrea 660
Blastocladia 132, 134, 156, 157
Blastocladial simplex 134
Blastomyces 425, 427, 431, 434, 435, 441, 442; braziliensis 373, 379; dermatitidis 373, 379, 384, 424, 425, 430, 431, 433, 436–444, 461
Blastospora 422
Boletellus 231; ananas 231
Boletinus 199; kunmingensis 199; pictus 201; punctatipes 200; punctatipes 200; punctatipes 200; punctatipes 200; punctatipes 200; punctatipes 200; punctatipes 201; 202; aereus 211; albidus 223; albofarinaceus 209;

foletus 201, 202; aereus 211; albidus 223; albofarinaceus 209; atroviolaceus 203; brunneissimus 228; candidus 223; castaneus 202; Cheoi 215, 216, 217; chrys-

enteron 214; communis 214; communis var. mutatis 214; crocipodius 210; decorus 216; edulis 227; eximius 228; felleus 207; flavidus 202; flavus 203; Frostii 221. 222: fuscescens 210: Gertrudiae 227; granulatus 206; instabilis 215; javanicus 207, 208; Kauffmanii 226; Iuteoporus 210; luteus 203; magnificus 221; mi-nutus 201; nigropunctatus 214; ornatipes 225, 226; paludosus 213, 215, 218; placidus 205; por-phyrosporus 206; pseudostrobilomyces 212; punctato-fumosus 207; punctilifer 216; puniceus 217; Queletii 221, 222; radi-cans 223; Ravenelii 212; recedens 219; regius 224; retipes 225, 226; rimosellus 226; rimosus 210; roseolus 207, 208; rubropunctus 219; rugosellus 219; san-guineus 224, 225; Satanas 220; sinicus 220, 222; speciosus 224; splendidus 223; subpaludosus 213, 218; subsplendidus 222; sylvestris 218; Taianus 220; tes-selatus 210; thibetanus 204; tomentipes 214; umbrinellus 209; umbrinus 209, 229; unicolor 218; velatus 208; violaceo-fuscus 210; virens 206; vitellinus 223; yunnanensis 213, 217 Botryodiplodia Theobromae 54, 82

Botrytis allii 364-366 Bovistella atrobrunnea 649, 650 Bovistoides simplex 583, 585 Brachysporium 54, 80 Broomeia 647 Broomeiaceae 647, 648 Bubakia crotonis 10 Bulbopodium 681

Caeoma 246; pusillum 309
Calbovista 648
Calocera cornea 602
Calostoma 657; cinnabarina 501; microsporum 655
Calvarula 644
Campanularis anomalus 685
Candida 374, 377, 380, 382, 383, 430; albicans 369–375, 377, 381, 383, 385, 425, 461, 463, 465, 467–470, 472–475; albicans var. stellatoidea 425; albicans var. tropicalis 425; guillermondi 425; Krusei 425; parakrusei 425
Cantharellus carbonarius 628; ci-

barius 501; floccosus 501 Catenaria 136, 137, 156 214; d; cro-; edus 207; Frostii Ger-6; in-

s 210; l; mis 214; idosus ; porrobilomosus niceus radi-

recees 225, mosus rubro-; sans 220; s 224; dosus l; syl-; tes-

4; to-209; r 218; s 210; yun-54, 82

550

501;

383, -381, 467stel-

steltropi-425;

pusillu Chamoni

Catenochytridium carolinianum 133, 137, 156; laterale 133, 135, 144, 156

Catenomyces 136; persicinus 136, 137, 156

Catharinia 273 Centrospora acerina 324 Cephalosporium 55, 80, 122

Cephalothecium 55; roseum 55, 398, 401

Ceratostomella 114, 117, 124; adiposum 124; fimbriata 123; paradoxa 123; ulmi 476

Cercoseptoria domingensis 361 Cercospora 4, 192, 324, 327, 352, 356; Achyranthina 352; Achyranthis 352; Apii 353; Aspleni 353; bangalorensis 353; Calotropidis 353; canescens 354; Cannabis 354; Capparidis 354; carbonacea 354; Caricae 322; centrostachydis 352; Clauseniae 354; Cocculi 355; Coffeicola 355; consimilis 355; daizu 360; difformis 322, 325, 327; exotica 358; fusimaculans 355; garbini-ana 360; granuliformis 325, 327; Hibiscina 355; Holarrhenae 355; Ii 325; Ixorae 356; Jasminicola 356; kaki 356; Lettsomiae 356; lilacina 325; macrospora 324; Mali 357; Melochiae 362; minima 357; murina 325; mysorensis 357; Nelumbonis 358; Nerii-Indici 358; Nymphaeacea 358; Panici 355; Panici-milacei 355; Panacis 358; Paramignyae 358; penicillata vor. Apii 353; petila 359; Piricola 357; Pouzolziae 358; Punicae 359; Rubi 360; septorioides 360; Shoreae 360; sojina 360; Sorghi 360; Sorghi var. Maydis 360; sororiae 323, 324, 325, 327; subsessilis 361; Tectoniae 361; Tinosporae 361; Vignicaulis 354; Violae 324–327; Violae-sylvaticae 325; Violae-tricoloris 324, 326, 327; Wal-theriae 361; Wrightiae 362; Zizyphi 362

zyphi 362 Cercosporella 178, 192; subulata 177, 178

177, 178 Cercosporina Cannabis 354; kikuckii 354 Chaconia 421; baphiae 421

Chaetomella 55 Chaetomium 82, 83; funicolum 42, 55, 56, 82-84; indicum 56, 82;

pusillum 755–757 Chamonixia 642, 667 Cheilymenia calvescens 725 Chitonia 263

Chrysocelis ascotela 421, 422 Chrysomyxa 718 Chytridium 332-335; cocconeidis

Chytridium 332-335; cocconeidis 332; Rhizophydii 333, 335; versatile 332; versatile var acaulis 332 Chytriomyces 157, 331, 332, 334, 335; hyalinus 332; stellatus 332

Citromyces 544; minutus 511 Cladochytrium 127; hyalinum 135, 137, 156; replicatum 130, 134– 137, 139, 142, 144, 147, 153, 156; tenue 136

Cladosporium herbarum 56, 57, 80, 181, 307

Clathrella 644, 645 Clathrogaster 660-664; Beccarii 664; vulvarius 664

Clathrospora 272, 289, 292–294 Clathrus 644, 645; cibarius 645 Claudopus 628, 629

Claudopus 628, 629 Clavaria 637 Clethria 644

Clitocybe 627, 629; laccata 501

Clitopilus 628 Clypeus 628

Coccidioides immitis 430, 461 Cochliobolus 708, 716; heterostrophus 710

Cochlonema 103; euryblastum 88 Coemansia 174–176; aciculifera 170 Coilomyces Schweinitzii 649 Coleosporium 718, 719, 723

Coleroa 752; chaetomium 752; salisburgensis 757, 758; straussii 755, 757

Colletotrichum 712; gloesporides 476

Collybia 265; conigena 197; platyphylla 501; radicata 489; tenuipes 268

Coniophora 634, 746 Coniothyrium 57, 82

Copelandia 685
Coprinus 255, 669, 670, 674-676, 679, 680, 684, 739, 746, 747; angulatus 670, 706; asterophorus 672, 706; atramentarius 674, 679; Brassicae 672, 673, 706; calvptratus 672, 706; cinchonensis 672, 706; cothurnatus 739-743, 747; dilectus 740; domesticus 677; ebulbosus 673, 679, 682-684, 706; ephemerus 672, 674, 682, 684; hexagonosporus 673, 706; insignis 674, 706; jalopensis 676, 706; Jonesii 676, 706; lagopus 207. rotundisporus 676; laniger 677, 706; mexicanus 677, 706; micaceus 681; mycelio

cephalus 673, 678, 741, 742, 744, 745, 747; narcoticus 739; niveus 680, 740; phaeosporus 678; plicatilis 682; pseudoradiatus 678, 706; pulchrifolius 678, 706; quadrifidus 673, 679, 682, 683, 706; radians 676, 677, 679; roseotinctus 740; rotundisporus 679, 706; semilanatus 680, 706, 740; Seymourii 680, 706; silvaticus 681, 706; Spraguei 682, 706; sterquilinus 672; subpurpureus 684, 706; variegatus 679, 682, 706; vermiculifer 744

Corditubera 640; microspora 640;

Staudtii 640

Cordyceps 4, 405, 407, 408, 410-412, 415; australis 411; bicephala 411; curculionum 402, 405, 411, 416; Huberiana 411; Humberti 402, 404; necator 411; oxycephala 404; paludosa 415; peltata 407, 409, 410, 415, 416; proliferans 411; Puiggarii 402-404, 416; sphaecocephala 402; sphecocephala 402, 404; (Torrubia) sphaecophila 402; sphecophila 402; sphecophila 402; subsessilis 405, 406, 416

Corticium 248, 633; alutaceum 169, 172; bombycinum 168; confluens 169; lacteum 172; microsporum 158, 168; pruinatum 248; radio-

sum 169, 172, 173; varians 197 Cortinarius 262, 681 Corynites 646; Ravenelii 646

Crinipellis 268

Crinodendron 636 Crucibulum vulgare 501

Cryptococcus neoformans 461, 463–465, 467–470, 472–475

Cunninghamella echinulata 58, 76,

Curvularia 80, 712; brachyspora 58, 84; falcata 58; geniculata 712; lunata 58

Cyathus 614, 624; stercoreus 614-626; stercoreus form Lesueurii 621, 624; stercoreus var. minor 622, 624; stercoreus form rufipes 624; vernicosus 621

Cyclomyces 4 Cylindrochytrium johnstonii 157 Cylindrosporium calamagrostidis 181, 305, 307, 308, 313; glyceriae 302

Cystoderma 454, 705; amianthinum 457, 458; amianthinum var. sublongisporum 460; amianthinum f. typicum 455; carcharias f. album 459; cinnabarinum 459; granulosum var. adnatifolium 459; granulosum var. occidentale 458; granulosum var. typicum f. typicum 459; Gruberianum 458; haematites 458; paradoxum 454, 455; subpurpureum 457, 458

Dacrymyces 593; abietinus 598, 599; abietnus var. **triseptata 598**, 604; deliquescens 593, 599; Ellisii 599; minor 593, 599; palmatus 599

Dacryomitra ceracea 601, 602; stipitata 593, 601, 602, 604 Dactylaria thaumasia 96, 105 Daedalea 4; quercina 501

Dasyscyphella 724; indica 724 Deconica bulbosa 696; crobula 262; pyrispora 703; subviscida 704

Dendrogaster 643 Dendryphium 16; comosum 16; costaricense 16

Desmaturus 645 Desmella 420; aneimiae 420; berberidis 420; quitensis 420; superficialis 420, 422; tenella 420, 422

ficialis 420, 422; tenella 420, 422 Diaporthe 241, 246; Ailanthi 241; alnea 241; detrusa 241 Dichotomophthora portulacae 342 Didymosphaeria inaequalis 751 Dimerosporiopsis englerianus 757

Diplocystis 647 Dipodascus 164 Dispira cornuta 400

Dissoderma 454 Dycticia 644

Eccilia 628, 629 Edythea 417, 418, 420; berberidis 418, 420; quitensis 418, 420; tenella 418, 420, 422

Elasmomyces alpinus 643; Mattirolanus 659; Rodwayi 642; sessile 659

Endochytrium operculatum 130, 133–135, 137, 139, 144, 147, 148, 156

Endomycopsis albicans 365 Endostigme 749; chlorospora 749; cinerascens 750, 751; crataegi 749; ditricha 749; fraxini 749; inaequalis 749-751; pirina 749; tremulae 749

Entoloma 628, 629 Entophlyctis aurea 449 Eocronartium muscicola 586, 603 Epiclinum Cumminsii 322

Erinella 724; corticola 724 Eriosphaeria 759; salisburgensis 757, 759

Eucordyceps 410 Eucystoderma 454 lentale cum f. n 458; m 454,

s 598, ta 598, 9; El-1; pal-12; sti-

24 la 262; 704 n 16;

5

; bersuper-20, 422 ii 241;

ne 342 51 s 757

peridis (0; te-

2; ses-130, 7, 148,

a 749; ataegi 749; 749;

5, 603

gensis

Eumycena 266, 267 Euryachora 302 Euzygodesmus 634 Exidia 598; nucleata 593; recisa 593; spiculosa 593

Favolus 4; rhipidium 500 Fimetaria 611 Fistulina hepatica 608–610 Fomes 4; annosus 501; Feei 501; pinicola 501, 502; roseus 501; senex 501 Fusarium 60, 79, 631; culmorum

usarium 60, 79, 631; culmorum 401; javanicum 58; lateritium 58; moniliforme 58, 387, 388; oxysporum cubense 476; roseum 59; Scirpi var. longipes 59; semitectum 59; semitectum var. majus 59; Solani 59, 60

Fusicladium 749; Peucedani 322

Gasterella 639
Gasterellaceae 639
Gasterellopsis 639

Gastrosporium 660, 661, 665; simplex 665 Gautieria 642, 660, 661, 666, 667;

morchelliformis 667 Gautieriaceae 661, 666 Gautieriales 661, 666

Gautieriales 601, 666

Geaster 547, 548, 554, 555, 562, 563, 569, 581; arenarius 548, 563, 569, 571; Bryantii 548, 550, 552, 555; caespitosus 649; campestris 548, 556, 557, 559, 562; coronatus 548, 566; Drummondii 548, 549, 555, 559; elegans 548, 552, 553, 555; ellipticis 554; fimbriatus 549, 576, 578; floriformis 548, 577, 578; floriformis 548, 577, 578; Hariotii 548, 552, 555; Hieronymi 548, 567, 569; hygrometricus var. giganteus 580; juniperinus 566, 569; lignicola 649; limbatus 548, 569, 570, 573, 585; limbatus var. pacificus 569; mammosus 548, 560, 562; minimus 548, 565, 566, 569; papyraceus 649; pectinatus 548, 550, 552, 555; pluriosteus 548, 550, 552, 555; pluriosteus 548, 550, 552, 574; saccatus form minor 548, 574, 578; saccatus form simili 548, 559, 562, 566; tomentosus 548, 564, 569; triplex 548, 572, 573; vulgaris 580; xerophilus 548, 554, 555; xylogenus 548, 562, 563

Geasteroides 579; texensis 579 Geastrum 547, 581, 582; fibrillosum 580; hygrometricum 580; hygrometricum var. giganteum 654; mirabile 649; Schweinitzii 649 Geopetalum 628; carbonarium 628;

Geopetalum 628; carbonarium 628 petaloides 628

Geopyxis coccinea 484; floccosa 491; hesperidea 485; occidentalis 485

Gibbera 752, 753, 759; salisburgensis 757, 758; straussii 757, 758; vaccinii 752, 759

Glaziella aurantiaca 445 Gliocladium 60, 79, 386; fimbriatum 79; roseum 60, 79, 386 Gliomastix convoluta 61, 81 Glocosporium 319, 321; affine 319

Gloeosporium 319, 321; affine 319 Gloeotulasnella pinicola 602 Glutinipes 266

Guepinia spathularia 602 Gymnoascus 524; Reesii 544 Gymnoglossum 643, 666; cambodgense 643; candidum 643; connectens 643; elasmomycetoides 643; foetidum 643; globosum 643; majus 643; megasporum 643; olivaceum 643; radiatum

643; stipitatum 643; utriculatum 643 Gymnomyces 642; vesiculosus 641, 642

Gymnosporangium 242; clavipes 30; inconspicuum 30 Gyroceras 17 Gyrodon 205, 229

Helicobasidium 169, 249; purpureum 249, 254 Helicogonium 164, 165; Jacksonii

158, 160, 168 Helicoceras 17

Helicogloea 590, 603; Lagerheimi 587, 603; **sebacinoidea 588**, 603 Helicoma 16, 17

Helicomina 16, 17; caperoniae 15, 16, 17, 18, 20

Helminthosporium 181, 344, 355, 708, 711, 712, 714, 716, 752; cookei 711; maydis 710; portulacae 342, 343, 344, 345; sativum 398, 401; sorghi 711, 712, 714, 715; sorghicola 713, 714, 715; 16; sudanensis 712, 714

Helotium 165, 166; albidum 165; citrinulum 165; erraticum 165; gemmarum 165; midlandensis 165; scutula 165; scutula var. fucatum 165

Helvella 268

Hemiarcyria 125; serpula 125

Hemitrichia 125, 126; karstenii 125; serpula 125, 126; Vesparium 125

Hendersonia 181; crastophila 190; culmicola f. minor 190; simplex

Heterosporium phlei 181

Histoplasma capsulatum 373, 425, 430, 461

Hoehneliogaster 640 Hohenbuehelia 628

Humaria 498, 499, madi 725; pall 724, 725; ahpallidisetosa 725: plumbea 726

Humarina 498, 724; plumbeo-atra 726; purpurea 726; umbrina 726, 727; waterstonii 727; zizyphi 727

Humicola 62, 79, 84

Hydnangium 640, 642; carneum 640; nigrescens 641; purpureum 641; vesiculosum 641

Hydnum 637

Hygrophilus 264 Hygrophorus acutoconicus 268 Hymenochaete 248; tomentosa 248 Hymenogaster 658, 666

411; australiensis Hymenostilbe

411, 412; formicarum 411; melanopoda 411; melanopus 411 Hymenostomum 490; microstomum 490; squarrosum 490

Hypochanum Hypocrea 408, 410 Hypodendrum 4 Hypomyces 408

Hyporhamma 125, 126; reticulatum 125 Hypoxylon malleolus 500

Ileodictyon 644, 645 Illosporium malifoliorum 388 Inocybe 268, 628

Ionomidotis 483 Irpex farinaceus 500, 502

Isaria melanopus 411 Itajahya 647; galericulata 647

Karlingia rosea 332, 453 Kickxella 174, 176 Krombholzia crocipodia 210; luteopora 210 Kunkelia nitens 7

Kupsura 646; sphaerocephala 646

Lachnea 498, 499, 725; coccinea 484; mirabilis 487 Lactaria 506

Lactarius indigo 501, 502; Peckii 501

Laestadia 751 Leccinum nigrescens 210 Lenzites 4; Berkleyi 501; betulina 501, 502; saepiaria 501 Lepiota 459; cinnabarina 459; pro-

cera 501 Lepista 627

Leptographium 124 Leptonia 628, 629

Leptosphaeria 270, 273, 278, 300; arenaria 186; Yerbae 278, 293 Leptostroma Avenae 315

Leptostromella Solani 317, 327 Leptothyriella Liquidambaris 318 Leptothyrium 315; anthelminthici 314; Avenae 315, 327; Cheno-podii 315; fimbriatum 315; Smilacis 317

Leucogaster 640, 668 Leucopaxillus 629

Leucophleps 668; candida 668 Leucophizon 665; nidificum 665, 666

Leucotelium 417, 418; cerasi 418,

Lichthemia Regnieri 62, 76 Linderomyces 262

Lophodermium arundinaceum 307, 311

Lycogala 668; mexicanum 651 Lycogalopsis 647

Lycoperdellales 668 Lycoperdellon 667, 668; minutum 668; Torrendii 668

Lycoperdon 565, 651, 652; acuminatum 651, 652; albidum 648; albinum 648; Curtisii 651, 652 Lysurus 388, 645; Gardneri 646;

Mokusin 645, 646; pusillus 646

Macowania 642 MacOwanites 642; agaricinus 642; alpinus 643; magnus 643

Macrometrula 263, 264; rubriceps 264 Macrophoma 300; phlei 301; sporoboli 300, 301, 313

Maravalia 418, 422 Martensella 173-176; Corticii 168-175; pectinata 175

Mastigosporium 308 Megachytrium Westonii 136, 139 Melampsora 241, 247, 717-719, 723;

Kusanoi 719 Melanospora 114 Memnoniella 83; echinata 62, 63,

Mesopsora 719; Hypericorum 717-719

Metarrhizium 83, 368; Anisopliae 63; brunneum 63; glutinosum 36, 81, 84, 363, 367, 368, 477 Michenera 249; artocreas 249

betulina 59; pro-

78, 300; 3, 293

327 ris 318 ninthici Cheno-5; Smi-

668 m 665, asi 418.

ım 307,

551

inutum acumin 648; 652

652 ri 646; us 646

; spo-

ii 168-

6, 139 9, 723;

52, 63, n 717–

sopliae um 36, Microsphaera 8; alni 8, 10; alni var. cinnamomi 8, 9, 19; alni var. lonicerae 8

Microsporum 80, 234; audouini 234, 235; canis 234; gypseum 63, 234, 461

254, 401 Microstoma 483, 486, 488, 490, 491, 494; hiemale 487, 488, 496, 497; protracta 486, 489–491, 496 Milesia 250, 251, 253, 254; Polygoni 250; Polypodii 250

Milesina 250 Mimema 418

Mitrula 719, 720; gracilis 717, 719, 721, 722; muscicola 719, 720, 722; phalloides 268

Monascus 515 Monilia 108, 112; albicans 385; candicans 248

Monoblepharella 157; Taylorii 134 Montagnea 255, 256; arenaria 256; candollei 256

Montagnites 255, 256; arenarius 255, 256, 258–260; arenarius var. texensis 261; candollei 256; pallasii 256; radiosus 256; tenuis 256

Morchella 268 Morganella 650, 652; mexicana 650 Mortierella tuberosa 170 Mucor 373; fumosus 63, 76; hiemalis 347; Mucedo 347, 605; racemosus 347-350; sexualis 347

Mutinus 646
Mycena 265–268; acuto-conica 268; albiscolor 267; albidula 268; albissima 267; alnicola var. odora 267; arenaria 267; Atkinsoniana 267; bisphaerigera 268; Brownii 267; cayugaensis 267; cheboyganensis 267; cineraria 267; cylindrospora 267; delicatella 268; epipterygia var. lignicola 267; fagicola 267; filigrinella 267; galericulata 267; Gaultheri 267; griseoviridis 267; griseoviridis 267; griseoviridis 267; griseoviridis 267; griseoviridis 267; Kauffmaniana 267; Kuehneriana 267; kauffmaniana 267; Kuehneriana 267; kauffmaniana 268; litoralis 267; McMurphyi 268; misera 268; olida var. americana 267; pallida 268; papillata 268; paucilamellata 267; plumbea var. robusta 267; pseudoclavicularis 267; pseudoclavicularis 267; pseudoinclinata 267; pusillissima 268; Rickeni 267; rubrotincta 267; semivestipes 268; serotina 268; sertulosa 267;

subcana 267; subconcolor 267; subfusca 267; subimmaculata 268; subvestita 268; Swartzii 268; tenuiceps 267; tenuicula 267, 268; thujina 267; translucentipes 268; turbinata 268; umbrina 267; vestita 268

Mycenastraceae 648 Mycenastrum 648 Mycenella 266, 267 Mycopharus 645

Mycosphaerella 120; citrullina 120; tulasnei 307, 308 Myriogonium 158, 165, 167, 176;

Myriogonium 158, 165, 167, 176; Odontiae 159, 161, 163, 164, 169 Myrothecium 81, 363, 364, 368; inundatum 64, 81, 363–367; roridum 63, 81, 363–368; verrucaria 36, 64, 81, 84, 363–367

Myxosporium 64 Myrica 635 Myriostoma 547, 582; coliforme 583, 585

Naematoloma 669, 670, 692, 693, 703, 706; campestre 692, 706; dispersum 694; elongatipes 693; humidicola 693, 706; petasiforme 694, 706

Naucoria humidicola 693 Nematoctonus haptocladus 194 Neopaxillus 262; echinosporus 262 Nephrochytrium 153, 157 Nidulariopsis 656, 657; iowensis

657; melanocarpa 656, 657 Niesslia 754, 756; chaetomium 756; exilis 756; pusilla 756

exilis 756; pusilla 756 Nolanea 628, 629 Nowakowskiella 127, 130, 132, 136, 139, 144, 145, 147, 153, 154, 157; delica 130, 140, 152; elegans 130; elongata 130, 151, 152; granulata 132; hemisphaerospora 130, 135, 146, 154; macrospora 132, 148, 157; profusa 127–130, 137, 139, 147–154, 156, 157; ramosa 127–133, 135–137, 139, 140, 142, 144–148, 150–155

Octaviania 640 Odontia 159; sudans 158, 159, 167, 169

Oidium 248, 254, 633, 634; effusum 249; Morgani 249; pulveraceum 249; tomentosum 248

Olpidium 450, 452; Rhizophlyctidis 449, 452, 453; Uredinis 450 Omphalia 265, 268; vestita 268 Omphalopsis 268

Oospora 80; lactis 64

Ophiobolus 716; graminis 398; heterostrophus 708, 716 Ophiocordyceps 410; peltata 410 Ophiostoma 114, 117, 124 Orbilia 483

Ormathodium 14, 16; ambrosiae 14, 15, 20; costaricense 16; styracis 14

Otidea 483

Otidella fulgens 268

Ovularia 749; baldingerae 310; haplospora 309; holci-lanati 310; hordei 301, 309, 310; lolii 310; pulchella 309, 310; pulche'la var. agropyri 310; pusilla 309, 310

Paecilomyces 80; varioti 64 Panaeolus 669, 684-686, 689, 692; anomalus 684, 706; castaneifolia **685**, 706; digressus 696; foenisecii 685; **fontinalis 690**, 706; fraxinophilus 686, 706; papili-onaceus 690; reticulatus 686, 706; retirugis 687; rufus 688, 706; semiovatus 688, 689; solidipes 688, 689, 706; subbalteatus 687, 688, 690; variabilis 689, 690, 706; venenosus 690, 706. Papulospora 391-394, 396-398, 400;

stoveri 394-396, 399, 400, 401 Patella 409, 725; gilva 726 Patouillardinia 597; cinerea 597 Paxillus 262, 627, 629; lepista 627 Pellicularia 248, 254, 634; lem-

Penicularia 246, 254, 054; lembospora 249; pruinata 248
Penicillium 69, 70, 78, 79, 84, 507, 515, 524, 527, 535, 543, 544; aculeatum 535, 537, 538, 543, 546; baarnense 527; bacillosporum 521; Biourgeianum 64, 65; bre-feldianum 514, 515; brevi-compactum 477; capsulatum 528, 529, 530, 543, 545; chrysogenum 65, 79, 738; citrinum 65, 79; corylophylum 66; cyaneum 66; diversum 539, 540-543, 546; didiversum 539, 540-543, 546; diversum var. aureum 540, 541, 543, 546; ehrlichii 515, 544; expansum 477; funiculosum 535, 538, 543; gladioli 477; helicum 515, 516, 518, 520, 543, 545; herquei 66, 538, 544; implicatum 66; javanicum 511, 515, 543; lapidosum 524, 525, 527, 543, 545; lavendulum 530, 531-533, 543, 545; levitum 511, 513-515, 543, 545; levitum 511, 513-515, 543, 545; levitum 66; luteum 66, 67, 78, 79, 518, 521, 524, 543; majuscu-79, 518, 521, 524, 543; majuscu-lum 67; mangini 527; minutum 511; multicolor 67; notatum 5, 67, 79; pallidum 532, 533, 543;

parvum 508, 509, 511, 515, 527, 543, 545; patulum 477; piceum 533, 534, 535, 543, 546; piscarium 67; purpurogenum 67, 79, 538, 543; roseo-purpureum 68; ro-543; roseo-purpureum 68; ro-tundum 518, 519, 543, 545; rubescens 533; rugulosum 79, 538; rugulosum var. atricolum 68; sanguineum, 68; spinulosum Steckii 68; striatum 521, 522, 524, 527, 543, 545; sulfureum 68: Sweicickii 68; tardum 68, 541–543; thomii 527, 543; vermiculatum 518, 520; verruculosum 538; virido-albus 69; Westlingi 69; wortmanni 69, 521, 524, 544 Peniophora 168, 169, 249, 634; Aegerita 249; candida 249

Peridermium 242, 251 Peronospora 6, 7; fragariae 7; parasitica 6; potentillae 7, 18, 20; rubi 7; sparsa 7

Pestalotia 70, 71 Peziza 637; coccinea 482, 484; cruciata 486, 491; Dudleyi 483, 484; floccosa 482, 483, 491, 495; hesperidea 485; hiemalis 487; melastoma 482; mirabilis 487, 490; occidentalis 482, 485; protracta 482, 486, 491; saccardiana 727; sepiatra 727; sepiatra 727; violacea 726

Phaeoseptoria 190, 192, 312; airae 191; calamagrostidis 307; festucae 190, 191, 306, 307; poae 190, 191

Pharus 645

Phleospora 180, 181, 192; graminivorum 181: idahoensis 180, 181, 192

Phlyctidium mycetophagum 335 Phlyctorhiza variabilis 332, 335

Pholiota 4; aurivella 254 Phoma 71, 82, 241; Ailanthi 241; alnea 241; detrusa 241 Phragmidium 30; rubi-idaei 478 Phyllachora 241; Trifolii 241, 246

Phylloporus 262 Phyllosticta 300, 301, 320; destructiva var. Evonymi 321; healdii 182, 183, 192; panici 183; rhoi-cola 320; rogleri 183, 301; Smi-lacis 317; sorghina 183

Phymatotrichum omnivorum 398 Physalacria 603

Physocladia 139 Physopella fici 10

Phytophthora erythroseptica 347 Pilacre 412

Pilobolus crystallinus 605; longi-pes 605, 606

15, 527, piceum scarium 79, 538, 68; ro-3, 545; um 79. ricolum ulosum 1, 522, eum 68: 8, 541ermicuulosum estlingi

7; par-18, 20;

24, 544 634;

4: cru-3, 484; 5: hesmela-', 490; otracta a 727; 27; vi-

airae festuie 190.

grami-180, 335 335

i 241; 478 1, 246

struc ealdii rhoi-Smi-398

347

ongi-

Piricularia parasitica 308 Pirostoma Nyssae 318 Plasmopara gonolobi 6 Platygloea Peniophorae 168 Plectania 482, 483, 485, 489, 490, 494; coccinea 484, 486, 496, 497; floccosa 483, 491; hiemalis 487; occidentalis 485, 486, 496; protracta 482, 483, 487, 490, 491, 496, 497

Pleosphaerulina 273 Pleospora 269-271, 273-276, 278, 281, 288-290; abbreviata 291; abscondita 279, 293; Alismatis 282, 283, 293; alpestris 289; amplispora 289, 294; Anthyllidis 289; Anthyllidis var. Aconiti 282, 293; Armeriae 286; atromaculans 283; Balsamorrhizae 286; Berberidis 283, 294; Boldoae 280, 293; calvescens 278; carpinicola 283; chlamydospora 287; coloradensis 286; Compositarum 294; cybospora 294; delicatula 278; diaporthoides 280, 293; dura 279, 293; Ephedrae 289; Forsteri 278; fuegiana 278; Galii 279, 293; helvetica 290; Henningsiana 290, 294; herbarum 272, 273, 276, 277, 283-287, 289-294; herbarum var. occidentalis 284, 285, 291, 294; Herniariae 285, 294; hispida 290; infectoria 280, 293; infec-toria var. nigriseda 283; infectoria var. sacchari 293; interme-dia 279; Inulae-candidae 291; Jaapiana 286, 294; Juglandis 283; Jaapiana 200, 294, Jugandus 290, kansensis 283, 293; laricina 290, 294; laxa 287; longispora 287, 294; Maireana 294; maritima 286, 294; media 284, 285, 289— 291; mollis 279, 293; montana 287, 294; multiseptata 289, 294; Niessleana 279, 293; njegusensis 294; obligasca 289; oblongata 279, 293, 294; obtusa f, fibrincola 294; orbicularis 283; oligostachyae 280, 293; oligotricha 283; papaveracea 278; parvula 278; pellita 278, 293; phyllophyla 291; pleosphaeroides 282, 293; pul-chra 287; pustulans 290; Rhodotypi 289; richthophensis 280, 282, 283, 293; rubicola 289, 294; rubicunda 282, 293; Shepherdiae 283; socialis 280; stenospora 286, 294; Thuemeniana 278, 283, 290, 294; Thurgoviana 282, 293; tomentosa 282; vagans 278, 293; vulgaris 272, 273, 276, 277, 279, 280, 292, 294, 201, 202, 282-284, 291-293

Pleurotus ostreatus 501, 502 Polycephalomyces 414; formosus 414, 415; paludosus 413, 414–416; ramosus 414, 415

Polychytrium 133, 156; stromaphilum 134

Polyphagus Euglenae 147, 156, 157 Polyporus 4; abietinus 500; adustus 500, 501; arcularius 500; australis 500, 502; cinnabarinus 194, 196, 197; cinnamomeus 500, 501; cristatus 500, 502; Curtisii 500; cuticularis 500, 502; dichrous 500; distortus 500; dryadeus 606-612; gilvus 500, 501; glomeratus 3; hirsutus 500; liceritus 500; liceritus 500; liceritus 600, luciricus 613; heidus noides 500; longipes 612; lucidus 500; perennis 500; pinsitis 500, 502; pocula 500; rhipidium 500; sanguineus 501, 502; Schweinitzii 501; semipileatus 500, 502; sulphureus 501, 606, 609,

tulipiferus 501; versicolor 501 Polythrincium 241; Trifolii 241,

Pompholyx 656 Poria 3-5 Prospodium 421 Protogaster 639 Protophallus 643, 644

Protubera 643, 644; brunnea 644; jamaicensis 644; Maracuja 643,

Proventuria 748 Psathyra cinchonensis 700; lateritia 698; mammillata 700

Psathyrella 264, 268; disseminata 250; petasiformis 694

Pseudocolus 645 Pseudocoprinus disseminatus 250 Pseudodiscosia 193; avenae 180 Pseudomonas coronafaciens v. at-

ropurpureum 296 Pseudomycena 266

Psilocybe 669, 670, 693, 694, 702, 703; atrobrunnea 701; atrorufa 698; bulbosa 694, 706; caespitosa 696, 706; californica 696, 706; castaneifolia 685; castanella 696, 706; coprophila 697, 702; fuligi-706; coprophila 697, 702; funga-nosa 697, 706; graveolens 697, 707; lateritia 698, 707; latispora 698, 707; mammillata 700, 707; modesta 700, 707; nigrella 700, 707; pallidispora 701, 707; panae-oliformis 701, 707; phyllogena 700, 702, 703, 707; plutonia 702, 707; pyrispora 703, 707; silvatica 698; subviscida 696, 704, 707; tomentosa 705, 707; vialis 705, Puccinia 30, 249, 417, 418; cerasi 422; donidiae 30; globosipes 30; graminis 244; graviae 30; helianthi 254; obliqua 6; sorghi 253; tumidipes 21, 23, 27, 30-32; Caricae 322; pullulans 71

Pyrenochaeta 72, 82, 313; elymi 299; exosporioides 299; graminis 299; leptospora 299; luzulae 299;

oryzae 299

Pyrenophora 274, 307, 716, 748; hyphasmatis 278; moravica 283; notarisii 748; phaeocomoides 283; rosae 748

Pyronema confluens 164 Pythium 85, 156, 157, 447; debary-anum 332; helicoides 95; irregulare 99; ultimum 87

Radiigera cinnamomea 652 Ramularia 11, 310; pulchella 309; pusilla 309; sidae 9, 11, 19 Ramulaspera 310; holci-lanati 310 Rhacophyllus 250; lilacinus 250 Beurmanni Rhinocladium equi 112; equinum 107, 108 Rhizidiomyces bivellatus 129, 130,

157 Rhizoctonia 249, 401; Crocorum 249, 254; solani 391-401, 477 Rhizomorpha 250; subcorticalis

250

Rhizophlyctis 447, 449, 450, 452, 453; (Karlingia) rosea 449, 453 Rhizophidium 328, 329; coronum 156; gibbosum 330; keratinophilum 329, 332, 335; marshallense 450, 452, 453; nodulosum 328, 329-335

Rhizopodella 482 Rhizopus 347; arrhizus 72, 76 Rhodopaxillus 627 Rhodophyllus 628, 629; abortivus

628, 629 Rhodotus subpalmatus 268 Rhopalophlyctis 157

Roestelia 242, 251 Rozella 157

Russula 506, 643; adusta 364; cinerascens 506; foetens 501; magna 506; pungens 506; patriotica 635; rubescens 506; tricolor 635

Saccharomyces 425, 430 Sarcopodium 80; fuscum 72 Sarcoscypha 482, 486, 488, 489; al-pina 487, 489; coccinea 484, 486; cruciata 487, 488; floccosa 483, 491; javensis 483, 486; occidentalis 485, 486; protracta 483, 487, 496

Schizophyllum commune 501 Scleroderma 582, 656; aurantium 653; aurea 653; Bovista 653; cepa 653, 654; columnare 653; geaster 655; lycoperdoides 653, 654; pteridis 654; tenerum 653; tuberoideum 654; verrucosum 653

Sclerophoma 301 Scleroplea 273

Sclerotinia baccata 487, 496; hiemalis 487

Scolecotrichum Alatroemeriae 322; Caricae 322; punctulatum 322 Scopulariopsis 80; brevicaulis 72 Scypharia coccinea var. hiemalis 487

Sebacina 166, 194, 197, 590, 597, 598, 603; adusta 590, 595, 597, 604; cinerea 594, 604; epigaea 594, 598; Eyrei 594, 604; podlachica 598; prolifera 164; vari-septata 595, 597, 603 Secotium 666; albipes 657, 658;

erythrocephalum 658; nubigenum 658; epandus 658; Rod-wayi 642; sessile 659

Sedecula 656

Sedeculaceae 655 Selenophoma everhartii 307

Septochytrium macrosporum 134, 157; variabile 137, 156

Septogloeum 180, 181; oxysporum 181

Septomyxa 80; affinis 72 Septoria 181, 184, 185, 189, 192,

eptoria 181, 167, 160, 167, 305, 305, 306, 312; agropyrina 304, 306; alopecuri 192; andropogonis f. sporobolica 186; avenae 185, 192, 301, 302, 304–306, 303; 313; bromi var. elymina 303; calamagrostidis 305; calamagrostidis f. koeleriae 186, 192; cenchrina 188; elymi 302, 304, 313; elymi forma elymina 304; elymina 304; glycericola 183, 192, 302; infuscans 181, 189, 304; nelicae 305; microspora 184; nodorum 184, 185, 192, 301, 302, 304, 305, 313; pacifica 304; pas-serinii 184, 185, 192; phleina 182; quinqueseptata 186. 188, 192; secalis var. stipae 301; tandilen-

sis 188; triseti 182 Simblum 646; sphaerocephalum 646

Siphonaria 154, 157 Solutoparies Pythii 130 Sorataea 417, 418; amiciae 417, 418,

422

501 urantium sta 653; are 653: ides 653. rum 653: rucosum

496; hieriae 322; m 322 ulis 72 hiemalis

590, 597, 95, 597 epigaea )4; pod-64; vari-57, 658;

nubige-8; Rod-17

ım 134, sporum

89, 192, opyrina andro-36; ave-304-306, a 303: calama-6, 192; 12. 304, na 304: a 183.

89, 304; 184; 01, 302, 4; pasna 182; 192: ndilenphalum

17, 418,

Spermospora 177, 178, 180, 308; subulata 178, 192, 308, 313

Sphaceloma 630, 631 Sphaerella 751; cinerascens 750, 751; (Sphaeria) cinerascens 750; inaequalis 750, 751

Sphaeria chaetomium 756; cincinnata 753; cinerascens 750, 751; exilis 756; fimicola 122; Helvellae 114, 121

Sphaerobolus iowensis 656, 657; f. europaea 656; stellatus 656 Sphaeronaema 114, 116, 121, 124; fimicola 121, 122; Helvellae 114,

Sphaeronaemella 114; carnea 121, 122; fimicola 114, 115, 119, 121, 122; Helvellae 114, 116, 121 Sphaeropsis 72, 300

Sphenospora 420; berberidis 420; kevorkianii 421, 422 Spilosticta 749; bistortae 749; inaequalis 749, 751

Spirochaeta cytophaga 83 Spondylocladiella botrytioides 168 Sporobolomyces 480, 481 Sporophlyctis rostrata 130, 157

Sporothrix 112; Schenckii 112 Sporotrichum 72, 80, 106, 107, 111-113; asteroides 106-108, 112; Beurmanni 107, 112; Carougeaui 106; Councilmani 107; cracoviense 106; Dori 107; equi 106; Fonsecai 106; Grigsbyi 106; Jeanselmei 106, 107; Schenckii 106-108, 112, 425; Schenckii var. Beurmanni 106; Schenckii var. Councilmani 106; Schenckii var. Fioccoi 106; Schenckii var. Greconis 106

Stachybotrys 72, 83; alternans 367;

Stagonospora 190; alopecuri 190; arenaria 185, 301, 304–306, 312– 313; brachypodii 190; foliicola 306; glyceriae 184, 190; glyceria-cola 189, 190, 192; maculata 189; subseriata 189, 190; subseriata subseriata 189, 1 var. maculata 189

Staheliomyces 646; cinctus 646 Stemphylium consortiale 72 Stereum 5; ochraceo-flavum 500; rameale 500; sepium 500 Stictopatella 321; Evonymi 321;

Iridis 320, 327

Stilbella 412 Stilbum 412, 414; burmense 410, 411, 412, 416; formicarum 411, 412; hirsutum 412; ramosum 412-414, 416; vulgare 412 Streptomyces griseus 469

Strobilomyces 229; floccopus 213, 229, 230; glabriceps 229; reti-sporus 230; velutipes 213

Stropharia 670, 689

Stylopage 88, 99, 103; araea 99, 100, 102; cephalote 87; hadra 99; leiohypha 99; rhabdospora 87, 88; rhicnacre 101, 102, 105; rhynchospora 99, 102

Stypella 598 Suaeda intermedia 30

Suillus atroviolaceus 203 Syncephalastrum racemosum 73,

Teichospora 275, 289, 290 Terrostella 547, 563, 579, 585; tex-ensis 579, 585

Thallospora 11; aspera 9, 11, 12, 18 - 20

Thelephora 637; biennis 634 Thielavia 166; Sepedonium 73, 82, 165; terricola 165

Thielaviopsis paradoxa 123 Tomentella 634; biennis 634; phy-lacteris 634

Torrubia sphecocephala 402; sphecophila 402 Torrubiella 407

Torula rhododendri 757

Trametes 4; malicola 501, 502; sepium 501

Tremella fuciformis 592, 603; lutescens 590, 591; mesenterica 591; mycophaga 168; mycophaga var. obscura 593; rufobrunnea 591, 604; simplex 168

Tremellodendron merismatoides 500 Tremellogaster 660-662; surina-

mensis 663 Tremellogasteraceae 661, 662

Tremellogastrales 661 Trichaster 547, 563, 585; texensis 580

Trichia 125; clavata 125; reticulata 125; serpula 125

Trichoderma 398, 469; lignorum 397, 401, 477, 736; viride 73, 74, 79, 80, 84, 368, 477 Tricholoma 627; nudum 197; prae-

magnum 3 Trichomonascus 176; mycophagus

Trichophyton 74, 469; album 235; discoides 234, 235; gypseum 461, 463, 467, 468, 470, 472, 473, 475; interdigitale 369, 384, 385; mentagrophytes 234, 235, 369, 461; purpureum 235, 369, 370, 385, 463, 467, 468, 470, 471, 473, 475;

rubrum 232, 234–237, 239, 240, 369–379, 382, 383, 385, 463, 467, 468, 470, 471, 473, 475; schoenleini 461
Trichurus terrophilus 363
Tritirachium 75, 80; dependens 74, 80
Tylopilus 207, 209

Underwoodia columnaris 268 Uredo 241, 246, 247, 251, 252, 723; (Melampsora?) hyperici-humifusi 718; Hypericorum 717 Uromyces 249 Uropyxis amorphae 30, 32; quitensis 420

Ustilago sphaerogena 10

Venturia 748, 749, 751, 753, 755; arctostaphyli 752-754; aucupariae 749; cassandrae 752; chaetomium 748, 756; chlorospora 749; cincinnata 753, 754; compacta 753, 759; dianthi 748; dickei 748, 753; eres 748; gaultheriae 753; inaequalis 749, 751, 752; kalmiae 753; macrotrichia 748; myrtilli 753; pirina 749; pulchella 753; pyrina 749; rosae 748; rumicis 749; straussii 755–757
Vermicularia luzulae 299
Verticillium 75, 80, 122
Volvaria 264
Volvariella 264

Xanthomonas 181; translucens var. phlei-pratensis 182 Xerocomus 213, 216–218 Xeromphalina 629; campanella 629 Xylaria cubensis 500

Ypsilospora 417, 420, 421; baphiae 420, 421

Zygodesmus 633-635; fuscus 633, 634; Hypochnoides 633; ochraceus 633 Zygorhynchus Moelleri 75, 76 753; Imiae yrtilli 753; imicis

is var.

la 629

phiae

s 633, ochra-6